

DIRECTIONS FOR USE

LAWN MASTER CRABGRASS CONTROL should be distributed evenly over 2000 sq. ft. of lawn area during the early spring of the year, after the normal spring clean-up to avoid the necessity of raking or disturbing the soil surface after application. Being a pre-emergence compound LAWN MASTER must be applied prior to the germination of crabgrass which can appear any time during or after the end of May. This product will not provide control if applied after crabgrass appears. For best results apply at the rate of 10 lbs. per 1000 sq. ft. of lawn area. (This bag covers 2000 sq. ft.) It is best to use a lawn spreader. When using one of the following spreaders adjust the setting on the number indicated in the table below and apply to the lawn first in one direction and then in a crosswise direction assuring complete coverage.

SUGGESTED SPREADER SETTINGS

LAWN BEAUTY	6
SCOTT'S	5
JACKSON	4

These spreader settings are meant for a guide only. They are in no sense a guarantee of exact application in as much as age, wear of spreader, speed of operation and roughness of terrain can cause variations in rates applied. Where some spring seeding is necessary LAWN MASTER CRABGRASS CONTROL should not be applied until the newly seeded areas have exhibited greening and preferably after the first mowing. Where LAWN MASTER CRABGRASS CONTROL is applied prior to spring seeding there should be a delay of at least 60 days before seeding to assure maximum germination of the lawn grasses.

LAWN MASTER CRABGRASS CONTROL contains the herbicide Dacthal® contains no arsenicals or other harmful residual metals to effect permanent turf vigor.

LAWN MASTER CRABGRASS CONTROL is a highly effective pre-emergence material for controlling not only crabgrass but also the following annual grasses and broadleaf weeds in turf.

Common Chick Weed	Fall panicum	PURSLANE	WITCHGRASS	Green Foxtail
TEXAS MILLET	HAIRY CRABGRASS	CARPET WEED	LAMB QUARTERS	JOHNSON GRASS (from seed)

CAUTION: Do not apply LAWN MASTER CRABGRASS CONTROL on Dichondra lawns or to newly seeded areas until greening appears. Do not take internally or contaminate food or foodstuffs. Avoid contact with eyes. Wash after handling or using. Keep children and pets off treated lawn area until the product has been washed off the grass by sprinkling or rain and the grass is dry.

Do not reuse this container for any other purpose.

FEDERAL CHEMICAL COMPANY

HOME OFFICE: LOUISVILLE, KENTUCKY 40202

5-8-67
JAN 1 1968
ACCEPTED

FREIGHT SHIPPING BAG
Meeting requirements of Package 1348
APPLICABLE FREIGHT CLASSIFICATION
Guaranteed by 

NET WEIGHT 7 POUNDS

NET WEIGHT 7 POUNDS

LAWN MASTER

LAWN WEED KILLER

Ingredient Statement - (Same As Front)

FEDERAL CHEMICAL COMPANY
HOME OFFICE: LOUISVILLE, KENTUCKY 40202

DIRECTIONS FOR USE

LAWN MASTER LAWN WEED KILLER is a combination of 2,4-D Amine and 2,4,5, TP on a light weight inert carrier. Its function is to control broadleaf weeds such as:

<u>Plaintain</u>	<u>Dandelion</u>	<u>Wild Carrot</u>
<u>Buckhorn</u>	<u>Common Chickweed</u>	<u>Wild Onion</u>
<u>Thistle</u>	<u>Mouse-Eared Chickweed</u>	<u>Pig Weed</u>
<u>Rag Weed</u>	<u>Sour Grass</u>	<u>Lambs Quarters</u>
<u>White Clover</u>	<u>Clover</u>	And many others

The combination of ingredients is very effective, economical and easy to use.

LAWN MASTER LAWN WEED KILLER can be used in the Spring, Summer or Fall. However, the best results are obtained when the weeds are growing vigorously. When using this product strive to distribute it uniformly and to retain the granules on the leaves of weeds. The following suggestions will assist:

1. The contents of this bag are to cover a 2500 sq. ft. area (50'x50'.) This can best be accomplished using a spreader.
2. Apply between mowings to provide sufficient leaf surface to be treated.
3. Sprinkle lawn prior to application or apply after a light rainfall.
4. Delay application when rain is predicted.
5. Do not water lawn for 24 hours after application.
6. Avoid application on windy days to avoid drift to surrounding plants.
7. Do not treat newly seeded lawns until grasses are well established or at least until after third mowing.

SUGGESTED SPREADER SETTINGS:

Lawn Beauty	9
Scott's	6
Jackson	5

These suggestions are for your guidance, and are in no sense a guarantee of exact application. Age, condition of spreader, speed of operation and roughness of terrain can cause variations in the rates applied.

CAPTION

GENERAL SAFETY

Avoid inhalation or contamination of food stuffs. Wash hands after using. Avoid contact with skin, eyes or clothing. Avoid contamination of irrigation ditches or water used for domestic purposes. Store away from children and domestic animals.

PLANTS AND SOILS

Do not allow dust to drift onto susceptible plants such as ornamentals, vegetables, etc. Do not apply to rhododendra lawns or to stoloniferous grasses (those with underground stems) such as bent's, bermuda or St. Augustine. Do not use on clover if you desire it to remain in the lawn. Certain desirable grasses may exhibit slight browning but will recover after a short period if the product has been used as directed. Vapors from this product may cause injury to susceptible plants growing in the immediate vicinity.

FROM WEEDING & ENVIRONMENTAL

This product is toxic to fish. Keep out of lakes, streams, ponds, or public water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not store near fertilizer, seeds, insecticides or fungicides.

PREPARATION

Do not reuse container. Destroy by burning in a safe place away from water supplies. Do not burn.

FEDERAL CHEMICAL COMPANY
LOUISVILLE, KENTUCKY 40202