Mauget. IMICIDE

SYSTEMIC INSECTICIDE
IN READY TO USE CAPSULES

FOR TREE INJECTION USE FOR SEASONAL CONTROL OF CERTAIN INSECTS ON ORNAMENTAL TREES

CAUTION

FIRST AID				
 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouting possible. Call a poison control center or doctor for further treatment advise. 				
Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes Call a poison control center or doctor for treatment advice.				
 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 				
Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.				
HOT LINE NUMBER				
product container or label with you when calling a ontrol center or doctor, or going for treatment. You contact 1-800-535-5053 for emergency treatment on.				

NOTE TO PHYSICIAN

There is no specific antidote available. Treat Patient symptomatically.

Mfg. By: Town, State: EPA Reg. No.: J.J. Mauget Co. Arcadia, CA 91006

7946-16

EPA Est. No.:

7946-CA-1

Net Contents: 25 capsules plus 25 feeder tubes per cartoon. 25 capsules @ 2mL, 50 mL net, or 25 capsules @ 3mL, 75 mL net, or

25 capsules @ 4 mL, 100 mL net
Shipping box: 12 Cartons as above.

Rev. 11/26/01D

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Causes eye irritation Wash thoroughly with soap and water after handling. Avoid breathing vapors. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT:

Some Materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

APPLICATOR AND OTHER HANDLERS MUST WEAR:

- Long-sleeved shirt and long pants.
- · Shoes plus socks

100.0%

- Chemical resistant gloves, such as polyethylene or butyl rubber or neoprene rubber or viton
- · Protective eyewear

ENVIRONMENTAL HAZARDS:

This pesticide is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS:

Do not use or store near heat or open flame.

A	C	C	E	P	Ţ	E	D	
MAP 2002 Under the Pederal Insecticide,								
Pangicide, and Rodenticide Act, as amended, for the pesticide Registered under								
EPA	Re	ea 3. N		94	1/-	16		

GENERAL DIRECTIONS

Measure the tree at chest height in inches. If measuring the circumference of the tree, divide this number by six (6) to determine the number of capsules needed. If measuring the diameter of the tree, divide this number by two (2) to determine the number of capsules needed. It is preferred to apply the injector units around the tree at the root flare.

Apply the following dosage depending on tree diameter;

2 ml - 2 to 10 inches dbh

3 ml - 10 to 36 inches dbh; or

4 ml - 36 inches dbh and above

Consult the enclosed pamphlet "Directions for use and application of Mauget "Micro-Injection System" for additional instructions. Applicators shall remove capsules promptly after treatment. Trees in advanced stages of insect infestations may not respond to treatment. The health of the tree and the environmental conditions will determine the rate of uptake.

RECOMMENDED TARGET INSECTS ON ORNAMENTAL TREES

ADELGIDS

APHIDS

BLACK VINE WEEVIL LARVAE

BRONZE BIRCH BORER

COTTONWOOD LONGHORNED BORER *

ELM LEAF BEETLE

EUCALYPTUS LONGHORNED BORER

FLATHEADED BORER (including Alder Birch Borer)

JAPANESE BEETLE

LACEBUGS

LEAFHOPPERS

LEAFMINERS

MEALYBUGS

PINE TIP MOTH LARVAE

PSYLLIDS (INCLUDING LERP PSYLLID)

ROYAL PALM BUGS

SCALE INSECTS

THRIPS

WHITEFLIES

RESTRICTIONS

Do not inject trees that are less than two inches in diameter.

This product is not to be used on trees which will produce food within the year following treatment.

2/5

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.


STORAGE: Store capsules in an upright position, above 45°F, in a cool, dry place.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Dispose of empty capsules in a sanitary landfill or by incineration if approved by State and local authorities.

NOTICE OF WARRANTY

J.J. MAUGET CO. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USE WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CON-DITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.



^{*} Imicide is not registered in California for controlling this insect


DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with it's labeling and the Worker Protection Standard, 40 CFR 170. This standard contains requirements for the protection of agricultural workers on farms, forest, nurseries and greenhouses, and the handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.



CAPSULE POSITION

Furn the capsule right side up to allow the naterial to flow through the tube into the cylem and phloem tissue in the tree.

10. REMOVAL & DISPOSAL

Allow time for the tree to absorb all the liquid material. Turn the capsule upside down for a minute before removal. Applicators shall remove capsules promptly after treatment. Follow instructions on product label for capsule disposal.

11. MICRO MINI-FEEDER TUBE

For established trees with thin bark (fess than 3/8" thickness). Use a 7/64" drill bit to produce a micro-injection site.

12. MICRO MINI-INSERTION TOOL

Because the 7/64* micro-injection site is so small, it is recommended that the Micro-mini insertion tool (316-NAM) be used to prevent plugging of the feeder tube & insure a clear pathway to the cambium tissue. Be sure to place the tube with the notch up.

DIRECTIONS


for use and application of



Micro-Injection system



REFER TO PRODUCT LABEL FOR ADDITIONAL PRECAUTIONS AND LIMITATIONS ON THE USE OF THIS PRODUCT



WHAT'S THE SYSTEM?

- A)... The Manget two piece capsule.
- B)...Insert hole.
- C)...Feeder lube.
- D) . Flanged gun-sile end,
- E)... Tapered beveled end.

2. WHAT TOOLS ARE NEEDED?


- A)... Electric Drift
- B)... 11/64" Drill bit.
- C)... Plastic Mallet (Hammer).
- D)... Tape Measure
- E)... Consult product labet for specific Personal Protective Equipment (PPE) required.

3. HOW MANY CAPSULES?

Measure the tree at chest height, if measuring the circumference, divide this number by 6 (six) to determine the number of capsules needed. If measuring the diameter, divide this number by two (2) to determine the number of capsules needed.

4. PRESSURIZING CAPSULE

Place capsule on firm flat surface and compress by pressing with bottom of foot, heal of hand or under some conditions with a plastic or rubber mallet.



5. HOW DO LORILL?

Pre-Drill the injection site at a slight downward angle at the root flair (approximately 6" above ground level) using a clean 11/64" drill bit. Drill to a depth of 3/8" (into the healthy xylem lissue). For the Micro-mini feeder tube, see #'s 11 & 12. Disinfect drill bit and Micro minimsertion tool prior to use on each tree.

6. HOW DEEP?

It is important that the feeder tube be set to the proper depth into the xylem. If set too deeply, flow is restricted by blockage in the heartwood; if set too shallow, leaks may occur. The feeder tubes are "beveled" cut to allow for 1/4" +/- error.

7. COMBINING CAPSULE & TUBE

Place by hand, the feeder tube's flange end with flange notch upwards into a compressed inverted capsule. Push the end of the tube flush with the internal capsule seal.

8. PLACING UNIT IN TREE

Firmly seat the beveled end of feeder tube with the attached inverted capsule into the predrilled injection hole. Tap the top corner of the capsule directly behind the feeder tube with a plastic mallet while supporting the capsule with the other hand. This action will simultaneously seal the feeder tube into the tree while breaking the capsule seal and releasing the material into the tree.