

PM 13

7948-19

4/4

Mauget®

ABACIDE

2 ml.

See carton label and pamphlet for complete directions & precautions.

ACTIVE INGREDIENTS

ABAMECTIN

INERT INGREDIENTS

1.0%

89.0%

TOTAL 100.0%

KEEP OUT
OF REACH
OF CHILDREN

Net Contents: 2 ml.

MADE IN U.S.A. J.J. MAUGET CO. LOS ANGELES, CA 90065

2/4

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Prolonged or frequently repeated exposure may cause allergic skin reactions in some individuals. Harmful if absorbed through skin.

Personal Protective Equipment:

Wear chemical resistant gloves and protective eyewear when handling and applying the product.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops. Use for tree micro-injection only as a post-bloom application.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns or grounds.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal of micro-injection capsules. Do not reuse micro-injection capsules.

STORAGE: Store in a cool, dry place out of the reach of children. Store capsules in an upright position in closed carton. Keep out of direct sunlight when possible.

DISPOSAL: Dispose of empty capsules in a sanitary landfill or by incineration if approved by State and local authorities.

Mauget *

ABACIDE™

INTERNAL SYSTEMIC TREATMENT BY MICRO-INJECTION
FOR CONTROL OF INSECTS/MITES .
ON ORNAMENTAL TREES.

RECOMMENDED ONLY FOR USE BY PROFESSIONAL
ARBORISTS (APPLICATORS)

ACTIVE INGREDIENT:

Abamectin B₁ [A mixture of avermectins containing > 80% avermectin B_{1a} (5-O-demethyl avermectin A_{1a}) and < 20% avermectin B_{1b} (5-O-demethyl-25-de (1-methylpropyl)-25-(1-methylethyl) avermectin A_{1b})] 1.0%

INERT INGREDIENTS: 99.0%
TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

WARNING
STATEMENT OF PRACTICAL TREATMENT

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

IF SWALLOWED: Immediately give one to two glasses of water and induce vomiting by gently touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON THE SKIN: Wash with plenty of water.

IF IN EYES: Flush with plenty of water.

Get medical attention in all cases.

Recommendations for Medical Treatment for Abamectin B₁ Acute Toxicity:

Early signs of intoxication include mydriasis (dilated pupils), ataxia (unsteadiness), and muscle tremors. Toxicity following accidental ingestion of the concentrate can be minimized by inducing vomiting within one half hour of exposure.

If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels) as indicated by clinical signs, symptoms and measurements.

In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin B₁ is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that influence GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic abamectin B₁ exposure.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

MFG. BY:

TOWN, STATE:

EPA ESTABLISHMENT NO.:

EPA REGISTRATION NO.:

NET CONTENTS:

J.J. MAUGET CO.
LOS ANGELES, CA 90065

7946-CA-1

7946-19

300 units 2 ml. or 0.7 wt. oz. ea.

GENERAL

Protective eyewear and chemical handling or installing the capsule in the eyes or skin.

ABACIDE™ insecticide is intended (applicators) on ornamental trees (Leafminers, Elm Leaf Beetle, Sycamore) It can be applied in commercial and exterior plantscapes and other woody shrubs are grown.

Measure the tree at chest height circumference of the tree, divide the number of capsules needed. tree, divide this number by two (2) capsules needed. Apply the injection flare. Consult the enclosed pamphlet application of Mauget Injector Unit. Applicators shall remove capsules if phytotoxicity has been noted in small holes and discontinue use if phytotoxicity is observed.

RESTRICTIONS

Do Not inject trees that are less than 2 inches in diameter.

Do not inject trees within two weeks of chemical treatment.


Do not treat trees that are suffering from moisture or herbicide damage.

This product is not to be used on trees within the year following treatment.


NOTICE OF WARRANTY

J.J. Mauget Co. makes no warranty for any purpose or otherwise expressed or implied in this product or its uses which extend beyond the normal conditions in accord with the label.


3/4




9. CAPSULE POSITION
Turn the capsule right side up to allow the material to flow through the tube into the xylem and phloem tissue in the tree.



10. REMOVAL & DISPOSAL
Allow time for the tree to absorb all the liquid material. Turn the capsule upside down for a minute before removal. Applicators shall remove capsules promptly after treatment. Follow instructions on product label for capsule disposal.



11. MICRO MINI-FEEDER TUBE
For established trees with thin bark (less than 3/8" thickness), use a 7/64" drill bit to produce a micro-injection site.




12. MICRO MINI-INSERTION TOOL
Because the 7/64" micro-injection site is so small, it is recommended that the Micro-mini insertion tool (316-NAM) be used to prevent plugging of the feeder tube & insure a clear pathway to the cambium tissue. Be sure to place the tube with the notch up.

DIRECTIONS

Mauget

Micro-Injection system

save trees



REFER TO PRODUCT LABEL FOR ADDITIONAL PRECAUTIONS AND LIMITATIONS ON THE USE OF THIS PRODUCT