

7946-26

01/02/2009

1  
9

JAN 02 2009

Frederick T. Smith  
SciReg, Inc.  
12733 Director's Loop  
Woodbridge, VA 22192

Subject: ArborFos (EPA Reg. No 7946-26)  
Label Amendment to include sub-labels A and B  
Application Dated: 10/2/08

Dear Mr. Smith:


The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(5), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit three (3) copies of your final printed labeling before you release the product for shipment. Final printed labeling means the label or labeling of the product when distributed or sold. Clearly legible reproductions or photo reductions will be accepted for unusual labels, such as those silk-screened directly onto glass or metal containers or large bags or drum labels.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(b). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

If you have any questions contact John Fournier at 703-308-0169 or by email at: [fournier.john@epa.gov](mailto:fournier.john@epa.gov). A stamped copy of the label is enclosed for your records.

Sincerely,



Linda A. Hollis, Chief  
Biochemical Pesticides Branch  
Biopesticides and Pollution  
Prevention Division (7511P)

Enclosure

*Concurrent*

JIP  
Fournier  
08

2  
1  
9

Master label encompassing Sub-label A – For Nonresidential Use and Sub-label B – For Residential Use

**Mauget**

**ArborFos<sup>TM</sup>**

SYSTEMIC FUNGICIDE  
IN READY TO USE CAPSULES  
FOR TREE INJECTION USE FOR SEASONAL SUPPRESSION OF  
CERTAIN DISEASES

MFG. BY:	J.J. MAUGET CO.
TOWN, STATE:	Arcadia, CA 91006
EPA REGISTRATION NO:	7946-26
EPA ESTABLISHMENT NO:	7946-CA-1
E-MAIL ADDRESS:	www.Mauget.com

**ACTIVE INGREDIENT:**

Mono- and di-potassium salts of Phosphorous Acid .....45.8%

**OTHER INGREDIENTS:** ..... 54.2%

Total ..... 100.0%

Contains 5.174 lbs/gallon of the active ingredient, mono- and di-potassium salts of Phosphorous Acid. Equivalent to 3.353 lbs Phosphorous Acid/gallon.

Net Contents:

**ACCEPTED**

JAN 02 2009

Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended, for  
the pesticide registered under  
EPA Reg. No. 7946-26

**Mauget**

**ArborFos<sup>TM</sup>**

SYSTEMIC FUNGICIDE  
IN READY TO USE CAPSULES  
FOR TREE INJECTION USE FOR SEASONAL SUPPRESSION OF  
CERTAIN DISEASES

MFG. BY:	J.J. MAUGET CO.
TOWN, STATE:	Arcadia, CA 91006
EPA REGISTRATION NO:	7946-26
EPA ESTABLISHMENT NO:	7946-CA-1
E-MAIL ADDRESS:	<a href="http://www.Mauget.com">www.Mauget.com</a>

**ACTIVE INGREDIENT:**

Mono- and di-potassium salts of Phosphorous Acid .....45.8%

**OTHER INGREDIENTS:** ..... 54.2%

Total ..... 100.0%

\*Contains 5.174 lbs/gallon of the active ingredient, mono- and di-potassium salts of Phosphorous Acid. Equivalent to 3.353 lbs Phosphorous Acid/gallon.

Net Contents:

# Mauget

## ArborFos<sup>TM</sup>

SYSTEMIC FUNGICIDE  
IN READY TO USE CAPSULES  
FOR TREE INJECTION USE FOR SEASONAL SUPPRESSION OF  
CERTAIN DISEASES

MFG. BY: J.J. MAUGET CO.  
TOWN, STATE: Arcadia, CA 91006  
EPA REGISTRATION NO: 7946-26  
EPA ESTABLISHMENT NO: 7946-CA-1  
E-MAIL ADDRESS: www.Mauget.com

### ACTIVE INGREDIENT:

Mono- and di-potassium salts of Phosphorous Acid .....45.8%

OTHER INGREDIENTS: .....54.2%

Total 100.0%

Contains 5.174 lbs/gallon of the active ingredient, mono- and di-potassium salts of Phosphorous Acid. Equivalent to 3.353 lbs Phosphorous Acid/gallon.

### Net Contents:

- \_\_\_\_\_ 24 capsules plus 24 feeder tubes per carton
- \_\_\_\_\_ 24 capsules @ 0.17 fl. oz. (5 mL) ea, 4.06 fl. oz. (120 mL) net
- \_\_\_\_\_ 24 capsules @ 0.25 fl. oz. (7.5 mL) each, 6.0 fl. oz. (180 mL) net
- \_\_\_\_\_ 24 capsules @ 0.34 fl. oz. (10 mL) each, 8.12 fl. oz. (240 mL) net

Shipping box: 12 cartons as above.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### FIRST AID

<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-535-5053** for emergency treatment information.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing vapors or spray mist. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

### PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Protective eyewear.
- Long pants and long-sleeved shirt.
- Shoes plus socks.
- Waterproof gloves.

Follow manufacturer's instructions for maintaining/cleaning personal protective equipment (PPE). If no such instructions for washables, use hot water and detergent. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before drinking, eating, chewing gum, using tobacco or using the toilet.
- Remove PPE clothing immediately if the pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool place over 45°F with capsules in an upright position. Keep out of direct sunlight when possible.

**PESTICIDE DISPOSAL:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Dispose of empty capsules in a sanitary landfill or by incineration if approved by state and local authorities. If burned, stay out of smoke.

### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

### RESTRICTIONS

Do not inject trees that are less than four inches in diameter.

### GENERAL DIRECTIONS

**Important:** Preventative application is more effective than therapeutic treatment in trees showing disease symptoms. Trees in advanced stages of disease development may not respond to treatment. The health, species of the tree, and the environmental conditions will determine the rate of uptake. Infected trees will absorb the material more slowly due to the vascular plugging caused by the disease. If ArborFos is not absorbed within 24 hours, the tree is considered high risk and has a poor chance of survival. For all rates of application, see #3 below. Thoroughly irrigate trees/plants prior to or at the same time of injection. Test for phytotoxicity prior to regular use.

5  
9

PLANT	DISEASE
ORNAMENTAL & FOREST TREES	PHYTOPHTHORA spp. (including <i>P. ramorum</i> , <i>P. cinnamomi</i> , <i>P. palmivora</i> , <i>P. cactorum</i> ) ANTHRACNOSE; APPLE SCAB ( <i>Venturia inaequalis</i> ); BACTERIAL BLIGHT ( <i>Xanthomonas</i> spp.) ARMILLARIA ROOT ROT ( <i>Armillaria</i> spp.); PYTHIUM spp.
CONIFERS GROWN IN COMMERCIAL NURSERIES, PLANTATIONS, and FOREST (INCLUDING CHRISTMAS TREE FARMS)	PHYTOPHTHORA spp. (including <i>P. ramorum</i> , <i>P. cinnamomi</i> , <i>P. palmivora</i> , <i>P. cactorum</i> ) ARMILLARIA ROOT ROT ( <i>Armillaria</i> spp.); PINE PITCH CANKER ( <i>F. subglutinans</i> ); PYTHIUM spp.
CROP TREES	
AVOCADOS	ROOT ROT ( <i>Phytophthora cinnamomi</i> , <i>P. citricola</i> ); ARMILLARIA ROOT ROT ( <i>Armillaria</i> spp.)
CITRUS (mature trees)	ROOT, FOOT and COLLAR ROT ( <i>Phytophthora</i> spp. incl. <i>P. nicotianae</i> , <i>P. citrophthora</i> ) ARMILLARIA ROOT ROT ( <i>Armillaria</i> spp.)
COCONUTS	BUD – NUT FALL ( <i>Phytophthora palmivora</i> ); ARMILLARIA ROOT ROT ( <i>Armillaria</i> spp.)
APPLES, PEARS, LOQUATS, QUINCE	FIRE BLIGHT ( <i>Erwinia amylovora</i> ); APPLE SCAB ( <i>Venturia inaequalis</i> ); ARMILLARIA ROOT ROT ( <i>Armillaria</i> spp.)
TREE NUTS SUCH AS ALMONDS WALNUTS	ROOT and COLLAR ROT ( <i>Phytophthora</i> spp.); ARMILLARIA ROOT ROT ( <i>Armillaria</i> spp.)

**1. The MAUGET SYSTEM**

- (A) Mauget compressible capsule with insert hole
- (B) Feeder tube with flanged gun-sight and opposite tapered beveled end

**2. TOOLS**

- (A) Portable electric drill
- (B) 11/64 in. (0.4 cm) drill bit
- (C) Plastic mallet
- (D) Tape measure
- (E) Insertion tool (optional)

**3. RATE**

Measure the diameter of the tree at chest height in inches.

The dosage is:

0.17 fl. oz. (5 ml) per inch DBH

Round down to the lower whole number.

Injection sites can be placed on first and second vertically spaced circumferential rings where necessary. Vertical injection sites should not be aligned.

**4. PRESSURIZING THE CAPSULES**

Apply the appropriate amount of pressure on the top of the capsule in order to compress.

**5. DRILLING THE TREE HOLE**

Pre-drill spaced injection sites at a slight downward angle at the root flair/buttrass area (approximately 6.0 to 8.0 in., 15 to 20 cm) above ground level, using a clean 11/64 in. (0.4 cm) drill bit (except monocotyledons, conifers, etc.). Drill to a depth of 3/8 to 1/2 in. (0.95 to 1.3 cm) into healthy xylem tissue under the bark. For mini-micro feeder tube, see Step 10. Disinfect drill bit, insertion tool (if used), as well as mini-micro insertion tool prior to use on each tree.

**6. TREE HOLE DEPTH**

It is important that the feeder tube be set to the proper depth in the conductive xylem tissue. If set too deeply, flow is restricted by blockage in the heartwood; if set too shallow, leakage may occur. The feeder tube dispensing end is beveled to allow for a 1/4 in. (0.6 cm) plus tolerance.

**7. COMBINING CAPSULE AND FEEDER TUBE**

Several methods of combining the micro-injector capsule with the feeder tube are acceptable including placing by hand, the feeder tube's flange end, with the flange notch upward, into the micro-injector insert hole of a compressed upright micro-injector capsule. Push the flange end of the feeder tube flush with the membrane located at the inner end of the insert hole.

**8. PLACING THE FEEDER TUBE IN THE TREE**

Firmly seat the beveled, dispensing end of the feeder tube, with the attached upright micro-injector capsule, into the pre-drilled tree injection hole. Tap the rear side, opposite the insert hole of the micro-injector capsule either with an optional mallet, hammer or push forward with the palm your hand. This action will simultaneously seat the feeder tube in the injection hole while breaking the micro-injector capsule membrane for releasing the capsule contents into the feeder tube and into the tree. Another method is to place the feeder tube in the pre-drilled hole of the tree using the optional insertion tool. Then place the compressed micro-injector capsule onto the feeder tube in place.

**9. REMOVAL**

Uptake in the tree usually occurs within several minutes. Capsules may be temporarily rotated in place to see if any liquid is left. When empty, turn the micro-injectors upside down for one minute before removal. Applicators must remove micro-injectors promptly after treatment. Empty micro-injectors must not be left on the tree. The health and species of the tree, and local environmental conditions will determine the rate of uptake. If the capsule does not completely empty within a few hours, invert and carefully remove the micro-injector and enclose it in a heavy duty plastic bag for disposal in accordance with state and local regulations.

**10. MINI-MICRO FEEDER TUBE**

For established trees with thin bark (less than 3/8 in. (0.9 cm) thickness), use a 7/64 in. (0.3 cm) drill bit to produce a micro-injection site for a mini-micro feeder tube.

**WARRANTY STATEMENT**

J.J. Mauget Co. makes no warranty of merchantability, fitness for any purpose or otherwise expressed or implied concerning this product or its uses which extends beyond the use of the product under normal conditions in accord with the statements made on this label.

Sub-label B – For Residential Use

**Mauget**

**ArborFos™**

SYSTEMIC FUNGICIDE  
IN READY TO USE CAPSULES  
FOR TREE INJECTION USE FOR SEASONAL SUPPRESSION OF  
CERTAIN DISEASES

MFG. BY:	J.J. MAUGET CO.
TOWN, STATE:	Arcadia, CA 91006
EPA REGISTRATION NO:	7946-26
EPA ESTABLISHMENT NO:	7946-CA-1
E-MAIL ADDRESS:	<a href="http://www.Mauget.com">www.Mauget.com</a>

**ACTIVE INGREDIENT:**

Mono- and di-potassium salts of Phosphorous Acid ..... 45.8%

**OTHER INGREDIENTS:** ..... 54.2%

Total ..... 100.0%

**Contains 5.174 lbs/gallon of the active ingredient, mono- and di-potassium salts of Phosphorous Acid. Equivalent to 3.353 lbs Phosphorous Acid/gallon.**

Net Contents:

# Mauget

## ArborFos™

SYSTEMIC FUNGICIDE  
IN READY TO USE CAPSULES  
FOR TREE INJECTION USE FOR SEASONAL SUPPRESSION OF  
CERTAIN DISEASES

MFG. BY: J.J. MAUGET CO.  
TOWN, STATE: Arcadia, CA 91006  
EPA REGISTRATION NO: 7946-26  
EPA ESTABLISHMENT NO: 7946-CA-1  
E-MAIL ADDRESS: www.Mauget.com

### ACTIVE INGREDIENT:

Mono- and di-potassium salts of Phosphorous Acid .....45.8%

OTHER INGREDIENTS: .....54.2%

Total 100.0%

Contains 5.174 lbs/gallon of the active ingredient, mono- and di-potassium salts of Phosphorous Acid. Equivalent to 3.353 lbs Phosphorous Acid/gallon.

### Net Contents:

\_\_\_\_\_ 24 capsules plus 24 feeder tubes per carton

\_\_\_\_\_ 24 capsules @ 0.17 fl. oz. (5 mL) ea, 4.06 fl. oz. (120 mL) net per carton

Shipping box: 12 cartons as above.

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

### FIRST AID

#### IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

#### IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

#### IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

#### IF INHALED

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-535-5053** for emergency treatment information.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing vapors or spray mist. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Wear appropriate personal protective equipment

### ENVIRONMENTAL HAZARDS

For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool place over 45°F with capsules in an upright position. Keep out of direct sunlight when possible.

**CONTAINER DISPOSAL:** Dispose of empty capsules in a sanitary landfill or by incineration if approved by state and local authorities. If burned, stay out of smoke.

### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

### RESTRICTIONS

Do not inject trees that are less than four inches in diameter

### GENERAL DIRECTIONS

**Important:** Preventative application is more effective than therapeutic treatment in trees showing disease symptoms. Trees in advanced stages of disease development may not respond to treatment. The health, species of the tree, and the environmental conditions will determine the rate of uptake. Infected trees will absorb the material more slowly due to the vascular plugging caused by the disease. If ArborFos is not absorbed within 24 hours, the tree is considered high risk and has a poor chance of survival. For all rates of application, see #3 below. Thoroughly irrigate trees/plants prior to or at the same time of injection. Test for phytotoxicity prior to regular use.

PLANT	DISEASE
ORNAMENTAL TREES	PHYTOPHTHORA spp. (including P. ramorum, P. cinnamomi, P. palmivora, P. cactorum) ANTHRACNOSE; APPLE SCAB (Venturia inaequalis); BACTERIAL BLIGHT (Xanthomonas spp.) ARMILLARIA ROOT ROT (Armillaria spp.); PYTHIUM spp.; PINE PITCH CANKER (F. subglutinans)
AVOCADOS	ROOT ROT (Phytophthora cinnamomi, P. citricola); ARMILLARIA ROOT ROT (Armillaria spp.)
CITRUS (mature trees)	ROOT, FOOT and COLLAR ROT (Phytophthora spp. incl. P. nicotianae, P. citrophthora) ARMILLARIA ROOT ROT (Armillaria spp.)
COCONUTS	BUD – NUT FALL (Phytophthora palmivora); ARMILLARIA ROOT ROT (Armillaria spp.)
APPLES, PEARS, LOQUATS, QUINCE	FIRE BLIGHT (Erwinia amylovora); APPLE SCAB (Venturia inaequalis); ARMILLARIA ROOT ROT (Armillaria spp.)
TREE NUTS SUCH AS ALMONDS and WALNUTS	ROOT and COLLAR ROT (Phytophthora spp.); ARMILLARIA ROOT ROT (Armillaria spp.)

**1. The MAUGET SYSTEM**

- (A) Mauget compressible capsule with insert hole
- (B) Feeder tube with flanged gun-sight and opposite tapered beveled end

**2. TOOLS**

- (A) Portable electric drill
- (B) 11/64 in. (0.4 cm) drill bit
- (C) Plastic mallet
- (D) Tape measure

**3. RATE**

Apply the compressed capsule every 3 inches around the circumference of the tree. Injection sites can be placed 6 – 8 inches above the soil surface.

**4. PRESSURIZING THE CAPSULES**

Apply the appropriate amount of pressure on the top of the capsule in order to compress.

**5. DRILLING THE TREE HOLE**

Pre-drill spaced injection sites at a slight downward angle at the root flair/buttress area (approximately 6 to 8 in.), above ground level, using a clean 11/64 in. drill bit (except monocotyledons and conifers). Drill to a depth of 3/8 to 1/2 in. into healthy xylem tissue under the bark. For conifers, palms, and other monocots, vary the depths of the injection sites. Depth should be 3 inches or less. Follow manufacturer's directions and precautions in the use of power drills.

**6. TREE HOLE DEPTH**

It is important that the feeder tube be set to the proper depth in the conductive xylem tissue. If set too deeply, flow is restricted by blockage in the heartwood; if set too shallow, leakage may occur.

**7. COMBINING CAPSULE AND FEEDER TUBE**

Carefully insert the feeder tube's flange end, with the notch upward, into the insert hole of a compressed micro-injection capsule without breaking the membrane. Be sure the flange end of the feeder tube is flush against the wall of the capsule insert hole.

**8. PLACING THE FEEDER TUBE IN THE TREE**

Firmly seat the feeder tube, with the attached upright micro-injector capsule, into the pre-drilled tree injection hole. Tap the rear side, opposite the insert hole of the capsule to break the capsule membrane and release the capsule contents into the feeder tube and into the tree.

**9. REMOVAL**

Uptake in the tree usually occurs within several minutes. Capsules may be temporarily rotated in place to see if any liquid is left. When empty, turn the micro-injectors upside down for one minute before removal. Remove micro-injectors promptly after treatment. Empty micro-injectors must not be left on the tree. The health and species of the tree, and local environmental conditions will determine the rate of uptake. If the capsule does not completely empty within a few hours, invert and carefully remove the micro-injector and enclose it in a heavy duty plastic bag for disposal in accordance with state and local regulations.

**10. MINI-MICRO FEEDER TUBE**

For established trees with thin bark (less than 3/8 in. (0.9 cm) thickness), use a 7/64 in. (0.3 cm) drill bit to produce a micro-injection site for a mini-micro feeder tube.

**WARRANTY STATEMENT**

J.J. Mauget Co. makes no warranty of merchantability, fitness for any purpose or otherwise expressed or implied concerning this product or its uses which extends beyond the use of the product under normal conditions in accord with the statements made on this label.



9  
9

Mauget®

# ARBORFOS

5 ml.

EPA REG. NO. 7946-26  
EPA EST. NO. 7946-CA-1


Refer to the outer box label for additional labeling.  
The registered label shall be in the possession of the user during application.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**ACTIVE INGREDIENT:**

Mono- and di-potassium salts of Phosphorous Acid.....	45.8%
OTHER INGREDIENTS:.....	<u>54.2%</u>
<b>TOTAL.....</b>	<b>100.0%</b>

Made in U.S.A. J.J. Mauget Co.  
5435 Peck Road  
Arcadia, CA 91006-5847



Net Contents 5 mL or 0.17 fl. oz.