



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

August 27, 2024

Carrie Tackema
Senior Regulatory Specialist
SciReg, Inc.
12733 Director's Loop
Woodbridge, VA 22192

Subject: Label Amendment – Revising master labeling including adding a sub-label and other minor changes
Product Name: Dinocide
EPA Registration Number: 7946-35
Application Date: 12/08/2023
Case Number: 492770

Dear Ms. Tackema:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or

claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Joseph Belsky at 202-566-2495 or at belsky.joseph@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'Kable Bo Davis', enclosed within a hand-drawn oval.

Kable Bo Davis
Senior Regulatory Specialist
Office of Pesticide Programs
Registration Division, Immediate Office

Enclosure

Master Label
DINOCIDE®

Sublabel A: For ready to use capsules

Sublabel B: For ready to use bottles

GROUP	4A	INSECTICIDE
		
DINOCIDE®		
SYSTEMIC INSECTICIDE FOR TREE INJECTION USE IN READY TO USE CAPSULES AND BOTTLES		
INTENDED FOR USE BY PROFESSIONAL APPLICATORS		
MFG. BY: TOWN, STATE: EPA REGISTRATION NO: EPA ESTABLISHMENT NO:		J.J. MAUGET CO. Arcadia, CA 91006 7946-35 7946-CA-1

ACTIVE INGREDIENT:

Dinotefuran* 12%

OTHER INGREDIENTS: 88%

Total 100%

*Contains 0.13 g/mL N-methyl-N'-nitro-N'-[(tetrahydro-3-furanyl)methyl]guanidine

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

{Note to EPA Reviewer: The Master label constitutes two sublabels. The first aid statements, precautionary statements, PPE, agricultural use requirements box, directions, disease box, and other mandatory label language is consistent between the two sublabels. The difference between the sublabels is to detail out application information specific to each product presentation. Sublabel A application information is specific to applying the product with ready to use capsules. Sublabel B application information is specific to applying the product with injection equipment for product sold in ready to use bottles.}

ACCEPTED
08/27/2024
Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 7946-35



**SYSTEMIC INSECTICIDE
FOR TREE INJECTION USE
IN READY TO USE CAPSULES
INTENDED FOR USE BY PROFESSIONAL APPLICATORS**

MFG. BY:
ADDRESS
TOWN, STATE:
PHONE:

J.J. MAUGET CO.
5435 Peck Road
Arcadia, CA 91006
626-444-1057

EPA REGISTRATION NO:
EPA ESTABLISHMENT NO:

7946-35
7946-CA-1

GROUP

4A

INSECTICIDE

ACTIVE INGREDIENT:

Dinotefuran*..... 12%

OTHER INGREDIENTS:..... 88%

Total..... 100%

*Contains 0.13 g/mL N-methyl-N'-nitro-N"-[(tetrahydro-3-furanyl)methyl]guanidine

Net Contents:

Includes: 24 capsules plus 24 feeder tubes per carton/ pack

____ 24 capsules @ 2mL, 48 mL net, or
____ 24 capsules @ 3mL, 72 mL net, or
____ 24 capsules @ 4mL, 96 mL net, or
____ 24 capsules @ 6mL, 144 mL net

Shipping box: 12 cartons/ pack as above.

Batch Code:

KEEP OUT OF REACH OF CHILDREN CAUTION

[See inside label for [complete][additional] [First Aid,][Precautionary Statements,][Directions for Use,] [and] [Storage and Disposal.]

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

FIRST AID**IF SWALLOWED**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything to an unconscious person.

IF ON SKIN

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

IF INHALED

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact INFOTRAC 1-800-535-5053 for emergency treatment information.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow contact with water or oxidizing agents. Hazardous chemical reactions may occur. Do not use or store near heat or open flame.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not dispose equipment washwaters or rinsate into a natural drain or water body.

This product is toxic to honeybees. The persistence of residues and potential residual toxicity of Dinotefuran in nectar and pollen suggests the possibility of chronic toxic risk to honeybee larvae and the eventual instability of the hive.

This product is toxic to bees exposed to treatment for more than 38 hours following treatment.

Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

Dinotefuran and its degradate, MNG, have the properties and characteristics associated with chemicals detected in ground water. The high water solubility of dinotefuran, and its degradate, MNG, coupled with its very high mobility, and resistance to biodegradation indicates that this compound has a strong potential to leach to the subsurface under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Periodic monitoring of shallow groundwater in the use area is recommended.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

APPLICATORS AND OTHER HANDLERS MUST WEAR:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of polyethylene or butyl rubber or neoprene rubber or Viton ≥14 mil
- Shoes plus socks
- Protective eyewear such as goggles, face shield or safety glasses

Resistance Management

DINOCIDE® contains a Group 4A insecticide. Insect biotypes with acquired resistance to Group 4A may eventually dominate the insect/mite population if Group 4A insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by DINOCIDE® or other Group 4A pesticides. To delay insecticide resistance, consider:

- Avoiding the consecutive use of DINOCIDE® or other Group 4A insecticides that have a similar target site of action, on the same insect species.
- Basing insecticide use on a comprehensive IPM program.
- Monitoring treated insect populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Restrictions

- Do not inject trees that are less than two inches in diameter at breast height (DBH) (6 inches in circumference).
- This product is NOT to be used on trees which will produce food within the year (365 days) following treatment unless food crop on treated tree is discarded and destroyed.
- Do not apply this product, by any application method, to linden, basswood, or other *Tilia* species in the state of Oregon.
- Do not apply this product more than one time per year.

Read entire label, use strictly in accordance with precautionary statements and directions, and with applicable state and federal regulations. Failure to follow label directions may result in poor control or tree injury.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and the handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

There are no reentry or Personal Protective Equipment requirements for this product.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

There are no reentry or protective clothing requirements for this product.

USE SITES

DINOCIDE® is for use on hardwood or deciduous trees, evergreens and palms grown in forests, woodlands, ornamental landscapes such as residential and commercial settings, median strips, roadways, Christmas tree farms, cemeteries, parks, and interior and exterior plantscapes. This product is not registered for sale or use in Nevada.

TARGET INSECTS

on Forest, Park, Landscape, Palm, and Ornamental Trees

ADELGIDS APHIDS BLACK VINE WEEVIL ADULT BRONZE BIRCH BORER COCONUT RHINOCEROS BEETLE [*] COTTONWOOD LONGHORNED BORER DOUGLAS FIR GALL MIDGE DOUGLAS FIR CONE MOTH LARVAE ELM LEAF BEETLE EUCALYPTUS LONGHORNED BORER FLATHEADED BORER (including Alder Borer and Birch Borer; excluding Emerald Ash Borer) JAPANESE BEETLE	LACEBUGS LEAFHOPPERS LEAFMINERS LOBATE LAC SCALE[*] MEALYBUGS PINE TIP MOTH LARVAE PSYLLIDS (including Lerp Psyllid) ROYAL PALM BUGS SCALE INSECTS (armored and soft, including Asian Cycad Scale) SPOTTED LANTERNFLY SPRUCE BUDWORM THRIPS WHITEFLIES
---	--

[*Not registered for sale or use in California]

FACTORS AFFECTING APPLICATION

Applications are most effective when made prior to insect infestation and in conjunction with good cultural management practices. The species and health of the tree, as well as local environmental conditions, will determine the rate of uptake when using the Mauget System. Uptake time in the tree usually occurs within several minutes to over an hour, but trees in advanced stages of insect infestation may not respond to treatment. If DINOCIDE® is not absorbed within 24 hours (barring any applicator error or malfunction of injection device, or environmental factors affecting transpiration) the tree may be considered high risk with a possible poor chance of survival.

Environmental Conditions

This technology relies on the natural uptake rate of the tree; and thus, factors that affect the transpiration rate can greatly affect the uptake rate. Transpiration is dependent upon a number of factors, such as soil moisture, soil and air temperatures, and time of day. For optimum uptake, apply when soil moisture is adequate and soil temperatures are above 45°F. Preferred conditions for injections are morning to early afternoon hours, with warm temperatures (55-85°F / 13-30°C), accompanied by low humidity, clear skies and a slight breeze. Sunny conditions along with moist soil and a well-hydrated tree will also increase the transpiration rate and will therefore improve uptake. Conversely, cool temperatures, cloudy and/or evening skies and trees under moisture stress will slow down the rate of uptake. Extreme heat and cold temperatures will adversely affect rates as well.

Trees that have healthy vascular systems will have correspondingly higher uptake rates. Trees in advanced stages of pest development may not respond to treatment, as vascular plugging caused by disease inhibits transpiration. If DINOCIDE® has not started to absorb within two hours, consider removing the device (following the proper sequence provided in the removal instructions) and drill a new hole in a different area of the trunk and inject again. The injection devices need to be evenly spaced at points on the trunk free of visible decay areas and wounds from the point of injection to where branching begins. If DINOCIDE® has not started to absorb within one hour after the second attempt, the vascular system of the tree may be too compromised for treatment or there is significant decay in that local injection area.

DO NOT inject trees that are drought stressed. Applications to drought or heat stressed trees may result in injury to tree tissue, poor treatment, and subsequently poor control. Avoid treating trees that are moisture stressed or suffering from herbicide damage.

Monitor Tree Health and Pest Infestations

Preventative application is more effective than therapeutic treatment in trees showing insect infestation symptoms. Effective injection treatment is favored by a full canopy (i.e., leaves) and a healthy vascular system. Once these tissues are compromised by pest damage (larval galleries, defoliation, leaf mining, etc.), an effective and uniform application of DINOCIDE® may be difficult to achieve and subsequent control may be poor. For optimal results, treat at least 2 to 3 weeks before pests historically infest the host tree. As a result of systemic movement and longevity of DINOCIDE® in trees, the interval may be extended much earlier to 6 months should tree dormancy, adverse weather, management, asynchronous life cycle of pests, etc., allow earlier application timing.

DINOCIDE® may also be effective as a remedial treatment against some pests, such as those with slower development or if multiple life stages are susceptible to the active ingredient. Pests that attack the stem and branches may disrupt vascular tissue resulting in poor distribution in an infested tree. However, control may be achieved if larvae come into contact or feed on DINOCIDE® treated tree tissues.

APPLICATION INSTRUCTIONS

Timing of Application:

Preventive applications approximately 2 to 4 weeks prior to the anticipated feeding damage will provide better management, but rescue treatments will also perform well with acceptable minimal damage. DINOCIDE® can also be used after damage has occurred against listed insect pests that produce large amounts of feeding debris. Focus timing and treatment on the most susceptible stage of the listed pest.

Sequential and/or Combination Treatments:

When treating for beetles that carry fungi (ambrosia), an additional treatment of fungicide may improve management strategies. Materials to consider are fungicides labeled for use against vascular-inhabiting fungi.

DINOCIDE® may be sequentially applied with other insecticides, such as abamectin, for a more broad-spectrum treatment.

Application Tips:

To account for trunk flare, place injection sites evenly around the base of the root flare within 6 to 8 inches of the root crown. For pines and other resinous conifer species, injection sites may be higher up on the trunk (see Step 2 below). Follow good injection practices. Disinfect drill bit prior to use on each tree.

Tree measurement guidance:

Dosages are based on the circumference OR the diameter (inches or centimeters) of the tree at breast height ("DBH"). DBH is the outside bark diameter of the trunk at 4.5 feet (1.4 m) above the ground on the uphill side of the tree. For the purposes of determining breast height, the ground includes the duff layer that may be present, but does not include unincorporated woody debris that may rise above the ground line.

The diameter is determined by measuring the circumference of the tree at breast height, and dividing circumference (in inches) by three (3). To determine DBH for multi-stemmed woody ornamentals, measure the DBH of each stem or branch and add together for the total DBH per tree.

MAUGET CAPSULES



Mauget compressible capsule with feeder tube insert hole
Feeder tube* with flanged gunsight ("V" notch) and opposite tapered beveled end


*Standard feeder tube provided is 2 3/4" using an 11/64" drill bit. Additional sizes available are Micro-Mini (1 1/4"), Micro (1 3/4") and Large (4 1/2"). Micro-Mini and Micro would be used on trees with thin bark, using a 7/64" drill bit. Large would be used on trees with thick bark, using the standard 11/64" drill bit.

TOOLS

Portable electric drill
11/64 in. (0.4 cm) drill bit
Plastic mallet
Tape measure
Insertion tool (optional)
Mauget Compression tool (optional)

1. NUMBER OF CAPSULES

- Measure the circumference or diameter of the tree at chest/breast height (approximately 4.5 feet from the soil-line) using a tape measure or an arborist diameter tape.
- If measuring the circumference (CBH), divide this number by six (6) to determine the number of capsules needed.
- If measuring the diameter (DBH), divide this number by two (2) to determine the number of capsules needed.
- If the number of capsules results in a fraction, round down to the lower whole number.
- To account for trunk flare, place injection sites evenly (every 6-8 inches) around the base of the root flare within 6 to 8 inches of the root crown.
- For pines and other resinous conifer species, injection sites may be higher up on trunk.



The following dosage, per capsule, depends on tree diameter:

- 2mL capsules – 2 to 10 inches DBH
- 3mL capsules – 11 to 36 inches DBH
- 4mL capsules – 37 inches DBH and above
- 6mL capsules – narrow-leaved evergreen trees

For heavier infestation and/or more persistent insects, use 4 mL capsules on all tree sizes. For narrow-leaved evergreen trees such as pine, spruce, or hemlock, use the 6 mL capsules at 4-inch circumference spacing (DBH divided by 1.3). Trees in advanced stages of insect infestation may not respond to treatment. The health, species of the tree and the environmental conditions will determine the rate of uptake.

For palms and other monocotyledons, when using multiple injection sites, alternate drilling depth to capture scattered vascular bundles, taking care that the depth of any single site is less than 1/3 the diameter of the tree. If using a single injection site, one feeder tube can be used to administer the contents of all capsules. Injection sites may be covered with Lac Balsam or similar for aesthetic purposes.

2. DRILLING THE TREE HOLE

- Unless otherwise noted, predrill, at approximately 6 in. intervals, injection sites at a slight downward angle at the root flair/butress area (approximately 6.0 to 8.0 in., 15 to 20 cm) above ground level, using a clean 11/64 in. (0.4 cm) drill bit (except monocotyledons, conifers, etc.).
- For conifers and other species with high resin pressure, place injection sites higher on the trunk (36-48") and to a deeper drill depth of 2+ inches. After drilling the injection hole, immediately insert the feeder tube and capsule into the hole to reduce resin flow pressure.
- Drill to a depth of 3/8 to 1/2 in. (0.95 to 1.3 cm) into healthy xylem tissue under the bark.
- It is important that the feeder tube be set to the proper depth in the conductive xylem tissue. If set too deeply, flow is restricted by blockage in the heartwood; if set too shallow, leakage may occur.
- The feeder tube dispensing end is beveled to allow for a 1/4 in. plus tolerance.
- Disinfect drill bit and insertion tool (if used) prior to use on each tree.
- Note: Feeder tube can be placed in the hole at this time to hold/mark the spot until capsules are ready to use.

3. PRESSURIZING THE CAPSULES

- To pressurize capsules firmly press down on colored top until it clicks using your hands/thumbs or a Mauget compression tool.
- In high altitude conditions, pressing the colored top down to the locking position may be difficult. This is an indication of high internal capsule pressure that results in faster diffusion of product into the tree.
- To maintain optimal pressure, monitor the capsules after installation and re-pressurize units when half of the dosage is administered.

If the center button depresses too easily and drainage is slow, pre-puncturing the capsule port membrane with the feeder tube prior to installation will equalize pressure to allow for diffusion. See 4 h-k below.

4. COMBINING CAPSULE AND FEEDER TUBE - Three methods of combining the capsule with the feeder tube are acceptable.

- By hand, place the feeder tube's flange end, with the flange notch upward, into the insert hole of a pressurized/compressed upright capsule.
 - Push the flange end of the feeder tube flush with the membrane located at the inner end of the insert hole.
 - Go to Step 5a.
- OR:**
- Place the feeder tube in the predrilled hole using the optional insertion tool.
 - Remove the insertion tool.
 - Place the pressurized/compressed capsule onto the feeder tube.
 - Go to Step 5b.
- OR:**
- With capsule and feeder tube pointing upward, fully insert the feeder tube into the capsule, breaking the capsule membrane.
 - Insert the beveled end of the feeder tube and capsule into the predrilled hole.
 - Tap the rear side of the capsule with a mallet or the heel of your hand to firmly seat the feeder tube into the hole.
 - Pressurize the capsule by carefully pressing down the colored top of the capsule until it clicks using either your thumbs or a Mauget compression tool.
 - Go to Step 6.

5. PLACING THE FEEDER TUBE IN THE TREE

- Firmly seat the beveled, dispensing end of the feeder tube, with the attached upright capsule, into the predrilled injection hole.
- Tap the rear side, opposite the insert hole of the capsule with a plastic mallet or the heel of your hand. This action will simultaneously seat the feeder tube in the injection hole while breaking the capsule membrane for releasing the capsule contents into the feeder tube and into the tree.

6. REMOVAL

- Uptake in the tree usually occurs within several minutes.
- Capsules may be temporarily rotated in place to see if any liquid is left.
- When empty, turn the capsules upside down for one minute before removal.

- d. Applicators must remove capsules promptly after treatment; empty capsules must not be left on the tree.
- e. The health and species of the tree, and local environmental conditions will determine the rate of uptake.
- f. If the capsule does not completely empty within a few hours, invert and carefully remove the capsule and enclose it in a heavy-duty plastic bag for disposal in accordance with state and local regulations.

Retreatment

At time of initial application, make note of the health level of each tree. Reevaluate health level in treated trees at 12-month intervals to determine the need for retreatment. Consider preventive applications 12-36 months after the initial treatment. Evaluate trees in high pest pressure areas or highly valued trees for retreatment if symptoms progress or 12 months after each treatment. Repeat treatment after 12 months, if needed. Follow application procedures described above for repeat injections; new drill holes will be required for subsequent treatments. Stagger the holes equally in subsequent applications to ensure proper uptake.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep pesticide in original container. Store in a cool (45°F-90°F), dry place out of direct sunlight and out of reach of children and animals.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or dispose of in a sanitary landfill, or by incineration if approved by state and local authorities. Do not burn unless allowed by state and local ordinances. If burned, stay out of smoke.

NOTICE OF WARRANTY

To the extent consistent with applicable law, J.J. Mauget Co. makes no warranty of merchantability, fitness for any purpose or otherwise expressed or implied concerning this product or its uses which extends beyond the use of the product under normal conditions in accord with the statements made on this label.

List of optional label claims:

- Begins absorbing upon injection
- Connect directly to the tree
- Convenient (Simple) (Design)
- Easy(-ier) to use (convenient, handy, useful, well-situated)
- Easy to apply
- Easy to store
- Easy (to use)
- Easy to use application
- Economical
- Even if it rains - it won't lose its effectiveness
- Ideal for targeting specific infested locations (in orchards)
- It works!
- Long Lasting Protection
- Long lasting treatment
- No mix, no mess
- No mixing (necessary) (required)
- Once it enters the tree it won't wash off
- Only takes a few minutes to inject
- Outdoor use only
- Preventative treatment
- Provides maximum control
- Ready to Use
- Shake well before application (applying)
- Ready to Use applicator (device)
- Quick and easy to use
- Ready to Use – No Mixing, No Measuring
- Requires no mixing
- Results that show
- Easier, more economical! Simple (and easy) (to use)
- Systemic treatment of undesirable pests (and disease)
- (The) Next Generation of Tree Care (tree injection)
- The key ingredient in this product (active ingredient) targets the specific pest
- Trusted results for satisfaction
- Kills listed pests
- Works!

Promotional options:

- Save up to \$(x) on (your) next purchase
- (X)% free (more)
- Great value
- Bonus size
- Reasonably priced
- Additional savings
- Cost saving consumables
- Economical
- Invest in labor rather than equipment (machinery)

PACKAGING RELATED CLAIMS:

- Connect directly to tree
- Controlled delivery
- Convenient
- Quick connect applicator
- New(!)
- Quick connect applicator (device)
- Quick connect injection (applicator) (device)
- Quick & (and) easy to use
- Delivers direct treatment
- Direct injection (inject) (injecting)
- Easy
- Easy and convenient
- Easy to store (and dispose) (see instructions)
- Easy to use
- Eliminate(s) mixing
- The easy way to treat (with Dinocide®)
- The easier way to treat (with Dinocide®)
- No (more) hand fatigue
- No constant trigger squeezing
- No more pumping
- No pumping just inject
- No cumbersome set up

- No equipment clean up
- No costly machines or bulky equipment
- No high pressure forced delivery
- No product wastage
- No direct contact with product solutions if used according to directions
- Only takes a few minutes to apply (and lasts all season long)
- Precise control for maximum accuracy
- Prepackaged dinotefuran active ingredient
- Consistent injection for maximum accuracy
- No more tired (aching) hands
- Fast and easy application
- Easy to use application (injection)
- Easy to use injection
- Change the way you (tree inject) (inject)
- No more spray! Inject!
- Ideal for large or small jobs (areas)
- Great for large or small jobs (areas)
- Precise control
- The easier way to control pests (disease)
- The easier way to suppress pests (disease)
- Easier, more economical
- Save (Saves) time and energy
- Give your hands a break
- Convenient
- Effective
- Efficient (concentrations)
- Economical
- No more bending over
- Fast and easy application
- Full and efficient delivery
- Improved Applicator (application) Device!
- Simple!
- Improved Applicator (Application) System! (Device!)
- Improved Applicator (Application)!
- Improved!
- Improved for lasting (better) results!
- New and Improved!
- No (More) Hand Fatigue!
- No Mix, No Mess
- No more pumping, no more pulling, just inject
- No spray! Just inject!
- No more tired (aching) hands
- Minimal investment in equipment and consumables
- Precise control for maximum accuracy
- Quick & Easy to Use
- Quick (and) natural absorption
- Save (Saves) time and energy
- Simplified equipment
- Simplified calculations
- Simplified pre-filled ready to use devices
- Targeted injection
- Target infested spots
- The easy (-ier) way to control pests (and disease)
- The easy way to inject
- The fast and easy way to inject!
- You're always ready to inject!
- Change the way you (tree) inject
- (New) Comfort injector
- Treat(ment) of more trees in less time
- No complicated mixing steps of chemistries
- Only takes a few minutes to apply (and lasts all season long)



DINOCIDE®

**SYSTEMIC INSECTICIDE
FOR TREE INJECTION USE
IN LIQUID LOADABLE INJECTORS
INTENDED FOR USE BY PROFESSIONAL APPLICATORS**

MFG. BY:
ADDRESS
TOWN, STATE:
PHONE:

J.J. MAUGET CO.
5435 Peck Road
Arcadia, CA 91006
626-444-1057

EPA REGISTRATION NO:
EPA ESTABLISHMENT NO:

7946-35
7946-CA-1

GROUP

4A

INSECTICIDE

ACTIVE INGREDIENT:

*Dinotefuran 12%

OTHER INGREDIENTS: 88%

Total 100%

* Contains 0.13 g/mL N-methyl-N'-nitro-N'-[(tetrahydro-3-furanyl)methyl]guanidine

Net Contents:

_____ 1 bottle @ 1 liter (33.8 fl. oz.)

_____ 1 bottle @ 500 mL (16.9 fl. oz.)

Shipping box: 4 bottles as above

Batch Code:

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

[See inside label for [complete][additional] [First Aid,][Precautionary Statements,][Directions for Use,] [and] [Storage and Disposal.]

FIRST AID**IF SWALLOWED**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything to an unconscious person.

IF ON SKIN

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

IF INHALED

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact INFOTRAC 1-800-535-5053 for emergency treatment information.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not mix or allow contact with water or oxidizing agents. Hazardous chemical reactions may occur.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark.

Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not dispose equipment washwaters or rinsate into a natural drain or water body.

This product is toxic to honeybees. The persistence of residues and potential residual toxicity of Dinotefuran in nectar and pollen suggests the possibility of chronic toxic risk to honeybee larvae and the eventual instability of the hive.

This product is toxic to bees exposed to treatment for more than 38 hours following treatment.

Do not apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

Dinotefuran and its degradate, MNG, have the properties and characteristics associated with chemicals detected in ground water. The high water solubility of dinotefuran, and its degradate MNG, coupled with its very high mobility, and resistance to biodegradation indicates that this compound has a strong potential to leach to the subsurface under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Periodic monitoring of shallow groundwater in the use area is recommended.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION. Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

APPLICATORS AND OTHER HANDLERS MUST WEAR:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of polyethylene or butyl rubber or neoprene rubber or Viton >14 mil
- Shoes plus socks
- Protective eyewear such as goggles, face shield or safety glasses

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Resistance Management

DINOCIDE® contains a Group 4A insecticide. Insect biotypes with acquired resistance to Group 4A may eventually dominate the insect/mite population if Group 4A insecticides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by DINOCIDE® or other Group 4A pesticides. To delay insecticide resistance, consider:

- Avoiding the consecutive use of DINOCIDE® or other Group 4A insecticides that have a similar target site of action, on the same insect species.
- Basing insecticide use on a comprehensive IPM program.
- Monitoring treated insect populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Restrictions

- Do not inject trees that are less than two inches in diameter at breast height (DBH) (6 inches in circumference).
- This product is NOT to be used on trees which will produce food within the year (365 days) following treatment unless food crop on treated tree is discarded and destroyed.
- Do not apply this product, by any application method, to linden, basswood or other *Tilia* species in the state of Oregon.
- Do not apply this product more than one time per year.

Read entire label, use strictly in accordance with precautionary statements and directions, and with applicable state and federal regulations. Failure to follow label directions may result in poor control or tree injury.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and the handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

There are no reentry or Personal Protective Equipment requirements for this product.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

There are no reentry or personal protective clothing requirements for this product.

USE SITES

DINOCIDE® is for use on hardwood or deciduous trees, evergreens and palms grown in forests, woodlands, ornamental landscapes such as residential and commercial settings, median strips, roadways, Christmas tree farms, cemeteries, parks, and interior and exterior plantscapes.

This product is not for registered for sale or use in Nevada.

TARGET INSECTS

on Forest, Park, Landscape, Palm, and Ornamental Trees

ADELGIDS APHIDS BLACK VINE WEEVIL ADULT BRONZE BIRCH BORER COCONUT RHINOCEROS BEETLE[*] COTTONWOOD LONGHORNED BORER DOUGLAS FIR GALL MIDGE DOUGLAS FIR CONE MOTH LARVAE ELM LEAF BEETLE EUCALYPTUS LONGHORNED BORER FLATHEADED BORER (including Alder Borer and Birch Borer; excluding Emerald Ash Borer) JAPANESE BEETLE	LACEBUGS LEAFHOPPERS LEAFMINERS LOBATE LAC SCALE[*] MEALYBUGS PINE TIP MOTH LARVAE PSYLLIDS (including Lerp Psyllid) ROYAL PALM BUGS SCALE INSECTS (armored and soft, including Asian Cycad Scale) SPOTTED LANTERNFLY SPRUCE BUDWORM THRIPS WHITEFLIES
--	--

[*Not registered for sale or use in California]

FACTORS AFFECTING APPLICATION

Applications are most effective when made prior to insect infestation and in conjunction with good cultural management practices. The species and health of the tree, as well as local environmental conditions, will determine the rate of uptake when using the Mauget System. Uptake time in the tree usually occurs within several minutes to over an hour, but trees in advanced stages of insect infestation may not respond to treatment. If DINOCIDE® is not absorbed within 24 hours (barring any applicator error or malfunction of injection device, or environmental factors affecting transpiration) the tree may be considered high risk with a possible poor chance of survival.

Environmental Conditions

This technology relies on the natural uptake rate of the tree; and thus, factors that affect the transpiration rate can greatly affect the uptake rate. Transpiration is dependent upon a number of factors, such as soil moisture, soil and air temperatures, and time of day. For optimum uptake, apply when soil moisture is adequate and soil temperatures are above 45°F. Preferred conditions for injections are morning to early afternoon hours, with warm temperatures (55-85°F /13-30°C), accompanied by low humidity, clear skies and a slight breeze. Sunny conditions along with moist soil and a well-hydrated tree will also increase the transpiration rate and will therefore improve uptake. Conversely, cool temperatures, cloudy and/or evening skies and trees under moisture stress will slow down the rate of uptake. Extreme heat and cold temperatures will adversely affect rates as well.

Trees that have healthy vascular systems will have correspondingly higher uptake rates. Trees in advanced stages of pest development may not respond to treatment, as vascular plugging caused by disease inhibits transpiration. If DINOCIDE® has not started to absorb within two hours, consider removing the device (following the proper sequence provided in the removal instructions) and drill a new hole in a different area of the trunk and inject again. The injection devices need to be evenly spaced at points on the trunk free of visible decay areas and wounds from the point of injection to where branching begins. If DINOCIDE® has not started to absorb within one hour after the second attempt, the vascular system of the tree may be too compromised for treatment or there is significant decay in that local injection area.

DO NOT inject trees that are drought stressed. Applications to drought or heat stressed trees may result in injury to tree tissue, poor treatment and subsequently poor control. Avoid treating trees that are moisture stressed or suffering from herbicide damage.

Monitor Tree Health and Pest Infestations

Preventative application is more effective than therapeutic treatment in trees showing insect infestation symptoms. Effective injection treatment is favored by a full canopy (i.e., leaves) and a healthy vascular system. Once these tissues are compromised by pest damage (larval galleries, defoliation, leaf mining, etc.), an effective and uniform application of DINOCIDE® may be difficult to achieve and subsequent control may be poor. For optimal results, treat at least 2 to 3 weeks before pests historically infest the host tree. As a result of systemic movement and longevity of DINOCIDE® in trees, the interval may be extended much earlier to 6 months should tree dormancy, adverse weather, management, asynchronous life cycle of pests, etc., allow earlier application timing.

DINOCIDE® may also be effective as a remedial treatment against some pests, such as those with slower development or if multiple life stages are susceptible to the active ingredient. Pests that attack the stem and branches may disrupt vascular tissue resulting in poor distribution in an infested tree. However, control may be achieved if larvae come into contact or feed on DINOCIDE®-treated tree tissues.

APPLICATION INSTRUCTIONS

Timing of Application:

Preventive applications approximately 2 to 4 weeks prior to the anticipated feeding damage will provide better management, but rescue treatments will also perform well with acceptable minimal damage. DINOCIDE® can also be used after damage has occurred against listed insect pests that produce large amounts of feeding debris. Focus timing and treatment on the most susceptible stage of the listed pest.

Sequential and/or Combination Treatments:

When treating for beetles that carry fungi (ambrosia), an additional treatment of fungicide may improve management strategies. Materials to consider are fungicides labeled for use against vascular-inhabiting fungi.

DINOCIDE® may be sequentially applied with other insecticides, such as abamectin, for a more broad-spectrum treatment.

Application Tips:

To account for trunk flare, place injection sites evenly around the base of the root flare within 6 to 8 inches of the root crown. For pines and other resinous conifer species, injection sites may be higher up on the trunk (see instructions below). Follow good injection practices. Disinfect drill bit prior to use on each tree.

Tree measurement guidance

Dosages are based on the circumference OR the diameter (inches or centimeters) of the tree at breast height ("DBH"). DBH is the outside bark diameter of the trunk at 4.5 feet (1.4 m) above the ground on the uphill side of the tree. For the purposes of determining breast height, the ground includes the duff layer that may be present but does not include unincorporated woody debris that may rise above the ground line.

The diameter is determined by measuring the circumference of the tree at breast height, and dividing circumference (in inches) by three (3). To determine DBH for multi-stemmed woody ornamentals, measure the DBH of each stem or branch and add together for the total DBH per tree.




APPLICATION EQUIPMENT

DINOCIDE® is a solvent-based insecticide for use with liquid loadable injectors. Test DINOCIDE® compatibility with the injector and its liquid contact components before regular use. Clean the interior of the injector with a solvent cleaner to remove any residual liquid from prior use. Disinfect tree injector tips between each tree injection. Use liquid loadable injector equipment/devices with care and in a safe manner, in addition to following each manufacturer's directions and precautions.

DIRECTIONS

Number of Micro-Injection Sites:

Measure the tree at chest height in inches at a level of 4.5 feet from the soil-line using a tape measure (DBH) or an arborist diameter tape (circumference). If measuring the diameter at breast height (DBH), divide this number by two (2) to determine the number of injection sites needed. If measuring the circumference, divide this number by six (6) to determine the number of injection sites needed. If the number of injection sites results in a fraction, round down to the lower whole number.



Dosage:

Use the following dosage, depending on tree diameter:

- 1 mL per inch DBH for trees 2 to 10 inches DBH **or** 2 mL per injection site every 6 inches of circumference
- 1.5 mL per inch DBH for trees 10 to 36 inches DBH **or** 3 mL per injection site every 6 inches of circumference
- 2 mL per inch DBH for trees 36 inches DBH **or** 4 mL per injection site every 6 inches of circumference
- 3 mL per inch DBH for narrow-leaved evergreen trees 24 inches and above **or** 6 mL per injection site, spaced every 4 inches of circumference (DBH/1.3)

For heavier infestations and/or more persistent insects, use the 2 mL per inch rate on all tree sizes for both hardwood and narrow-leaved evergreen trees.

For narrow-leaved evergreen trees, use the closer injection interval of every 4 inches of circumference (DBH/1.3). Increase the dosage to 3.0 mL per inch with narrow-leaved evergreen trees 24 inches or greater in diameter.

Drilling the Hole:

Depending on the specific type of applicator, it may be necessary to predrill holes prior to insertion of the applicator tip into the tree.

To account for trunk flare, place injection sites evenly (every 6-8 inches) around the base of the root flare within 6 to 8 inches of the root crown avoiding root valleys.

Drill holes deep enough to allow the tree's vascular system to transport Dinocide® throughout the tree.

Make injection holes at least $\frac{1}{2}$ to $\frac{3}{4}$ inch into healthy xylem (white wood) with actual depth up to 2 inches or more from the outer trunk surface depending upon the tree species and outer bark thickness.

For conifer species with high resin pressure, place injection sites higher on the trunk (36-48") and to a deeper drill depth of 2+ inches.

Follow good injection practices. Disinfect drill bit prior to use on each tree.

Application and Use:

Set the injector for the appropriate amount of Dinocide® per site and follow injector manufacturer's instructions.

For treatment of pinewood wilt nematode, space injection sites every 4 inches of circumference.

For pines and other resinous conifer species, injection sites may be higher up on trunk (see above).

Uptake Time:

Uptake time in the tree usually occurs within several minutes to an hour dependent upon species of tree, environmental conditions, and the type of tree injector. Trees in advanced stages of insect infestation may not respond to treatment.

Retreatment

At time of initial application, make note of the health level of each tree. Reevaluate health level in treated trees at 12-month intervals to determine the need for retreatment. Consider preventive applications 12-36 months after the initial treatment. Evaluate trees in high pest pressure areas or highly valued trees for retreatment if symptoms progress or 12 months after each treatment. Repeat treatment after 12 months, if needed. Follow application procedures described above for repeat injections; new drill holes will be required for subsequent treatments. Stagger the holes equally in subsequent applications to ensure proper uptake.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep pesticide in original container. Store in an upright position, above 45°F, in a cool, dry place.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or dispose of in a sanitary landfill, or by incineration if approved by state and local authorities. Do not burn unless allowed by state and local ordinances. If burned, stay out of smoke.

NOTICE OF WARRANTY

To the extent consistent with applicable law, J.J. Mauget Co. makes no warranty of merchantability, fitness for any purpose or otherwise expressed or implied concerning this product or its use which extends beyond the use of the product under normal conditions in accord with the statements made on this label.

List of optional label claims:

- Begins absorbing upon injection
- Economical
- Even if it rains - it won't lose its effectiveness
- Ideal for targeting specific infested locations (in orchards)
- It works!
- Long Lasting Protection
- Long lasting treatment
- No mix, no mess
- No mixing (necessary) (required)
- Once it enters the tree it won't wash off
- Only takes a few minutes to inject
- Outdoor use only
- Preventative treatment
- Provides maximum control
- Ready to Use
- Shake well before application (applying)
- Quick and easy to use
- Ready to Use – No Mixing, No Measuring
- Requires no mixing
- Results that show
- Easier, more economical! Simple (and easy) (to use)
- Systemic treatment of undesirable pests (and disease)
- (The) Next Generation of Tree Care (tree injection)
- The key ingredient in this product (active ingredient) targets the specific pest
- Trusted results for satisfaction
- Kills listed pests
- Works!

Promotional options:

- Save up to \$(x) on (your) next purchase
- (X)% free (more)
- Great value
- Bonus size
- Reasonably priced
- Additional savings
- Cost saving consumables
- Economical

PACKAGING RELATED CLAIMS:

- Connect directly to tree
- Controlled delivery
- Convenient
- Delivers direct treatment
- Direct injection (inject) (injecting)
- Easy
- Easy and convenient
- Easy to store (and dispose) (see instructions)
- Easy to use
- Eliminate(s) mixing
- The easy way to treat (with Dinocide®)
- The easier way to treat (with Dinocide®)
- Only takes a few minutes to apply (and lasts all season long)
- Precise control for maximum accuracy
- Change the way you (tree inject) (inject)
- No more spray! Inject!
- Ideal for large or small jobs (areas)
- Great for large or small jobs (areas)
- Precise control
- The easier way to control pests (disease)
- The easier way to suppress pests (disease)
- Easier, more economical
- Save (Saves) time and energy
- Convenient
- Effective
- Efficient (concentrations)
- Economical
- No more bending over
- Fast and easy application
- Full and efficient delivery
- Simple!
- Improved Applicator (Application) System! (Device!)

- No Mix, No Mess
- No spray! Just inject!
- Quick (and) natural absorption
- Save (Saves) time and energy
- Simplified calculations
- Targeted injection
- Target infested spots
- The easy (-ier) way to control pests (and disease)
- Change the way you (tree) inject
- Treat(ment) of more trees in less time
- No complicated mixing steps of chemistries
- Only takes a few minutes to apply (and lasts all season long)