



Kauritil

Wettable Powder Copper Fungicide

ACTIVE INGREDIENT:

Copper Oxidechloride 80%
(Copper expressed as metallic 47%)

INERT INGREDIENTS 20%

EPA Reg. No. 7966-27

CAUTION:

KEEP OUT OF REACH OF CHILDREN.
HARMFUL IF SWALLOWED.

Dispose of wastes by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning.

TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS AND PONDS.

Net Weight ~~50 Pounds~~

BASF Wyandotte Corporation Parsippany, New Jersey 07054

DIRECTIONS FOR USE

General Instructions. Kauritil is an insoluble, copper fungicide with reduced phytotoxic potential. Repeat applications at 10-14 day intervals may be necessary to maintain disease control. Use sufficient spray volume to obtain good coverage of crop, and increase volume as crop grows larger.

Instructions given are based on general applications. Consult Experiment Station or Extension Service fungicide specialist for specific use directions designed for your area.

Mixing Directions. Fill the spray tank half full of water and, with agitator running, slowly add the required amount of Kauritil. Other spray materials should be added only after Kauritil is thoroughly mixed. Finish filling spray tank and apply at once. Kauritil is compatible with commonly used insecticides and nutrients, and with wettable sulfur.

CARROTS — *Cercospora Leaf-spot, Leaf Blight.* Use 3 to 4.5 pounds per acre beginning when disease threatens in area and continuing at 7 to 14 day intervals as necessary to protect foliage. Use sufficient spray volume for good coverage of crop.

CELERY — *Early and Late Blight.* Use 3 to 7 pounds per acre beginning as soon as plants become established in the field. Continue at 5 to 7 day intervals depending on disease severity. Use sufficient spray volume for good coverage of crop.

CITRUS (Florida) — *Melanose, Scab, Greasy Spot.* Use 6.5 pounds in 500 gallons of water per acre. Apply in dormant season before spring flush and repeat within two weeks after petal fall. For Melanose and Scab, repeat again in four weeks using 5 pounds. Do not apply after fruit reaches 3/4 inch diameter.

POTATOES — *Early and Late Blight.* Use 3 to 7 pounds per acre and begin spraying when disease is reported in the area. Repeat at 7 to 10 day intervals depending on disease severity. When disease is severe on large plants use 7 pounds at 7 day intervals. Use sufficient spray volume to obtain thorough coverage.

SUGAR BEETS — *Cercospora Leaf Spot.* Use 4 pounds per acre and begin spraying when symptoms first appear. Repeat at 7 to 10 day intervals for as long as disease threatens. Use sufficient spray volume to obtain thorough coverage of large plants.

TOMATOES — *Early Blight, Late Blight, Septoria Leaf Spot.* See Potatoes.

Kauritil® is a registered trademark and product of BASF Aktiengesellschaft, Ludwigshafen am Rhein, Federal Republic of Germany.