



Directions For Use

Apply only to corn to be used for animal feed.

Do not dilute this product. Apply undiluted product with a low pressure sprayer system with nozzles arranged so as to provide adequate and uniform coverage of kernels as they are augered into conventional storage bins.

NOTE: POCKETS OF UNTREATED CORN MAY DEVELOP MOLD GROWTH WHICH CAN SPREAD AND EVENTUALLY CAUSE SPOILAGE OF THE ENTIRE BIN. Stop auger immediately if spraying system fails. Make sure spray system is calibrated to deliver the required amount of spray, and that it provides uniform coverage of the kernels.

FIELD-SHELLED CORN Determine moisture content of corn and apply 3-9 fluid ounces of LUPROSIL (undiluted) per bushel according to the following chart and the directions given above.

Percent Moisture At Harvest	Fluid Ounces Per Bushel for Indicated Storage Period		
	To April 1	To July 1	To September 1
21 or less	3	4	6
22-25	4	6	8
26-30	6	8	9

After treatment follow normal aeration procedures to prevent moisture condensation and migration in storage.

Use Precautions

READ ENTIRE LABEL USE STRICTLY ACCORDING TO PRECAUTIONS AND DIRECTIONS FOR USE AND IN ACCORDANCE WITH SPECIFIC RECOMMENDATIONS OF STATE AGRICULTURAL EXPERIMENT STATIONS AND EXTENSION SERVICE SPECIALISTS.

Luprosil™

Preservative For Field-Shelled Corn

For the prevention of storage loss due to molds in high-moisture corn

Active Ingredient:
Propionic Acid 100%

CAUTION

KEEP OUT OF REACH OF CHILDREN
IRRITATING TO EYES, SKIN AND MUCOUS MEMBRANES.
AVOID CONTACT WITH EYES, SKIN OR CLOTHING.

In case of contact, flush eyes with plenty of water for at least 15 minutes and get medical attention promptly. For skin, remove contaminated clothing at once and wash skin with soap and water. Launder clothing before reuse.

For other Use Precautions, see side panel.
Destroy empty container. Do not reuse.

Net Contents 5 Gallons

BASF Wyandotte Corporation
Parsippany, New Jersey

Protective gloves and goggles should be worn when using the undiluted product.

When using in an enclosed space, provide good ventilation or use an efficient fan to remove fumes from the work area.

This product is corrosive in concentrated form. Rinse spray equipment thoroughly after use to prevent corrosion. Spillage of the product should be flushed away with plenty of water or neutralized with agricultural lime, washing soda or sodium bicarbonate.

GERMINATION OF SEED GRAIN MAY BE REDUCED BY TREATMENT AND PERSISTENT ODORS FROM CHEMICAL TREATMENT MAY PRODUCE OFF-FLAVORS IN FOODS. THEREFORE DO NOT USE ON GRAINS OTHER THAN THOSE INTENDED FOR ANIMAL FEED.

Warranty and Disclaimer

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use. The exclusive remedy of user or buyer and the limit of liability of BASF Wyandotte Corporation is the purchase price paid for the quantity of product involved.

EPA Reg No 7969-33

1172-US

LUPROSIL™ is a trademark and product of BASF AG
Ludwigshafen am Rhein, Federal Republic of Germany