

BASF



ACCEPTED
July 7, 1972
UNDER THE PATENT OFFICE
NOTICE TO APPLICANTS
IN LONDON LONDON REGISTER
NO. *296A-35*

Basanite™ Five

BASF



Basanite Five

ly pre-mix the proper amounts of BASANITE FIVE and oil. With vigorous agitation, add this pre-mix to the water. Add the remaining required amount of water. Agitation must be maintained during mixing and application to keep the spray uniform. mixed. FOR OIL SPRAYS, drain all water from sprayer tank, lines and pump. Add a small amount of oil to the tank, circulate through the pump system, then completely drain the system and discard the liquid. This is to remove any water remaining in the sprayer. Add desired amount of oil to the tank. With agitation, add the proper amount of BASANITE FIVE. Moderate agitation will maintain a uniform spray mix. **NOTE:** Small amounts of water in BASANITE FIVE—oil sprays may result in extremely thick gel-like emulsions which are difficult to apply. This problem can be avoided by carefully following the mixing instructions outlined.

GENERAL WEED CONTROL: Use 2 to 3 pints of BASANITE FIVE in 5 to 30 gallons of oil made up to 100 gallons with water. Spray weed growth thoroughly.

CONTACT PRE-EMERGENCE WEED CONTROL: BASANITE FIVE may be used to kill tiny weeds that emerge before certain crops. When practical, fitting the land several days before seeding will result in more weeds being up at the time of spraying. For use on land planted to the large seeded crops, beans, corn, cucurbits, potatoes and gladiolus, spray at least one day before first emergence of the crop using 1-1/2 pints of BASANITE FIVE per acre. Mix this amount with 2 gallons of oil and 30 gallons of water and apply as a fine spray to insure wetting of the weeds.

CROP USES

GRAPES AND BUSH FRUITS: Apply spray containing BASANITE FIVE when weeds are growing well but before they are 6 inches high. **ON GRAPES,** applications can be made up to 3 or 4 weeks after


oughly wet all weed foliage using approximately 100 gallons per acre. The higher dosage and larger amounts of oil are needed where grassy weeds are the major problem. **NOTE: Do not graze treated areas or use treated foliage for feed or forage.**

ALFALFA, TREFLOIL, CLOVER, PEAS AND SOY-BEANS GROWN FOR SEED: (Pre-harvest spraying to facilitate harvest.) Apply 3 to 6 days before harvest. For airplane application, use 2 to 3 pints of BASANITE FIVE in 4 to 10 gallons of diesel or fuel oil per acre. For ground application, use 2 to 3 pints of BASANITE FIVE per acre. This may be mixed with 8 to 15 gallons of oil for low-volume sprays or with 5 to 15 gallons of oil plus 25 to 40 gallons of water for high-volume sprays. The higher rates and volumes are suggested for use during cool, cloudy weather and where there is heavy foliage growth. **NOTE: Do not graze treated areas nor feed forage or seed from treated crops to livestock.**

POTATO VINES: (Top kill for harvest aid.) Spray 10 to 20 days before harvest. The concentration of the spray and the amount to use per acre are determined by variety and vigor of the vines, degree of kill desired and weather conditions at the time of application. Use the higher rates during cool, cloudy weather or where foliage growth is especially heavy. For LOW VOLUME SPRAYS, use 2 to 3 pints of BASANITE FIVE in 5 to 10 gallons of diesel or fuel oil per acre, applying by either air or ground equipment. For MEDIUM VOLUME SPRAYS, use 2 to 4 pints of BASANITE FIVE per acre in 5 gallons of fuel oil and 25 to 40 gallons of water, depending on density of vine growth. Adjustment of boom and nozzles to give coverage of all parts of vines is essential. **NOTE:** Browning of the vascular ring of potato tubers sometimes occurs after the use of vine killers. Do not spray exposed tubers nor graze treated areas.

the vines will need to be increased proportionately. These treatments will materially aid in the control of weeds. **USE PRECAUTION: Do not spray entire vines. Do not apply to young replants in mature replantings or to plants in "baby" yards. Do not apply within 14 days of harvest.**

IMPORTANT: Do not store at temperatures below 20°F. If stored for extended periods below 20°F, bring the contents of the container up to 32°F and agitate by rolling.



This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Birds and other wildlife in treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

Do not apply this product in any manner not specified on this label.

NOTE: Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by breaking or puncturing and burying with wastes.

WARRANTY AND DISCLAIMER

Some warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. **But neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.** The exclusive remedy of user or buyer and the limit of liability of BASF Wyandotte Corporation is the purchase price paid for the quantity of product involved.

0472-00-La24-US

EPA Reg. 7969-35

BASANITE™ is a trademark of BASF Wyandotte Corporation, Parsippany, New Jersey.