

February 16, 1984

TIME AND RATE OF APPLICATION

Apply the Basagran + Blazer 2L tank mix broadcast when weeds are actively growing and before weeds reach the maximum size listed in the Tank Mix Recommendation Table. Such applications generally correspond to the peanut growth stages of bunching to pegging.

Delay in application which permits weeds to exceed the maximum size stated will result in inadequate control.

Choose the rate of Basagran (1½ or 2 pints per acre) according to the size and species of the weeds to be controlled with Basagran alone (see Basagran label). Then add Blazer 2L at the rate of 1 pint per acre, if needed, to control the additional weed species, up to the maximum size, as shown in the Tank Mix Recommendation Table. Blazer 2L may also be included in the tank mix at a rate up to 2 pints per acre; however, this will increase the severity and/or frequency with which peanut injury is observed.

BASAGRAN + BLAZER 2L TANK MIX RECOMMENDATION TABLE

Tank Mix Rates of Application		
Basagran: 1½ or 2 pts./A depending on stage of growth of weeds controlled by Basagran alone.	Blazer 2L: 1 pt./A	
Additional Weeds Controlled With Tank Mix		
Species	Maximum Height	Number of Leaves
Common Ragweed (<i>Ambrosia artemisiifolia</i>)	6"	up to 10
Redroot Pigweed (<i>Amaranthus retroflexus</i>)	3"	up to 6
Smooth Pigweed (<i>Amaranthus hybridis</i>)	3"	up to 6
Black Nightshade (<i>Solanum nigrum</i>)	2"	up to 6
Sesbania (<i>Sesbania exaltata</i>)	6"	up to 4 pinnate
Morningglories (<i>Ipomoea</i> spp.)	4"	up to 4
Crotalaria (<i>Crotalaria spectabilis</i>)	6"	up to 6

WATER VOLUME AND SPRAY PRESSURE

Ground Equipment

Use a minimum of 20 gallons of total spray mixture per acre (broadcast basis) and 40 psi pressure with standard high pressure hollow cone or flat fan nozzles spaced 20 inches apart.

Air Equipment

Use a minimum of 10 gallons of water per acre.

Mixing

Fill the spray tank half full with water and add the recommended amount of Basagran and Blazer 2L while the agitator is running. Then add the remaining quantity of water.

RESTRICTIONS AND LIMITATIONS (PARTIAL LIST)

Do not apply tank mix if peanuts show injury (leaf phytotoxicity and/or plant stunting) produced by any other prior herbicide treatment because this injury may be enhanced and/or prolonged.

Do not apply tank mix during prolonged periods of drought or during unseasonably cold weather, as unsatisfactory weed control may result.

Do not apply tank mix to peanuts that have been subjected to stress conditions such as hail damage, flooding, drought, or unseasonably cold or widely fluctuating temperatures because injury may result.

Do not add a surfactant or oil concentrate except where specifically recommended.

Blazer is a trademark of the Rohm and Haas Company



BASF Wyandotte Corporation
Parsippany, New Jersey 07054

February 16, 1984

Supplemental Labeling

Basagran

herbicide

**TANK MIX WITH BLAZER® 2L HERBICIDE FOR
POSTEMERGENCE APPLICATION IN PEANUTS
(EXCEPT TEXAS AND OKLAHOMA)**

BASAGRAN (EPA Reg. No. 7980-45)

BLAZER 2L (EPA Reg. No. 707-15)

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on both EPA-registered labels are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

Basagran is a postemergence broadleaf weed herbicide that controls the broad spectrum of weeds listed below.

Balloonvine (<i>Cardiospermum halicacabum</i>)	Dayflower (<i>Commelina</i> spp.)	Smallflower Morningglory (<i>Jacquemontia tamnifolia</i>)
Beggarticks (<i>Bidens frondosa</i>)	Devilsclaw* (<i>Proboscidea louisianica</i>)	Spurred Anoda (<i>Anoda cristata</i>)
Bristly Starbur (<i>Acanthospermum hispidum</i>)		Tropic Croton (<i>Croton glandulosus</i>)
Cocklebur (<i>Xanthium pensylvanicum</i>)	Giant Ragweed (<i>Ambrosia trifida</i>)	Velvetleaf (<i>Abutilon theophrasti</i>)
Coffee Senna* (<i>Cassia occidentalis</i>)	Jimsonweed (<i>Datura stramonium</i>)	Wild Sunflower (<i>Helianthus annuus</i>)
Common Ragweed* (<i>Ambrosia artemisiifolia</i>)	Ladysthumb (<i>Polygonum persicaria</i>)	Yellow Nutsedge* (<i>Cyperus esculentus</i>)
Cypressvine Morningglory (<i>Ipomoea quamoclit</i>)	Pennsylvania Smartweed (<i>Polygonum pensylvanicum</i>)	
	Prickly Sida or Teaweed (<i>Sida spinosa</i>)	

*Add oil concentrate at the rate of 2 pints per acre (see Basagran product label for additional information). The addition of oil concentrate when tank mixing with Blazer 2L will probably increase the severity and/or frequency with which peanut injury is observed.

See Basagran label for specific recommendations.

The tank mix of Basagran plus Blazer 2L will also control certain weeds not controlled by Basagran alone (see Tank Mix Recommendation Table).

The tank mix is effective mainly through contact action. Therefore, weeds must be thoroughly covered with spray.

Large crop-and-weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage. Peanuts are tolerant to the tank mix of Basagran + Blazer 2L; however, under certain conditions, peanuts may burn, crinkle and bronze.

(over)

February 16, 1984

TIME AND RATE OF APPLICATION

Apply the Basagran + Blazer 2L tank mix broadcast when weeds are actively growing and before weeds reach the maximum size listed in the Tank Mix Recommendation Table. Such applications generally correspond to the peanut growth stages of bunching to pegging.

Delay in application which permits weeds to exceed the maximum size stated will result in inadequate control.

Choose the rate of Basagran (1½ or 2 pints per acre) according to the size and species of the weeds to be controlled with Basagran alone (See Basagran label). Then add Blazer 2L at the rate of 1 pint per acre, if needed, to control the additional weed species, up to the maximum size, as shown in the Tank Mix Recommendation Table. Blazer 2L may also be included in the tank mix at a rate up to 2 pints per acre; however, this will increase the severity and/or frequency with which peanut injury is observed.

BASAGRAN + BLAZER 2L TANK MIX RECOMMENDATION TABLE

Tank Mix Rates of Application		
Basagran: 1½ or 2 pts./A depending on stage of growth of weeds controlled by Basagran alone	Blazer 2L: 1 pt./A	
Additional Weeds Controlled With Tank Mix		
Species	Maximum Height	Number of Leaves
Common Ragweed (<i>Ambrosia artemisiifolia</i>)	6"	up to 10
Redroot Pigweed (<i>Amaranthus retroflexus</i>)	3"	up to 6
Smooth Pigweed (<i>Amaranthus hybridis</i>)	3"	up to 6
Black Nightshade (<i>Solanum nigrum</i>)	2"	up to 6
Sesbania (<i>Sesbania exaltata</i>)	6"	up to 4 pinnate
Morningglories (<i>Ipomoea</i> spp.)	4"	up to 4
Crotalaria (<i>Crotalaria spectabilis</i>)	6"	up to 6

WATER VOLUME AND SPRAY PRESSURE

Ground Equipment

Use a minimum of 20 gallons of total spray mixture per acre (broadcast basis) and 40 psi pressure with standard high pressure hollow cone or flat fan nozzles spaced 20 inches apart.

Air Equipment

Use a minimum of 10 gallons of water per acre.

Mixing

Fill the spray tank half full with water and add the recommended amount of Basagran and Blazer 2L while the agitator is running. Then add the remaining quantity of water.

RESTRICTIONS AND LIMITATIONS (PARTIAL LIST)

Do not apply tank mix if peanuts show injury (leaf phytotoxicity and/or plant stunting) produced by any other prior herbicide treatment because this injury may be enhanced and/or prolonged.

Do not apply tank mix during prolonged periods of drought or during unseasonably cold weather, as unsatisfactory weed control may result.

Do not apply tank mix to peanuts that have been subjected to stress conditions such as hail damage, flooding, drought, or unseasonably cold or widely fluctuating temperatures because injury may result.

Do not add a surfactant or oil concentrate except where specifically recommended.

Blazer is a trademark of the Rohm and Haas Company.



BASF Wyandotte Corporation
Parsippany, New Jersey 07054

February 16, 1984

Supplemental Labeling

Basagran[®]

herbicide

TANK MIX WITH BLAZER[®] 2L HERBICIDE FOR POSTEMERGENCE APPLICATION IN PEANUTS (EXCEPT TEXAS AND OKLAHOMA)

BASAGRAN (EPA Reg. No. 7988-45)

BLAZER 2 (EPA Reg. No. 707-150)

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on both EPA-registered labels are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

Basagran is a postemergence broadleaf weed herbicide that controls the broad spectrum of weeds listed below.

Balloonvine
(*Cardiospermum halicacabum*)

Beggarticks
(*Bidens frondosa*)

Bristly Starbur
(*Acanthospermum hispidum*)

Cocklebur
(*Xanthium pensylvanicum*)

Coffee Senna*
(*Cassia occidentalis*)

Common Ragweed*
(*Ambrosia artemisiifolia*)

Cypressvine Morningglory
(*Ipomoea quamoclit*)

Dayflower
(*Commelina* spp.)

Devilscraw*
(*Proboscidea louisianica*)

Giant Ragweed
(*Ambrosia trifida*)

Jimsonweed
(*Datura stramonium*)

Ladysthumb
(*Polygonum persicaria*)

Pennsylvania Smartweed
(*Polygonum pensylvanicum*)

Prickly Sida or Teaweed
(*Sida spinosa*)

Smallflower Morningglory
(*Jacquemontia tamnifolia*)

Spurred Anoda
(*Anoda cristata*)

Tropic Croton
(*Croton glandulosus*)

Velvetleaf
(*Abutilon theophrasti*)

Wild Sunflower
(*Helianthus annuus*)

Yellow Nutsedge*
(*Cyperus esculentus*)

*Add oil concentrate at the rate of 2 pints per acre (see Basagran product label for additional information). The addition of oil concentrate when tank mixing with Blazer 2L will probably increase the severity and/or frequency with which peanut injury is observed.

See Basagran label for specific recommendations.

The tank mix of Basagran plus Blazer 2L will also control certain weeds not controlled by Basagran alone (see Tank Mix Recommendation Table).

The tank mix is effective mainly through contact action. Therefore, weeds must be thoroughly covered with spray.

Large crop-and-weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage. Peanuts are tolerant to the tank mix of Basagran + Blazer 2L; however, under certain conditions, peanuts may burn, crinkle and bronze.

(over)