

**BASAGRAN^R Herbicide
SUPPLEMENTAL LABEL**

Tank mix with Cobra^R herbicide for postemergence use in soybeans

**Basagran (EPA Registration No. 7969-45)
Cobra (EPA Registration No. 239-2585)**

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on both EPA registered labels are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

DIRECTIONS FOR USE

It is a violation of Federal Law to use these products in a manner inconsistent with approved labeling.

GENERAL INFORMATION


The tank mix of Basagran herbicide plus Cobra will also control certain weeds not controlled by Basagran herbicide alone (See Tank Mix Recommendation Table).

The tank mix is effective mainly through contact action. Therefore, weeds must be thoroughly covered with spray.

Large crop-and weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage. Soybeans are tolerant to the tank mix of Basagran + Cobra; However, under certain conditions, soybean foliage may be affected.

TIME AND RATE OF APPLICATION

The rates of application and weed sizes for the use of this tank mix are given in the Rate and Time of Application Recommendation Table. Applications of this tank mix should be made when weeds are small and actively growing and before weeds reach the maximum size listed in the application table. Such applications should be applied within 14 to 28 days after planting. Soybeans are tolerant to this tank mix at all stages of growth; however, bronzing, speckling or burning of leaves may occur under conditions of high temperatures or humidity. Soybean plants will generally outgrow this condition within 10-14 days.



BEST AVAILABLE COPY

RESTRICTIONS AND LIMITATIONS (Partial List)

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

Do not apply the Basagran + Cobra Tank Mix less than 90 days before harvesting soybeans.

Do not graze animals on green forage or stubble. Do not utilize hay or straw for animal feed or bedding.

Do not apply Basagran + Cobra Tank Mix if rain is expected within 8 hours of application or unsatisfactory weed control may result.

Make no more than one application of this tank mix in one season.

BEST AVAILABLE COPY

**BASAGRAM + COBRA TANK MIX - SOYBEAN
RATE AND TIME OF APPLICATION TABLE**

PRODUCT	PRODUCT RATE	WEEDS CONTROLLED/WEED SIZE			ADDITIVE (RATE)
BASAGRAM	1-2 PINTS/A	BALLOONVINE BEGGARTICKS BRISTLY STARBUR COCKLEBUR COFFEE SENNA COMMON LAMBSQUARTERS COMMON PURSLANE COMMON RAGWEED CYPRESSVINE MORNINGGLORY DAYFLOWER DEVILSCLAW GALINSOGA GIANT RAGWEED JIMSONWEED	LADYSTHUMB PENNSYLVANIA SMARTWEED PRICKLY SIDA OR TEAWEEED REDWEED SHEPHERDSPURSE SMALLFLOWER MORNINGGLORY SPURRED ANODA TROPIC CROTON VELVETLEAF VENICE MALLOW WILD BUCKWHEAT WILD MUSTARD WILD POINSETTIA WILD SUNFLOWER		OIL CONCENTRATE (*) OR 28% NITROGEN SOLUTION (1 GALLON/A) IF VELVETLEAF IS THE PRIMARY WEED TARGET AND LAMBSQUARTERS IS NOT A PROBLEM. APPLICATION BY AIR OF 28% NITROGEN IS NOT RECOMMENDED
PLUS	PLUS				
			LEAF STAGE	MAX. HEIGHT	
COBRA	10 - 12 1/2 FLUID OUNCES	COMMON RAGWEED BLACK NIGHTSHADE MORNINGGLORIES CROTALARIA SESBANIA TALL WATERHEMP REDROOT PIGWEED SMOOTH PIGWEED	UP TO 10 UP TO 6 UP TO 4 UP TO 6 UP TO 4 PINNATE UP TO 4 UP TO 4 UP TO 4	6" 2" 4" 6" 6" 3" 3" 3"	NOTE: DO NOT INCLUDE OIL CONCENTRATE WITH 28% NITROGEN SOLUTION.
*ADD OIL CONCENTRATE TO THE TANK MIX AT THE RATE OF 1/2 - 1 PINT/A					

BASCOBRA

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF Corporation (BASF) or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

BASAGRAN is a registered trademark of BASF AG.

COBRA is a registered trademark of Valent Corporation