


BASAGRAN SUPPLEMENTAL LABEL



BASAGRAN^(R) + GRAMOXONE EXTRA^(R) TANK MIX* - PEANUTS

**BASAGRAN (EPA Reg. No. 7969-45)
GRAMOXONE EXTRA (EPA Reg. No. 10182-280)**

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on both EPA-registered labels are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

Directions for Use

It is a violation of Federal law to use these products in a manner inconsistent with approved labeling.

General Information

The tank mix of BASAGRAN plus GRAMOXONE EXTRA will also control certain weeds not controlled by BASAGRAN alone (see Tank Mix Recommendation Table).

Since this tank mix is effective mainly through contact action, thorough coverage of weeds is essential for effective weed control. Large crop and weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage. Crop foliage present at application may bronze or crinkle, but the plants will soon outgrow these effects and develop normally.

Time and Rate of Application

The rates of application and weed sizes for the use of this tank mix are given in the Rate and Time Application Table. This tank mix should be applied at the ground crack stage of peanuts to control an early flush of weeds. A second application may be applied up to 28 days after ground crack stage. Do not make more than two applications of this tank mix to the same crop.

Apply the BASAGRAN + GRAMOXONE EXTRA tank mix to weeds which are actively growing and before weeds reach the maximum size listed in the Application Recommendation Table.

Application to weeds which exceed the maximum size stated may result in inadequate control.

* tank mix not applicable to California

BEST AVAILABLE COPY

BASAGRAN SUPPLEMENTAL LABEL-Peanuts

Spray Additives

Always add a nonionic surfactant containing at least 50% surface active agent at the rates listed in the Tank Mix Recommendation Table. Do not use crop oil concentrate or any other oil-based additive with this tank mix.

Water Volume and Spray Pressure

Use a minimum of 20 gallons of total spray mixture per acre (broadcast basis) and 30-50 psi pressure with standard high pressure hollow cone or flat fan nozzles spaced 20 inches apart. Use only ground equipment to apply this tank mix.

Mixing

Fill the spray tank half full with water and add the recommended amount of BASAGRAN, GRAMOXONE EXTRA and nonionic surfactant while the agitator is running. Then add the remaining quantity of water.

Restrictions and Limitation (Partial List)

Read and follow the restrictions and limitations on the labels for BASAGRAN and GRAMOXONE EXTRA. The most restrictive labeling applies in tank mixes.

Do not apply tank mix if peanuts show injury (leaf phytotoxicity and/or plant stunting) produced by any other prior herbicide treatment because this injury may be enhanced and/or prolonged.

Do not apply tank mix during prolonged periods of drought or during unseasonably cold weather, as unsatisfactory weed control may result.

Do not apply tank mix to peanuts that have been subjected to stress conditions such as hail damage, flooding, drought, or unseasonably cold or widely fluctuating temperatures because injury may result.

Avoid drift to all other crops and non-target areas, other than peanuts may be severely injured by drift.

Crops

BEST AVAILABLE COPY

BASAGRAN SUPPLEMENTAL LABEL-Peanuts

BASAGRAN + GRAMOXONE EXTRA TANK MIX - PEANUTS RATE AND TIME OF APPLICATION RECOMMENDATION TABLE

Table 19

Product	Product Rate	Weeds Controlled	Weed Growth Stages		
			Leaf Stage	Max. Height	Additive
BASAGRAN	1 pint/A	Balloornvine	2-4	2"	Use suitable non-ionic surfactant at 8 oz./
		Beggarticks	up to 6	6"	
		Bristly Starbur	up to 4	2"	
		Cocklebur	2-6*	6"	
		Coffee Senna	(up to 1 pinnate)	2"	
		Common Ragweed	up to 6	3"	100 gallons water or as directed on respective label.
		Dayflower	up to 6	4"	
		Devilsclaw	up to 6	3"	
		Giant Ragweed	up to 4	6"	
		Jimsonweed	up to 6	6"	
		Ladysthumb	up to 6	6"	
		Pennsylvania Smartweed	up to 6	6"	
		Prickly sida or Teaweed	up to 4	2"	
		Spurred Amoda	up to 6	3"	
		Tropic Croton	up to 2	2"	
		Velvetleaf	up to 4	2"	
		Wild Sunflower	up to 4	5"	
—plus—	—plus—				
GRAMOXONE EXTRA	0.41 pint/A (6.5 fl.oz./A)	Smooth Pigweed	up to 6	4"	
		Redroot Pigweed	up to 6	4"	
		Tall Waterhemp	up to 6	4"	
		Sicklepod	up to 4	4"	
		Florida Beggarweed	up to 4	4"	
		Morningglories	up to 6	4"	
		Smallflower	up to 4	3"	
		Texas Panicum	up to 2	2"	
		Crabgrass smooth	up to 2	2"	
		large	up to 2	2"	
		Goosegrass	up to 2	2"	

* Do not treat earlier than leaf stage shown and do not count cotyledon leaves.

* Do not treat earlier than leaf stage shown and do not count cotyledon leaves.

BEST AVAILABLE COPY

BASAGRAN SUPPLEMENTAL LABEL-Peanuts

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF Corporation (BASF) or the Seller. All such risks shall be assumed by the Buyer.


BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OF HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

BASAGRAN is a registered trademark of BASF AG.
GRAMOXONE EXTRA is a registered trademark of ICI Americas Inc.

1989 BASF Corporation

BEST AVAILABLE COPY

draft 4



Page 1

February 28, 1990

**BASAGRAN® Herbicide
SUPPLEMENTAL LABEL**

Tank mix with Pinnacle® herbicide for postemergence use in soybeans

BASAGRAN (EPA Registration No. 7969-45)
PINNACLE (EPA Registration No. 352-525)

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on both EPA-registered labels are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

DIRECTIONS FOR USE

It is a violation of Federal Law to use these products in a manner inconsistent with approved labeling.


GENERAL INFORMATION

The tank mix of Basagran® herbicide plus Pinnacle® will control certain weeds not controlled by Basagran or Pinnacle alone (See Tank Mix Recommendation Table).

The tank mix is effective mainly through contact action. Therefore, weeds must be thoroughly covered with spray. Large crop- and weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage.

TIME AND RATE OF APPLICATION

The rates of application and weed sizes for the use of this tank mix are given in the Rate and Time of Application Recommendation Table. Applications of this tank mix should be made when weeds are small and actively growing and before weeds reach the maximum size listed in the application table. Such applications should be applied within 14 to 28 days after planting. Soybeans are tolerant to the tank mix of Basagran + Pinnacle after the first trifoliate soybean leaf has fully expanded, however, under conditions of high temperature or humidity some leaf-bronzing or leaf-speckling of soybean foliage may occur. Soybean plants will generally outgrow this condition within 10-14 days.



WATER VOLUME AND SPRAY PRESSURE

Apply recommended rates of this tank mix as follows:

Ground Equipment: Use a minimum 20 gallons of water per acre on a broadcast basis. Use 40 psi pressure (measured at the boom, not at the pump or in the line) when using flat fan nozzles and use 40-60 psi pressure when using hollow cone nozzles. Do not use flood, whirl chamber or controlled droplet application (CDA) nozzles.

Air equipment: Use a minimum of 5 gallons of water per acre. Use 40 psi pressure when using flat fan nozzles and use 40-60 psi pressure when using hollow cone nozzles. Consult the respective labels for special directions for aerial applications.

BASAGRAN® + PINNACLE® Tank Mix - Soybeans
Rate and Time of Application Table

Product	Product Rate	Weeds Controlled	Maximum Weed Size	Additive (Rate) *
BASAGRAN	1 1/2 pint per acre	Cocklebur	6"	Nonionic
		Common Lambsquarters	4"	Surfactant
+		Jimsonweed	6"	at 0.125% v/v
		Ladysthumb	6"	(1/5 pint/20gal)
PINNACLE	1/4 ounce per acre	Penn. smartweed	6"	
		Prickly sida/Teaweed	3"	+
		Redroot pigweed	8"	UAN solution**
		Smooth pigweed	8"	at 4% v/v
		Tall waterhemp	8"	(3.2qt/20gal)
		Velvetleaf	5"	
		Venice mallow	2"	
		Wild buckwheat	3"	
		Wild mustard	4"	
		Wild sunflower	5"	

* Under dry conditions a nonionic surfactant of 0.25% v/v (2/5 or 0.4 pint per 20gal) or a crop oil concentrate at 0.5% v/v (4/5 or 0.8 pint per 20gal) may be used to increase weed control. The use of crop oil concentrate or high rates of surfactant may increase injury of soybeans.

** UAN (Urea Ammonium Nitrate) solution is referred to as 28% to 32% nitrogen solution.

RESTRICTIONS AND LIMITATIONS (Partial List)

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

Do not apply the tank mix of Basagran + Pinnacle within 60 days of harvesting soybeans.

Do not graze animals on green forage or stubble. Do not utilize hay or straw for animal feed or bedding.

Do not apply the tank mix of Basagran + Pinnacle if rain is expected within 8 hours of application or unsatisfactory weed control may result.

Do not apply this tank mix through any type of irrigation system.

Do not cultivate within seven days before or after application of this tank mix.

Do not allow spray from either ground or aerial equipment to drift onto adjacent crops or land, as injury to other plants may occur. Consult the respective labels for details.

Thoroughly clean sprayer immediately after spraying. See label for Pinnacle "Sprayer Cleanup".

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF Corporation (BASF) or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

BASAGRAN is a registered trademark of BASF AG.

PINNACLE is a registered trademark of E. I. du Pont De Nemours & Co. (Inc.)