

Pin 5

7969-45

ACCEPTED

MAY 28 1972

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 7969-45

Supplemental Labeling

Basagran® herbicide

Tank mix with Poast Plus® herbicide for postemergence use in soybeans

BASAGRAN (EPA Registration No. 7969-45)

POAST PLUS (EPA Registration No. 7969-88)

All applicable directions, restrictions, precautions and conditions of sale and warranty on both EPA registered labels are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

Directions for use

It is a violation of Federal law to use these products in a manner inconsistent with approved labeling.

General Information

Basagran® herbicide and Poast Plus® herbicide may be tank mixed for postemergence control of broadleaf and grass weeds in soybeans. This tank mix is effective mainly through contact action. Therefore, weeds must be thoroughly covered with spray. Large crop and weed leaf canopies shelter smaller weeds and prevent adequate spray coverage.

Time and rate of application

Rates of application and weed sizes for this tank mix are given in Table 1. Apply when weeds are small and actively growing and before weeds reach the maximum size listed. Soybeans are tolerant to the tank mix of Basagran and POAST PLUS at all stages of growth; however, under conditions of high temperature or humidity, some leaf-bronzing or leaf-speckling may occur. Soybeans generally outgrow this condition within 10-14 days after application.

Separate applications of Basagran and POAST PLUS should be made if: a) all weeds to be controlled are not at the correct growth stage for treatment at the same time, or b) grasses to be controlled include rhizome johnsongrass, quackgrass, bermudagrass, wirestem muhly, shattercane, volunteer cereals, wild oats, red rice, or itchgrass. Refer to either the Basagran or POAST PLUS label for instructions for separate applications.

Water volume and spray pressure

Apply recommended rates of this tank mix as follows:

2.84

Ground equipment:

Use a minimum of 10 gallons of water per acre (broadcast basis) and a minimum of 40 psi pressure (measured at the boom, not at the pump or in the line). Use standard high pressure hollow cone or flat fan nozzles spaced 20 inches apart. Do not use flood or whirl chamber nozzles.

Air equipment:

Use a minimum of 5 gallons of total spray solution per acre.

Mixing

Fill tank of a thoroughly clean sprayer half to two-thirds full with clean water. Start agitation and add the recommended amount of product in the following order: Basagran, urea ammonium nitrate (UAN) or ammonium sulfate (AMS), oil concentrate or Dash® spray adjuvant, and allow to mix thoroughly. Add recommended amount of POAST PLUS and remaining volume of water. Maintain constant agitation during application.

Additives

At the low rate of POAST PLUS (24 fl. oz.) the additives, DASH plus UAN or AMS must be used. At the higher rate of POAST PLUS (36 fl. oz.), either DASH or oil concentrate must be used. UAN or AMS may also be added.

Restrictions and limitations (partial list)

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

Do not apply the tank mix of Basagran plus Poast Plus within 90 days of soybean harvest.

Do not graze treated soybean fields and do not feed treated soybean forage (green succulent) or ensilage to livestock. Treated soybean hay may be fed.

Conditions of sale and warranty

BEST AVAILABLE COPY

The Directions for use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF

2064
Corporation (BASF) or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions of use, subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Basagran is a registered trademark of BASF AG.
Poast Plus and Dash are a registered trademarks of BASF Corporation.

©1992 BASF Corporation

BGLABEL

4084

Table 1

BASAGRAN + POAST PLUS Tank Mix - Soybeans
Rate and Time of Application

Product	Product Rate per Acre	Weeds Controlled/Weed Size				Additive (Rate/Acre)	
		Broadleaves and Sedge				Dash or Oil Concentrate	UAN Solution or Ammonium Sulfate
Basagran	1-2 pts./A according to weed species and size (see label for Basagran).	Balloonvine		Ladysthumb			
		Beggarticks		Pennsylvania Smartweed			
		Bristly Starbur		Prickly sida or Teaweed			
		Canada Thistle*		Redweed			
		Cocklebur		Shepherdspurse			
		Coffee Senna		Smallflower Morningglory			
		Common Lambsquarters		Spurred Aster			
		Common Purslane		Tropic Croton			
		Common Ragweed		Velvetleaf			
		Cypressvine Morningglory		Venice Mallow			
		Dayflower		Wild Buckwheat			
		Devilsclaw		Wild Mustard			
		Galinsoga		Wild Poinsettia			
		Giant Ragweed		Wild Sunflower			
plus	plus	Jimsonweed		Yellow Nutsedge			
Poast Plus	24 fl. oz.	Annual Grasses **				Dash only	1/2-1 gallon
		Wild Proso Millet***	4-10"	Green Foxtail	3-8"		plus
		Fall Panicum	3-8"	Witchgrass	3-8"	2 pts.	UAN or
		Giant Foxtail	3-8"	Woolly Cupgrass	3-8"		2-1/2 lbs.
	36 fl. oz.			Volunteer Corn	1-12"		AMS
		Barnyardgrass	3-8"	Junglerice	3-8"	Dash	1/2 - 1 gallon
		Broadleaf Signalgrass	3-8"	Red Sprangletop	3-8"	or oil	UAN or
		Yellow Foxtail	3-8"	Texas Panicum	3-8"	concentrate	2-1/2 lbs.
		Seedling Johnsongrass		Goosegrass	3-6"	2 pts.	
			3-8"	Large Crabgrass	3-6"		plus
				Smooth Crabgrass	3-6"		AMS may be added to this tank mixture.

* Requires two applications of Basagran in accordance with the label for control.

** Tank mix does not control rhizome johnsongrass, quackgrass, bermudagrass, wirestem muhly, shattercane, volunteer cereals, wild oats, red rice, or itchgrass.

*** For control of wild proso millet only, include Poast Plus in the tank mix at 18 fluid ounces/A.

BEST AVAILABLE COPY