

Basagran[®]

herbicide

For use on succulent peas

EPA Reg. No. 7969-45

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA-registered product label must be followed. This labeling must be in the possession of the user at the time of herbicide application.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Directions For Use For Controlling Canada Thistle in Succulent Peas

Apply 2 pints of **Basagran**[®] herbicide per acre when plants are from 8 inches tall to bud stage. Make a second application at the same rate 7-10 days later.

Peas are tolerant to **Basagran** after 3 pairs of leaves (or 4 nodes) are present. Pea injury can be very pronounced.

Even at tolerant stages, yellowing, bronzing, speckling or burning of leaves may occur under certain conditions (see **Restrictions and Limitations**). This temporary injury is generally outgrown without delaying podset or maturity or reducing yield.

Tolerant pea types are garden peas, English peas and southern peas.

Restrictions and Limitations (partial list)

Do not apply more than 2 pints of **Basagran** (1.0 pounds of a.i.) per acre per application.

Do not apply more than a total of 4 pints of **Basagran** per acre in one season.

Do not apply more than a total of 2.0 pounds of bentazon a. i. (from all sources) per acre, per calendar year.

Do not apply **Basagran** to succulent peas within 10 days of harvest. In California, do not apply **Basagran** to succulent peas within 30 days of harvest.

Do not apply **Basagran** to pea fields until peas have at least 3 pairs of leaves (or 4 nodes) because severe crop damage may occur.

Do not apply **Basagran** to peas under stress from root rot.

Do not apply **Basagran** when peas are in bloom.

Do not add oil to Basagran for use on peas except for use in the Pacific Northwest (PNW).

Conditions of Sale and Warranty


The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product.

Crop injury, ineffectiveness or other unintended consequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer. BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

Directions For Use, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Basagran is a registered trademark of BASF AG.

© 2000 BASF Corporation
NVA 2000-04-004-0024



BASF Corporation
Agricultural Products
P.O. Box 10528
Research Triangle Park, NC 27709

BASF