

7969-45

9/27/2001

1/6

SEP 27 2001

Mr. Edward G. Jordan, Ph.D.
BASF Corporation
P.O. Box 13528
Research Triangle Park, NC 27709-3528

Dear Dr. Jordan:

SUBJECT: Basagran Herbicide On Flax - New Use
Supplemental Label
EPA Registration No. 7969-45
Your Submission Dated: September 20, 2001

The supplemental label referred to above, submitted in connection with registration under the Federal Insecticide, fungicide, and Rodenticide Act, as amended, is acceptable. The amended supplemental label should be incorporated into the master label of this product, at the next label printing. A stamped copy is enclosed for your record. Please submit one copy of your final printed label before you release this product for shipment.

The Health Effects Division's (HED) Risk Assessment identified multiresidue testing and field rotational crop studies as data gaps. As previously indicated in our March 14, 2000, letter to you (reducing the PHI in peas), these data gaps are being handled under the RED follow-up process.

As indicated in HED's review, there is a Codex MRL of 0.1 ppm for bentazon and its metabolites in/on linseed. In the interest of harmonization of the U.S. tolerance and the Codex MRL, you are encouraged to submit the residue data and Good Agricultural Practices (GAP) to Codex now that the registration and tolerance of 1.0 ppm for flax have been approved in the U.S.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

Supplemental Labeling

Not for use in California or Florida

wjm 3-8-01 NVA 2001-04-004-0102

Basagran®

herbicide

For postemergence use in flax

Basagran EPA Reg. No 7969-45

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the Precautionary Statement, Environmental Hazards, Agricultural Use Requirements, Storage and Disposal statements, and Conditions of Sale and Warranty.

General Information

Basagran® herbicide is intended for selective post-emergence control of certain broadleaf weeds and sedges. (See Application Rate Table for specific weeds.) Basagran does not control grasses. Basagran is effective mainly through contact action; therefore, weeds must be thoroughly covered with spray. Large weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage. Flax is tolerant to Basagran, however, some leaf speckling and leaf bronzing may occur under certain conditions. (See Restrictions and Limitations.)

Timing of Application

Apply Basagran early postemergence as a directed spray, when weeds are small and actively growing and before weeds reach the maximum size listed in the Application Rate Table.

Early application to weeds produces the most beneficial effect on weed control (exception: yellow nutsedge and Canada thistle), allows use of the lower rate (depending on weed species), and makes it easier to obtain thorough spray coverage. Delay in application which permits weeds to exceed the maximum size stated will result in inadequate control. Do not cultivate or mow within five days before or after application of Basagran.


Ground Application Methods and Equipment (Broadcast)

Water Volume: Use 10-20 gallons of spray solution per broadcast acre for optimal performance.

Spray Pressure: Use a minimum of 40 psi (measured at the boom, not at the pump or in the line).

Note: When using the lower volume (i.e., 10 gallons per acre) or when crop and weed foliage is dense, use a minimum of 60 psi for best results.

Application Equipment: Use standard high-pressure



pesticide flat fan or hollow cone nozzles spaced up to 20 inches apart. Do not use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles as erratic coverage can cause inconsistent weed control. Do not use selective application equipment such as recirculating sprayers or wiper applicators. Good coverage is essential for maximum control.

Special Information for Irrigated Areas

In irrigated areas, it may be necessary to irrigate prior to treatment with Basagran herbicide to ensure that weeds are growing actively. Weeds growing under drought conditions usually are not satisfactorily controlled.

Addition of Oil Concentrate to Spray Tank

A nonphytotoxic oil concentrate (commonly referred to as oil concentrate) should be added to the spray tank for certain weed problems as recommended in the Application Rate Table.

The oil concentrate must contain either a petroleum or vegetable oil base and must meet the following criteria: 1) be nonphytotoxic, 2) contain only EPA-exempt ingredients, 3) provide good mixing quality in the jar test (see the following page), and 4) be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers which provide good mixing quality. For vegetable oil concentrates, it has been observed that highly refined vegetable oils are more satisfactory than unrefined vegetable oils.

For additional information, see Jar Test for Estimating Suitability of Oil Concentrates at the end of this section.

With the addition of oil concentrate to Basagran, a slight leaf burn of desirable plants may occur, but all new growth is normal, and crop vigor is not reduced. The potential for leaf burn is increased when relative humidity and temperature are high. A few oil concentrates have exhibited excessive leaf burn. However, a directed spray application should reduce the potential for leaf injury. Refer to your supplier of Basagran for information concerning successful local experience prior to purchasing any oil concentrate.

Rate of Oil Concentrate

Use one to two (1-2) pints in 10-20 gallons of water per acre.

4/6

Mixing/Spraying

Clean sprayer thoroughly prior to application of **Basagran**[®] herbicide, particularly if a herbicide was used with the potential to injure the crop to be sprayed with **Basagran**.

Fill tank of a thoroughly clean sprayer 1/2-2/3 with clean water. Start agitation and add **Basagran** and allow to mix thoroughly. Add oil concentrate and remaining volume of water. Maintain constant agitation during application.

Jar Test for Estimating

Suitability of Oil Concentrates

- Water supply:** Use only water from intended source and at the source temperature.
- Amount of water in jar.** Ground application: For 20 gallons per acre spray volume, use 3 1/3 cups (800 ml) of water. For other spray volumes, adjust proportionately.
- Amount of herbicide and oil concentrate to add:** Add herbicide and oil concentrate at the rate of 1 teaspoon (5 ml) for each pint of recommended label rate.
- Add components in following sequence, gently mixing between adding components:**
 - Basagran**
 - Oil concentrate.**

5. **Cap jar, invert 10 cycles, let stand for 15 minutes. Evaluate.**

6. **Evaluation:** An ideal tank mix combination will be uniform; thus, the suitability of the oil concentrate is questionable if any of the following are observed:

Free oil at the surface-film or globules.

Flocculation-fine particles which may be suspended in the liquid or found as a precipitated layer at the bottom of the jar.

Clabbering-thickening texture (coagulated) resembling yogurt or a curd-like texture as with cottage cheese.

Application Rate Table For Flax

Weeds Controlled	Application Rate Table			
	1 1/2 pints per acre		2 pints per acre	
	Leaf stage	Maximum Height	Leaf stage	Maximum Height
Balloonvine	2-4	2'	4-6	3'
Beggaricks	Up to 6	6"	6-8	6"
Bristly Starbur	Up to 4	2'	4-6	3'
Cocklebur	2-6"	6"	6-10	10"
Coffee Senna	—	—	Up to 1 pinnate**	2'
Common Lambquarters [†]	Up to 6	1 1/2'	4-8**	2'
Common Purslane	Up to 4	1'	4-6	2'
Common Ragweed	—	—	4-6**	3'
Dayflower	Up to 6	4'	6-10	8"
Devilsclaw	—	—	Up to 6**	3'
Gainefoga	—	—	Cotyledon to 6**	2'
Giant Ragweed**	—	—	Up to 4	6"
Jimsonweed	Up to 6	6"	6-10	10"
Ladysthumb	Up to 6	6"	6-10	10"
Marshelder	Up to 4	2'	Up to 8	4'
Pennsylvania Smartweed	Up to 6	6"	6-10	10"
Prickly Side or Teaweed	Up to 6	3'	6-8	4'
Redweed	4-8	6"	6-10	8"
Sesbania	—	—	3-5**	3'
Shepherdspurse**	Up to 6	4'	6-10	8"
Spurred Anoda	Up to 6	3'	6-8	4'
Tropic Croton	Up to 2	2"	2-4	4"
Velvetleaf	Up to 6**	5"	4-6**	6"
Vernice Mallow	Up to 6	2'	6-10	4'
Wild Buckwheat	Up to 4	3"	4-6	5"
Wild Mustard	Up to 6	4"	6-10	8"
Wild Poinsettia	2-4**	4"	4-8**	6"
Wild Sunflower	Up to 4	5"	4-6	8"

For additional weeds, See Special Directions.

* Do not treat earlier than leaf stage shown, and do not count cotyledon leaves.

** Add oil concentrate according to the Directions For Use.

† Control may be partial or inconsistent.

** If after the first application, a second weed flush develops, re-treat according to this rate table.

** Do not treat rosette before the seed stalk appears.

Special Directions for Other Weed Problems

<p>Canada Thistle Apply 2 pints of Basagran per acre when plants are between 8 inches tall and the bud stage. Make a second application at the same rate 7-10 days later. Add oil concentrate to the spray solution of Basagran/water for each application, according to the Directions For Use.</p>
<p>Yellow Nutsedge Two applications are preferred for best results. Apply 1 1/2-2 pints of Basagran per acre when plants are 6-8 inches tall. If needed, make a second application at the same rate 7-10 days later. Add oil concentrate to the spray solution of Basagran/water for each application, according to the Directions For Use.</p>
<p>Musk Thistle Apply 2 pints of Basagran per acre when plants are in the rosette stage no larger than 10 inches in diameter. Make a second application at the same rate if needed. Add oil concentrate to the spray solution of Basagran/water for each application, according to the Directions For Use.</p>

Restrictions and Limitations

Do not apply **Basagran**[®] herbicide to crops listed on this label that have been subject to stress conditions such as hail damage, flooding, drought, injury from other herbicides, or widely fluctuating temperatures, as crop injury may result.

Do not apply **Basagran** during prolonged periods of drought or during unseasonably cold weather as unsatisfactory weed control may result. Rainfall or overhead irrigation soon after application (within 8 hours) may nullify the effectiveness of **Basagran**.

Do not apply more than 2 pints of **Basagran** (1.0 pounds of a. i.) per acre per application.

Do not apply more than a total of 4 pints of **Basagran** per acre per calendar year.

Do not apply more than a total of 2.0 pounds of bentazon a. i. (from all sources) per acre per calendar year.

Do not apply within 45 days of harvest.

Do not apply this product through any type of irrigation system.

Clean sprayer thoroughly prior to application of **Basagran**, particularly if a herbicide was used which has the potential to injure flax.

Environmental Hazards

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not contaminate water when disposing of equipment wash waters or rinsate.

Bentazon, which is present in this product, is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Notice: It is a violation of Federal law to use any pesticide in a manner that results in the death of an endangered species or in adverse modification of their habitat.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Do not store at less than 32° F and do not allow product to freeze.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

- **Plastic Containers:** Triple rinse the container (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke. Do not re-use empty container.

5/6

6/6

Conditions of Sale and Warranty
The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing

Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Basagran is a registered trademark of BASF AG.

© 2001 BASF Corporation

NVA 2001-04-004-0102

BASF Corporation
 P.O. Box 13526
 Research Triangle Park, NC 27709

BASF