

ACCEPTED  
MAY 22 1989  
7969-53

**BASF**

# Ronilan® WP

## Fungicide

A wettable powder containing

**Active ingredient**

3-(3,5-Dichlorophenyl)-5-ethenyl-5-methyl-2,4-oxazolinedione

50%  
50%

**Inert ingredients**

EPA Reg. No. 7969-53

KEEP OUT OF REACH OF CHILDREN.

### CAUTION

Causes eye irritation. Do not get on skin, in eyes or on clothing

**Statement of practical treatment**

If contacted, flush eyes immediately with water for 15 minutes. Get medical attention. In case of contact with skin or clothing, remove contaminated clothing, wash skin thoroughly with soap and water. This product is a potential skin sensitizer. If irritation persists get medical attention.

**Environmental hazards**

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**Net contents 3 lbs.**

(2 x 1½ lb bags Do not remove bags from carton except for immediate use )

**BASF Corporation** Chemicals Division  
100 Cherry Hill Road Parsippany New Jersey 07054

**Specimen Label**

### General information

Ronilan is effective for the control of brown rot blossom and twig blight, and fruit brown rot of stonefruit including apricots, cherries, nectarines, peaches, plums, and prunes. Ronilan is a contact fungicide, therefore, thorough coverage of plant parts to be protected is essential for effective disease control.

### Time and rate of application

Depending on conditions which favor disease development, one to three applications should be made during bloom for control of brown rot blossom and twig blight. One or two preharvest applications should be made for control of fruit brown rot (see table).

A rate of 1½ pounds product per acre is generally recommended. A one pound product per acre rate of Ronilan should be used only when low disease pressure can be predicted. Under high disease pressure or in conditions very favorable for disease development, use the higher treatment rates specified, multiple applications, and shorter application intervals. For large mature trees, use the higher treatment rates specified. Consistency of disease control with low recommended rates can be enhanced by 8-16 oz /100 gal of a non-ionic surfactant in tank mix with Ronilan.

### Method of application

Rates of Ronilan per 100 gallons are based on an application of 400 gallons per acre of dilute spray. Refer to the **Application Rates and Timing for Control of Brown Rot** if other than 400 gallons per acre is applied.

(Insert)

**Mixing instructions:** Thoroughly mix the required amount of Ronilan in a small volume of water. Partially fill the spray tank with clean water and begin agitation. Add the Ronilan mix to the tank and fill tank to the total volume required. Maintain agitation to keep the material in suspension and apply with properly calibrated spray equipment.

Ronilan is physically compatible with most pesticides used in a tank mix on stonefruit. To assure compatibility of Ronilan with other products, add correct proportions in a small container with water. Mix

Apply RONILAN as a spray, using sufficient water to obtain thorough coverage. Ground equipment: Apply in 50-400 gallons per acre. Air equipment: Apply in a minimum of 15 gallons per acre. Aerial application is recommended for bloom and petal fall applications only.

### Application Rates and Timing for Control of Brown Rot (*Monilinia* spp.)

Crop	Pounds of Product* per		Timing
	100 Gallons	Acre	
Apricots Cherries Nectarines Peaches Plums Prunes	0.25 to 0.5  Rate is based on 400 GPA	1.0 to 2.0	<b>Brown Rot Blossom &amp; Twig Blight</b> Apply at <b>early bloom</b> Apricots—Red Bud Peaches & Nectarines—Pink Bud Cherries—Early Popcorn Plums and Prunes—Green Tip Apply again at <b>full bloom</b> if conditions persist which favor disease development  Apply again at <b>petal fall</b> if conditions persist which favor disease development Do not apply more than 3 bloom treatments <b>Fruit Brown Rot</b> Apply when conditions favor disease development within 3 weeks of harvest. One additional application should be made 7 to 14 days later if these conditions persist. Do not apply more than 2 pre-harvest treatments Do not apply within 3 days of harvest
	Do not exceed 2 lbs product per acre per application		

\*See Time and rate of application

thoroughly and let the mixture stand for five minutes. The mixture is compatible if the combination remains mixed, or can be readily remixed.

### Restrictions and limitations for stonefruit

Do not apply Ronilan within 3 days of harvest.

Do not apply more than 2 pounds of Ronilan per acre per application.

Do not apply more than 10 pounds of Ronilan per acre per season (maximum of 6 pounds at bloom and 4 pounds preharvest).

Do not apply Ronilan during rain. Apply when conditions will permit spray to dry on the plants.

Do not apply this product through any type of irrigation system.

### Conditions of sale and warranty

The Directions for use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for use, subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

EPA Establishment No. 39578-TX-1

Ronilan is a registered trademark of BASF AG  
Made in West Germany

© 1988 BASF Corporation