

most reliable wild oat control, this product should be incorporated within 48 hours of application as described in the "FIELD PREPARATIONS" and "INCORPORATED APPLICATIONS" sections of this label.

NO-TILL APPLICATION

FOR WILD OAT CONTROL IN SPRING PLANTED CROPS IN MONTANA ONLY

This product may be applied prior to planting no-till spring wheat, durum wheat, and barley at a rate of 15 pounds per acre for all soil types.

No-till applications of this product may be made 3 weeks prior to soil freeze-up and ending before spring thaw (Average soil temperature at the 2-inch depth must be 40° F or less).

No-till applications may be made in standing stubble or to fields with surface residue. No-till applications should not be made under conditions which will not allow granule contact with the soil.

No-till applications are to be made in reduced tillage/no-till systems only. If this product is used in any other system, it must be applied and incorporated as described elsewhere in this label.

FOR FALL PLANTED CROPS

FOR USE IN COLORADO, KANSAS, NEBRASKA, OKLAHOMA AND SOUTH DAKOTA — This product can be surface-applied prior to seeding of winter wheat without incorporation when seeding is done with a hoe drill. Apply 15 pounds per acre for suppression of downy brome (*Bromus tectorum*), cheat (*Bromus secalinus*) and Japanese brome (*Bromus japonicus*).


FOR USE IN IDAHO, MONTANA, OREGON, UTAH and WASHINGTON — Apply this product at a rate of 15 pounds per acre as a surface application ahead of a Yielder™ type no-till drill to obtain suppression of wild oats and downy brome in winter wheat or barley. A flexible harrow attached to the drill or used shallowly following planting at a right angle to the drill rows may improve herbicide performance.

Yielder™ is a trademark of Yielder Drill, Inc.

843 53-000 43/53

If you have any questions concerning this Label Pamphlet or the proper application or use of Far Go herbicide, contact your local Monsanto Sales Representative.

In case of an emergency involving this product
Call Collect, day or night, (314) 694-4000



MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.



Page 1
May 23, 1988


Supplemental Labeling

Laddok^R
Herbicide

Not intended for use in California

For Postemergence Use in Residential and Non Residential Non Crop and
Non Food Crop Sites

Laddok EPA Reg. No. 7969-54



All applicable directions, restrictions, precautions and Conditions
of Sale and Warranty on the EPA registered label are to be
followed. This labeling must be in the possession of the user at
the time of herbicide application.

Directions for use

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.


General information

LADDOK selective herbicide is intended for the postemergence control
of a broad spectrum of broadleaf weeds. LADDOK does not control
grasses. LADDOK is effective mainly through contact action;
therefore, weeds must be thoroughly covered with spray. Large leaf
canopies shelter smaller weeds and prevent adequate spray coverage.
The desired plants recommended for use in this labeling are tolerant
to LADDOK when used as directed. Very slight leaf speckling of
desired plants may occur but these plants generally outgrow this
condition.

Timing of applications

Apply LADDOK early postemergence when weeds are small and actively
growing and before weeds reach the maximum size listed in the
Application Rate Table.

Early application to weeds produces the most beneficial effect on
weed control, allows use of the lower rate (depending on weed
species), and makes it easier to obtain thorough spray coverage.
Delay in application which permits weeds to exceed the maximum size
stated will result in inadequate control.



Water Volume and spray pressure.

Apply recommended rates of LADDOK as follows:

Ground equipment: Use a minimum of 20 gals. of water per broadcast acre and a minimum of 40 psi pressure (measured at the boom, not at the pump or in the line). When weed foliage is dense use up to 50 gals. of water and up to 80 psi pressure. Use standard high pressure pesticide hollow cone or flat fan nozzles for best weed coverage.

Special information on irrigation

In irrigated areas, it may be necessary to irrigate prior to treatment with LADDOK to ensure weeds are growing actively. Weeds growing under drought conditions usually are not satisfactorily controlled, so it will be necessary to irrigate these areas before application or wait until adequate rainfall has been received.

NON-FOOD USES


Turf

For use in St. Augustine grass, zoysia, and centipede grass. Apply no earlier than 15 days after sprigging or plugging.

Non-crop and Roadside


For use in switch grass and big bluestem in fields, ranges or roadside.

Applications can be made postemergence at any growth stage of the grasses if already established. Care should be taken in seedling grasses as injury may result. A small area should be tested before a general application is made.



Application Rate Table

Weeds Controlled	Application Rates for LADDOK					
	2 Pints/Acre		2 1/2 pints/Acre		3 1/2 pints/Acre	
	Leaf	Max.	Leaf	Max.	Leaf	Max.
Annuals	Stage	Height	Stage	Height	Stage	Height
Cocklebur	2-4**	3"	2-10**	8"	----	----
Common Lambsquarters	----	----	Up to 8	5"	8-12	8"
Common Ragweed	----	----	Up to 5	4"	4-7	5"
Giant Ragweed	----	----	Up to 4	4"	4-6	6"
Jimsonweed	2-4	3"	Up to 6	4"	6-10	8"
Ladysthumb	2-6	4"	Up to 10	9"	10-14	12"
Pennsylvania Smartweed	2-6	4"	Up to 10	9"	10-14	12"
Redroot Pigweed	----	----	Up to 10	6"	----	----
Smooth Pigweed	----	----	Up to 10	6"	----	----
Tall Waterhemp	----	----	Up to 8	2"	6-9	4"
Velvetleaf	2-4	3"	Up to 6	5"	6-8	8"
Venice Mallow	----	----	Up to 8	4"	----	----
Wild Buckwheat	----	----	Up to 4	3"	4-6	5"
Wild Mustard	----	----	Up to 6	4"	6-10	8"
Wild Sunflower	----	----	Up to 5	6"	4-6	8"
Beggarticks	----	----	-----	-----	Up to 6	6"
Bristly Starbur	----	----	-----	-----	Up to 4	2"
Dayflower	----	----	-----	-----	Up to 6	4"
Prickly Sida or Teaweed	----	----	-----	-----	Up to 6	3"
Spurred Anoda	----	----	-----	-----	Up to 6	3"
Annual Morningglory	----	----	Up to 4	4"	4-6	6"
Smallflower Morningglory	----	----	Up to 4	4"	4-6	6"
Perennials						
Yellow Nutsedge	See Special directions below.					
Canada Thistle	See Special directions below.					
Field Bindweed	See Special directions below.					
**Do not treat earlier than leaf stage shown and do not count cotyledon leaves.						



SPECIAL DIRECTIONS FOR OTHER WEED PROBLEMS**Canada Thistle**

Apply 3 1/2 pints of Laddok per acre when plants are from 8-10 inches tall to the bud stage. Make a follow-up application 7-10 days later using the same rate of Laddok, or 2 pints of Basagran^R per acre.


Yellow Nutsedge

A single application of Laddok at 3 1/2 pints per acre will provide suppression of yellow nutsedge.

For control of yellow nutsedge apply 3 1/2 pints of Laddok per acre when plants are 1 to 4 inches tall. Make a follow-up application 7-10 days later using the same rate of Laddok or 1 1/2 to 2 pints of Basagran per acre.

Field Bindweed

For suppression of field bindweed apply 3 1/2 pints of Laddok per acre when vines are 8 to 10 inches long.




Appendix-Scientific names

The following are scientific names for the weeds listed in this label. For specific recommendations on control of these weeds, refer to the Application Rate Table.

Broadleaf Weeds

Common Name	Scientific Name
Beggarticks	<i>Bidens frondosa</i>
Bristly Starbur	<i>Acanthospermum hispidum</i>
Canada Thistle	<i>Cirsium arvense</i>
Cocklebur	<i>Xanthium strumarium</i>
Common Lambsquarters	<i>Chenopodium album</i>
Common Ragweed	<i>Ambrosia artemisiifolia</i>
Dayflower	<i>Commelina spp.</i>
Field Bindweed	<i>Convolvulus arvensis</i>
Giant Ragweed	<i>Ambrosia trifida</i>
Jimsonweed	<i>Datura stramonium</i>
Ladysthumb	<i>Polygonum persicaria</i>
Morningglory, Annual	<i>Ipomea spp.</i>
Pennsylvania Smartweed	<i>Polygonum pennsylvanicum</i>
Prickly Sida or Teaweed	<i>Sida spinosa</i>
Redroot Pigweed	<i>Amaranthus retroflexus</i>
Morningglory, Smallflower	<i>Jacquemontia tamnifolia</i>
Smooth Pigweed	<i>Amaranthus hybridis</i>
Spurred Anoda	<i>Anoda cristata</i>
Tall Waterhemp	<i>Amaranthus tuberculatus</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Venice Mallow	<i>Hibiscus trionum</i>
Wild Buckwheat	<i>Polygonum convolvulus</i>
Wild Mustard	<i>Sinapis arvensis</i>
Wild Sunflower	<i>Helianthus annuus</i>
Yellow Nutsedge	<i>Cyperus esculentus</i>



Restrictions and limitations

Do not apply LADDOK when desired plants are under stress from prolonged cold, wet weather, poor fertility, or other factors.

Do not apply LADDOK if desired plants shows injury (leaf phytotoxicity and/or plant stunting) produced by any other prior pesticide applications, because this injury may be enhanced and/or prolonged.

Do not apply LADDOK during prolonged periods of drought or during unseasonably cold weather as unsatisfactory weed control may result.

Do not graze treated area or feed treated forage to livestock.

Rainfall or overhead irrigation soon after application (within 8 hours) may nullify the effectiveness of LADDOK.

Do not mix or apply LADDOK with any other pesticide or with fertilizer except as specifically recommended on this labeling.

Do not apply this product through any type of irrigation system.

Storage and disposal

Store above 15 degrees F.

Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Do not reuse empty container. Triple rinse container (or equivalent). Then offer for recycling or... : reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

In case of emergency

In case of large scale spillage regarding this product: Avoid contact, isolate area and keep out animals and unprotected persons. Confine spill and call:


Chemtrec 800-424-9300
BASF 201-316-3000 (Collect)

In case of medical emergency regarding this product, call:

1. Your local doctor for immediate treatment.
2. Your local poison control center (hospital).
3. BASF Corporation

Notice to User

Due to variability within species and in application techniques neither the manufacturer nor the seller has determined whether or not LADDOK can be safely used on other nonbearing food crops, ornamentals, nursery and other nonfood crops under all conditions. It is therefore recommended that the professional user should determine if LADDOK can be used safely prior to broad use.



LADDOK^R
POSTEMERGENCE FLOWABLE HERBICIDE

For Selective Postemergence Broadleaf Weed Control in
Field Corn, Seed Corn, Silage Corn, Sweet Corn and Popcorn;
and in Grain Sorghum

Active Ingredients:

Sodium salt of bentazon* (3-(Isopropyl)-1H-2,1,3-
benzothiadiazin-4(3H)-one, 2,2-dioxide).....19.14
Atrazine* (2-chloro-4-ethylamino-6-isopropylamino-
s-triazine).....17.54

Inert Ingredients.....63.44

*Equivalent to 1.66 pounds per gallon each of
bentazon and atrazine.

EPA REGISTRATION NO. 7969-54

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este
producto hasta que la etiqueta haya sido explicada ampliamente

Corrosive. Causes eye damage and skin irritation. Do not get
eyes, on skin or on clothing. Wear goggles or face shield and
rubber gloves when handling. Harmful if swallowed. May cause
allergic skin response.

If in eyes, immediately flush eyes with plenty of water. Get
medical attention. If on skin, immediately flush skin with
plenty of water. Get medical attention if irritation persists

ENVIRONMENTAL HAZARDS

Atrazine, which is present in this product, can travel (seep or
leach) through soil and can enter ground water which may be used
as drinking water. Atrazine has been found in ground water. Users
are advised not to apply Laddok to sand and loamy sand
soils where the water table (ground water) is close to the
surface and where these soils are very permeable, i.e.,
well-drained. Your local agricultural agencies can provide
further information on the type of soil in your area and the
location of ground water.

SHAKE WELL BEFORE USING
NET CONTENT - 2 1/2 Gallons

BASF CORPORATION
Parsippany, N.J. 07054

Not Intended For Use
in California

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

The following restrictions apply to use of this product after February 1, 1988.

Before using this pesticide on corn and sorghum in the counties listed below, you must obtain the PESTICIDE USE BULLETIN FOR PROTECTION OF ENDANGERED SPECIES for the county in which the product is to be used. The bulletin is available from your County Extension Agent, State Fish and Game Office, or your pesticide dealer. Use of this product in a manner inconsistent with the PESTICIDE USE BULLETIN FOR PROTECTION OF ENDANGERED SPECIES is a violation of Federal laws.

Alabama - Colbert, Greene, Jackson, Lamar, Lauderdale, Limestone, Madison, Marshall, Morgan, Pickens and Sumter

Arizona - Graham, Maricopa, Mohave, Pima, Pinal and Santa Cruz

Arkansas - Benton, Clay, Clark, Cross, Lawrence, Lee, Poinsette, Polk, Randolph, Sharp and St. Francis

Florida - Broward, Dade, Glades, and Palm Beach

Kentucky - Ballard, Butler, Edmundson, Green, Hart, Jackson, Laurel, Livingston, Marshall, McCracken, McCreary, Pulaski, Rockcastle, Taylor, Warren and Wayne

Mississippi - Claiborne, Copiah, Hinds, Itawamba, Lowndes, Monroe and Noxubee


Missouri - Barry, Benton, Camden, Christian, Dallas, Greene, Hickory, Jasper, Lawrence, Miller, Newton, Osage, Polk, St. Clair, Stone and Webster

Nevada - Clark

New Mexico - Chaves, DeBaca and Eddy

North Carolina - Edgecombe, Nash and Pitt

Ohio - Pickaway



(Endangered Species Restrictions)

Oklahoma - Delaware, Mccurtain and Pushmataha


Oregon - Lake

Tennessee - Bedford, Blount, Clairborne, Decatur, Franklin, Hancock, Hardin, Hickman, Knox, Lawrence, Lincoln, Loudon, Marshall, Maury, Meigs, Monroe, Rhea, Roane, Scott, Sequatchie, Smith, Sullivan and Wayne

Texas - Bastrop, Burleson, Comal, Harris, Hays, Jeff Davis, Pecos and Reeves

Utah - Utah and Washington

Virginia - Lee, Russell, Scott, Smyth, Tazewell, Washington and Wise



Reentry and Workers' Protection Statements

Do not apply this product in such a manner as to directly or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

Laddok selective herbicide is intended for the postemergence control of a broad spectrum of broadleaf weeds. Laddok does not control grasses. Laddok is effective mainly through contact action; therefore, weeds must be thoroughly covered with spray. Large crop-and-weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage.

Corn is tolerant to Laddok at all stages of growth. Very slight leaf speckling of corn may occur but plants generally outgrow this condition within 10 days. Corn types included are field, sweet and popcorn; and corn grown for seed or silage.

Always add UAN solution or oil concentrate according to the section entitled "Additive Information".

TIMING OF APPLICATIONS

Apply Laddok early postemergence when weeds are small and actively growing and before weeds reach the maximum size listed in the APPLICATION RATE TABLE FOR CORN AND GRAIN SORGHUM. Such applications generally correspond to the crop growth stages of one to seven leaves.

Early application to weeds produces the most beneficial effect on weed control, allows use of the lower rate (depending on weed species), and makes it easier to obtain thorough spray coverage. Delay in application which permits weeds to exceed the maximum size stated will result in inadequate control.

Do not cultivate within 5 days before or after application of Laddok in the following states: AZ, CO, CT, ID, IL, IN, IA, KS, KY, ME, MA, MI, MN, MO, MT, NB, NV, NH, NJ, NY, ND, OH, OR, PA, RI, SD, UT, VT, WA, WV, WI, WY. A cultivation 5 or more days after application may be necessary if all weeds are not controlled or if a second flush of weeds occurs.

WATER VOLUME AND SPRAY PRESSURE

Apply recommended rates of Laddok as follows:

Ground equipment: Use a minimum of 20 gals. of water per broadcast acre and a minimum of 40 psi pressure (measured at the boom, not at the pump or in the line). When crop and weed foliage is dense use up to 50 gals. of water and up to 80 psi pressure. Use standard high pressure pesticide hollow cone or flat fan nozzles spaced 20 inches apart. Do not use flood or whirl chamber nozzles.

Air equipment: Use a minimum of 5 gals. of water per acre and a maximum of 40 psi pressure. Use only diaphragm-type nozzles producing cone or fan spray patterns.

AERIAL APPLICATION-SPECIAL DIRECTIONS

To obtain uniform coverage and to avoid drift hazards, the following application equipment and practices should be used:

Nozzle height: Maximum of 10 feet above crop.

Nozzle orientation: Nozzles must be oriented so as to discharge straight back with the air stream (opposite the direction of travel of the aircraft) or at some angle between straight back and straight down.

Nozzles must be located no farther out than three-fourths the distance from the center of the aircraft to the end of the wing or rotor.

Water volume and spray pressure: See Air equipment.

Do not apply Laddok by aircraft when wind is blowing at a velocity above 10 mph. Coarse sprays (larger droplets) are less likely to drift.

Do not apply Laddok by air within 200 feet upwind of ornamental or sensitive nontarget crops such as soybeans, peanuts, cotton, sugarbeets, sunflowers or okra.

Applicator must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

SPECIAL INFORMATION FOR IRRIGATED AREAS

In irrigated areas, it may be necessary to irrigate prior to treatment with Laddok to ensure weeds are growing actively. Weeds growing under drought conditions usually are not satisfactorily controlled.

ADDITIVE INFORMATION

UAN Solution:

In corn and grain sorghum UAN (Urea Ammonium Nitrate) solution may be added to Laddok in the place of oil concentrate for most weeds. Consult the APPLICATION RATE TABLE FOR CORN AND GRAIN SORGHUM. UAN solution is commonly referred to as 28, 30 or 32% N. With the addition of UAN solution to Laddok on corn or grain sorghum, a slight leaf tip burn or yellowing and speckling may occur. New growth is normal and vigor is not reduced. Do not use brass or aluminum nozzles when spraying Laddok with UAN solution.

Rate of UAN Solution

Ground application - 1 gallon per acre.
 Air application - 1/2 gallon per acre.

Oil Concentrate:

If UAN solution is not used, a nonphytotoxic oil concentrate (commonly referred to as oil concentrate) should be added to the spray tank. Use oil concentrate if Canada thistle, yellow nutsedge or field bindweed are to be controlled. The oil concentrate must contain either a petroleum or vegetable oil base and must meet the following criteria: 1) be nonphytotoxic, 2) contain only EPA-exempt ingredients, 3) provide good mixing quality in the jar test (see below), and 4) be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers which provide good mixing quality. For vegetable oil concentrates, it has been observed that highly refined vegetable oils are more satisfactory than unrefined vegetable oils. For additional information see "Jar Test for Estimating Suitability of Oil Concentrates" at the end of this section.

Rate of Oil Concentrate:

Ground application-2 pints/acre (maximum)
 Air application-1 pint/acre (maximum)

Jar Test for Estimating Suitability of Oil Concentrates:

1. Water supply: Use only water from intended source and at the source temperature.
2. Amount of water in jar:
Ground Application-For 20 gal/A spray volume use 3 1/3 cups (800 ml) of water.
Air Application- For 5 gal/A spray volume use 5/6 cup (200 ml) of water, or
For 10 gal/A spray volume use 1 2/3 cups (400 ml) of water.
For other spray volumes, adjust proportionately to above.
3. Amount of herbicide and oil concentrate or UAN solution to add: Add herbicide and oil concentrate or UAN solution at the rate of 1 teaspoon (5 ml) for each pint of recommended label rate.
4. Add components in following sequence, gently mixing between component additions:
 1. UAN solution (if used)
 2. Laddok
 3. Oil concentrate (if used)
5. Cap jar, invert 10 cycles, let stand for 15 minutes, evaluate.
6. Evaluation: An ideal tank mix combination will be uniform; thus, the suitability of the oil concentrate is questionable if any of the following are observed:

Free oil at the surface - film or globules.

Flocculation - fine particles which may be suspended in the liquid or found as a precipitated layer at the bottom of the jar.

Clabbering - thickening texture (coagulated) resembling yogurt or a curd-like texture as with cottage cheese.

MIXING/SPRAYING

Fill tank of a thoroughly clean sprayer half to two-thirds full with clean water. Start agitation. Add UAN solution then add Laddok; allow to mix thoroughly. If UAN solution is not used add oil concentrate and remaining volume of water. Maintain constant agitation during application. Avoid allowing the mixture to stand overnight. Always clean sprayer thoroughly immediately after use by flushing the system with water and a strong detergent. Do not allow cleaning water to contaminate any streams or ponds.

APPLICATION RATE TABLE FOR CORN AND GRAIN SORGHUM

Weeds Controlled	Application Rates for Laddok					
	2 pints*		2 1/2 pints*		3 1/2 pints*	
	Leaf Stage	Max. Height	Leaf Stage	Max. Height	Leaf Stage	Max. Height
Cocklebur	2-4	3"	2-10**	8"	--	--
Common Lambsquarters	--	--	Up to 8	5"	8-12	8"
Common Ragweed	--	--	Up to 5	4"	4-7	5"
Giant Ragweed	--	--	Up to 4	4"	4-6	6"
Jimsonweed	2-4	3"	Up to 6	4"	6-10	8"
Ladythumb	2-6	4"	Up to 10	9"	10-14	12"
Penna. Smartweed	2-6	4"	Up to 10	9"	10-14	12"
Redroot Pigweed	--	--	Up to 10	6"	--	--
Smooth Pigweed	--	--	Up to 10	6"	--	--
Tall Waterhemp	--	--	Up to 8	2"	6-9	4"
Velvetleaf	2-4	3"	Up to 6	5"	6-8	8"
Venice Mallow	--	--	Up to 8	4"	--	--
Wild Buckwheat	--	--	Up to 4	3"	4-6	5"
Wild Mustard	--	--	Up to 6	4"	6-10	8"
Wild Sunflower	--	--	Up to 5	6"	4-6	8"
Beggarticks	--	--	--	--	Up to 6	6"
Bristly Starbur	--	--	--	--	Up to 4	2"
Dayflower	--	--	--	--	Up to 6	4"
Prickly Side or Teaweed	--	--	Up to 4	2"	Up to 6	3"
Spurred Anoda	--	--	--	--	Up to 6	3"
Morningglory, Annual	Up to 4	4"	4-6	6"
Smallflower Morningglory	Up to 4	4"	4-6	6"
Yellow Nutsedge	See Special Directions Below					
Canada Thistle	See Special Directions Below					
Field Bindweed	See Special Directions Below					

* Always add one gallon per acre of UAN solution or one quart per acre of oil concentrate to Laddok. See section Additive information.
 ** Do not treat earlier than leaf stage shown and do not count cotyledon leaves.

For suppression of Canada thistle apply 3 1/2 pints of Laddok per acre when plants are from 8-10 inches tall to the bud stage. Add oil concentrate for each application according to sections "Additive Information" and "Mixing/Spraying".

Yellow Nutsedge

A single application of Laddok at 3 1/2 pints per acre will provide suppression of yellow nutsedge.

Add oil concentrate for each application according to sections "Additive Information" and "Mixing/Spraying".

Field Bindweed

For suppression of field bindweed apply 3 1/2 pints of Laddok per acre when vines are 8 to 10 inches long..

Add oil concentrate for each application according to sections "Additive Information" and "Mixing/Spraying".


SORGHUM

Canada Thistle

Yellow Nutsedge

Field Bindweed

For Suppression of these weeds, apply 3 1/2 pints of Laddok per acre when Canada thistle plants are from 8-10 inches tall up to the bud stage, field bindweed vines are 8-10 inches long or Yellow nutsedge is 1-4 inches tall. Add oil concentrate according to sections "Additive Information" and "Mixing/Spraying". For best results, cultivate 7-14... days after application.



RESTRICTIONS AND LIMITATIONS

Do not apply Laddok when crop is under stress from prolonged cold, wet weather, poor fertility or other factors, or when crop is wet and succulent from recent rainfall as crop injury may occur.

Do not apply Laddok if crop shows injury (leaf phytotoxicity and/or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced and/or prolonged.

Do not apply Laddok during prolonged periods of drought or during unseasonably cold weather, as unsatisfactory weed control may result.

Seed corn producers should consult the seed corn company regarding tolerance of seed population inbred lines to Laddok.

Do not apply to grain sorghum that is heading out or blooming.


Do not make more than one application of Laddok per season.

Do not graze treated area or feed treated forage to livestock for 21 days following application.

Rainfall or overhead irrigation soon after application (within 8 hours) may nullify the effectiveness of Laddok.

Do not mix or apply Laddok with any other pesticide or with fertilizer except as specifically recommended on this labeling.

Do not apply this product through any type of irrigation system.



STORAGE AND DISPOSAL

Store above 15° F.

Do not contaminate water, food or feed by storage or disposal.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Triple rinse container (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.


IN CASE OF EMERGENCY

In Case of Large Scale Spillage Regarding This Product: Avoid contact, isolate area and keep out animals and unprotected persons. Confine spill and call:

Chemtrec 800-424-9300
BASF 201-263-3400 (Collect)

In Case of Medical Emergency Regarding This Product,
Call:

1. Your Local Doctor for Immediate Treatment.
2. Your Local Poison Control Center (Hospital).
3. BASF 201-263-3400 (Collect).



CONDITIONS OF SALE AND WARRANTY

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Basagran and Laddok are registered trademarks of BASF AG

