


## Supplemental Labeling

**POAST®  
HERBICIDE**

For use in sweet potatoes

POAST EPA Reg. No. 7969-58



All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA registered label are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

**REENTRY AND WORKERS' PROTECTION STATEMENTS**

Do not apply this product in such a manner as to directly or through drift expose workers or other persons, except those knowingly involved in the application. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must inform workers of areas or fields that may not be entered without specific protective clothing until sprays have dried, and appropriate actions to take in case accidental exposure. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is a reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: **WARNING.** Area treated with POAST herbicide on (date of application). Do not enter without appropriate protective clothing until sprays have dried. Causes eye irritation. If contacted, flush eyes immediately with water for 15 minutes. Avoid contact with skin. If contacted, flush skin immediately with plenty of water. Get medical attention if irritation persists.

In case of contact with skin or clothing, remove contaminated clothing, wash skin thoroughly with soap and water. Call physician if irritation occurs. Wash contaminated clothing before re-use.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water or wetlands (swamps, bogs, marshes or potholes). Do not contaminate water when disposing of equipment washwaters.

**ENDANGERED SPECIES CONCERNS**

The use of any pesticide in a manner that may kill or otherwise harm an endangered or threatened species or adversely modify their habitat is a violation of Federal Law.

The use of this product is contraindicated to prevent death or harm to Solano grass which occurs in Solano County, California. Before using this product in this county, you must obtain the EPA ENDANGERED SPECIES BULLETIN (EPA/ES-85-13), available from either your County Agricultural Extension Agent, the Endangered Species Specialist in the California Department of Fish and Game, or the Regional Offices of the U.S. Fish and Wildlife Service (Portland, Oregon) or the U.S. Environmental Protection Agency (San Francisco, California). **THIS BULLETIN MUST BE REVIEWED PRIOR TO PESTICIDE USE. THE USE OF THIS PRODUCT IS PROHIBITED IN THIS COUNTY UNLESS SPECIFIED OTHERWISE IN THE BULLETIN.**

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### GENERAL INFORMATION

POAST® herbicide is a selective broad spectrum postemergence herbicide for control of annual and perennial grass weeds. POAST does not control sedges or broadleaf weeds.

Essentially all grass crops such as sorghum, corn, small grains and rice, as well as ornamental grasses such as turf, are susceptible to POAST. Avoid all direct or indirect contact with any contact with any grass crop unless otherwise specified on the POAST label.

**Control Symptoms:** POAST rapidly enters the plant through the foliage and translocates throughout the plant. Control symptoms exhibited by the grass plant progress from a slowing or stopping of growth (generally within two days), to reddening of the foliage to leaf tip burn. Subsequently, burn back of the foliage occurs. These symptoms will generally be observed within three weeks depending on environmental conditions.

### APPLICATION INFORMATION

Apply POAST herbicide to actively growing grasses by ground equipment when the grasses are at the proper growth stage as specified in the annual grass and perennial grass rate recommendation tables. Do not apply to grasses under stress, such as stress due to lack of moisture, herbicide injury, mechanical injury or cold temperatures, since unsatisfactory control will result. Thorough spray coverage of grass foliage is essential. For broadcast application use standard high pressure pesticide hollow cone or flat

fan nozzles. Do not use flood or whirl chamber nozzles.

**Spray volume:** Under most conditions 10 gallons per acre of spray volume is optimal. A minimum of 5 gallons and a maximum of 20 gallons of spray solution per acre for broadcast applications may be used. In the Western States region a minimum of 10 gallons per acre is recommended. In the high plains of Texas and Oklahoma, a maximum of 10 gallons per acre is recommended.

**Spray pressure:** When using standard high pressure hollow cone or flat fan nozzles, adjust pressure to a minimum of 40 psi and a maximum of 60 psi to the nozzle. Always adjust spray pressure, spray volume and height of spray boom to ensure penetration of plant canopy and thorough coverage of grasses to be controlled. Refer to the nozzle manufacturer's directions for recommended height.

**Band applications:** Banding of POAST may be used to control grasses. Grasses which are not covered or only partly covered by spray will not be adequately controlled. When treating taller weeds, such as volunteer corn, the spray boom must be high enough to spray the top leaves and into the whorl of the plant. All recommendations are on a broadcast basis unless otherwise stated. When banding, rates of POAST, oil concentrate and water should be reduced in proportion to the area sprayed.

**Other spray equipment:** Do not use selective application equipment such as recirculating sprayers, wiper applicators, or shielded applicators. Application of POAST with control drop applicator (CDA) nozzles is not recommended due to erratic coverage which causes inconsistent weed control.

## CULTIVATION INFORMATION

Do not cultivate within 5 days prior to application of POAST or within 7 days following application.

A timely cultivation after 7 days may aid in providing season-long control. For control of Quackgrass delay cultivation until 14 to 21 days after application of POAST.

### ADDITION OF OIL CONCENTRATE

A nonphytotoxic oil concentrate (commonly referred to as oil concentrate) should always be added to the spray tank. The oil concentrate must contain either a petroleum or vegetable oil base and must meet the following criteria:

- 1) be nonphytotoxic,
- 2) contain only EPA-exempt ingredients,
- 3) provide good mixing quality in the jar test (see below), and
- 4) be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers which provide good mixing quality. For vegetable oil concentrates, it has been observed that highly refined vegetable oils are more satisfactory than unrefined vegetable oils. For additional information see Jar test for estimating suitability of oil concentrates.

**Rate of oil concentrate:**  
2 pints/acre

**Jar test for estimating suitability of oil concentrates:**

1. **Water supply:** Use only water from intended source and at the source temperature.
2. **Amount of water in jar:**  
For 20 gals./A spray volume use 3-1/3 cups (800 ml.) of water.

For 10 gals./A spray volume use 1-2/3 cups (400 ml) of water.

For 5 gals./A spray volume use 5/6 cup (200 ml) of water.


For other spray volumes, adjust proportionately to above.

3. **Amount of herbicide and oil concentrate to add:** Add herbicide and oil concentrate at the rate of 1 teaspoon (5 ml) for each pint of recommended label rate.
4. **Add components in following sequence, gently mixing between component additions:**
  1. UAN solution or ammonium sulfate (if used)
  2. Oil concentrate
  3. POAST®
5. **Cap jar, invert 10 cycles, let stand for 15 minutes, evaluate.**
6. **Evaluation:** An ideal tank mix combination will be uniform; thus, the suitability of the oil concentrate is questionable if any of the following are observed:
  1. Free oil at the surface - film or globules.
  2. Flocculation - fine particles which may be suspended in the liquid or found as a precipitated layer at the bottom of the jar.
  3. Clabbering - thickening texture (coagulated) resembling yogurt or a curd-like texture as with cottage cheese.

## MIXING/SPRAYING

Fill tank of a thoroughly clean sprayer half to two-thirds full with clean water. Start agitation and add UAN solution or ammonium sulfate (if used), and oil concentrate; allow to mix thoroughly. Add POAST and

remaining volume of water. Maintain constant agitation during application.



## RESTRICTIONS AND LIMITATIONS: Southeastern United States

- Do not apply within 30 days of harvest in the Southeastern United States.
- Do not apply more than 2-1/2 pints per acre per season in the Southeastern United States.
- Do not apply to grasses under stress, such as stress due to lack of moisture or herbicide injury, as unsatisfactory control may result.
- Do not apply if rainfall is expected within one hour following application as grass control will be unsatisfactory.
- Physical incompatibility, reduced weed control or crop injury may result from mixing POAST with other pesticides (fungicides, herbicides, insecticides or miticides), additives or fertilizers. BASF DOES NOT RECOMMEND THE USE OF POAST TANK MIXES OTHER THAN THOSE LISTED ON BASF LABELS, SUPPLEMENTAL LABELING, OR TECHNICAL BULLETINS. Local agricultural authorities may be a source of information when using other than BASF recommended combinations. Do not apply POAST with another pesticide whose label cautions against use in combination with oil adjuvants.

- 3 7 11
- Do not apply POAST through any type of irrigation system.



## RESTRICTIONS AND LIMITATIONS: Western Region States

- Do not apply within 60 days of harvest to sweet potatoes in California or Western Region states.
- Do not apply more than 5 pints per acre per season to sweet potatoes in California or Western Region states.
- Do not apply to grasses under stress, such as stress due to lack of moisture or herbicide injury, as unsatisfactory control may result.
- Do not apply if rainfall is expected within one hour following application as grass control will be unsatisfactory.
- Physical incompatibility, reduced weed control or crop injury may result from mixing POAST with other pesticides (fungicides, herbicides, insecticides or miticides), additives or fertilizers. BASF DOES NOT RECOMMEND THE USE OF POAST TANK MIXES OTHER THAN THOSE LISTED ON BASF LABELS, SUPPLEMENTAL LABELING, OR TECHNICAL BULLETINS. Local agricultural authorities may be a source of information when using other than BASF recommended combinations. Do not apply POAST with another pesticide whose label cautions against use in combination with oil adjuvants.

- Do not apply POAST through any type of irrigation system.

## RECOMMENDATION FOR GRASS CONTROL

- Apply POAST with ground equipment to actively growing grasses at sizes indicated in the following table.
- Always follow recommendations given in the Application Information section.

- Always adjust spray pressure, spray volume and height of spray boom to ensure penetration of plant canopy and thorough coverage of grasses to be controlled.

- Do not apply to drought-stressed grass or grass which has gone through an extended dry period.

- In irrigated areas, it may be necessary to irrigate prior to treatment with POAST herbicide to ensure weeds are growing actively.

- Always add 2 pints of oil concentrate per acres.

## SWEET POTATO SOUTHEASTERN UNITED STATES

GRASSES	TIME OF APPLICATION	RATE OF POAST	ADDITIVE/ OIL CONCENTRATE
Crabgrass, Large	Up to 6"	1 pint	2 pints
Goosegrass	Up to 6"		
Signalgrass, Broadleaf	Up to 8"		

## SWEET POTATO WESTERN UNITED STATES

GRASSES	TIME OF APPLICATION	RATE OF POAST PER ACRE	ADDITIVE/ OIL CONCENTRATE
Crabgrass, Large	Up to 4"	1-1/2 pints	2 pints
Goosegrass	Up to 4"		

**Procedure for cleaning spray equipment:**

Clean sprayer thoroughly prior to application of POAST®, particularly if a herbicide was used which has the potential to injure crops.

Consult the label of previously used herbicides for cleaning instructions. If no instructions are available the steps listed below are suggested for cleaning of spray equipment prior to or following applications of POAST.

**Step #1** Hose down thoroughly the inside as well as the outside of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of this rinse water.

**Step #2** Refill tank with water while adding 1 gallon household ammonia or 1 pint household dish washing detergent per 100 gallons of water. Or add a commercial sprayer cleaner according to the manufacturer's directions. Operate the pump to circulate the detergent solution through the sprayer system for 5 to 10 minutes and discharge a small amount of solution through the boom and nozzles. Let the solution stand for 24 hours.

**Step #3** Flush the detergent solution out of the spray tank through the boom.

**Step #4** Remove the nozzles and screens and flush the system with two tankfuls of water.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper

disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Triple rinse container (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**CONDITIONS OF SALE AND WARRANTY**

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User

58 11  
accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Poast is a registered trademark of BASF Ag.

:ksw


6911  
August 5, 1991

## Supplemental Labeling

# POAST® HERBICIDE

For Use in Rhubarb (IL, IN, MI, MN, OH AND WI only).

POAST EPA Reg. No. 7969-58



All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

### REENTRY AND WORKERS' PROTECTION STATEMENTS

Do not apply this product in such a manner as to directly or through drift expose workers or other persons, except those knowingly involved in the application. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must inform workers of areas or fields that may not be entered without specific protective clothing until sprays have dried, and appropriate actions to take in case of accidental exposure. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is a reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "WARNING. Area treated with POAST herbicide on (date of application). Do not enter without appropriate protective clothing until sprays have dried. If contacted, flush eyes immediately with water for 15 minutes. Get medical attention."

In case of contact with skin or clothing, remove contaminated clothing, wash skin thoroughly with soap and water. Call physician if irritation occurs. Wash contaminated clothing before re-use.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, marshes, or potholes). Do not contaminate water when disposing of equipment wash waters.

### Endangered Species Concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered or threatened species or adversely modify their habitat is a violation of Federal law.

# DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## GENERAL INFORMATION

POAST® herbicide is a selective broad spectrum postemergence herbicide for control of annual and perennial grass weeds. POAST does not control sedges or broadleaf weeds.

Since essentially all grass crops such as sorghum, corn and small grains, as well as ornamental grasses such as turf, are susceptible to POAST avoid all direct or indirect contact with any grass crop.

**Control Symptoms:** POAST rapidly enters the plant through the foliage and translocate throughout the plant. Control symptoms exhibited by the grass plant progress from a slowing or stopping of growth (generally within two days), to reddening or yellowing of the foliage to leaf tip burn. Subsequently, burn back of the foliage occurs. These symptoms will generally be observed within three weeks depending on environmental conditions.

## APPLICATION INFORMATION

Apply as a broadcast or spot spray at rates and growth stages listed in the weed tables. Do not exceed applications rates and use restrictions specified in **Restrictions and Limitations**. Apply POAST herbicide to actively growing grasses by ground equipment when the grasses are not at the proper growth stage as specified in recommendations for grass control. Do not apply to grasses under stress, such as stress due to lack of moisture, herbicide injury, mechanical injury or cold temperatures, since unsatisfactory control will result. Thorough spray

coverage of grass foliage is essential. For broadcast application use standard high pressure pesticide hollow cone or flat fan nozzles. Do not use flood or whirl chamber nozzles.

**Spray Volume:** Under most conditions 10 gallons per acre of spray volume is optimal. A minimum volume of 5 gallons and a maximum volume of 20 gallons of spray solution per acre for broadcast applications should be used.

**Spray Pressure:** When using standard high pressure hollow cone or flat fan nozzles, adjust pressure to a minimum of 40 psi and a maximum of 60 psi to the nozzle.

Always adjust the spray pressure, spray volume and height of spray boom to ensure penetration of plant canopy and thorough coverage of grasses to be controlled. Refer to the nozzle manufacturer's directions for recommended height.

**Tall Crop Application:** When a crop is 24 or more inches in height and the grasses may be below the crop canopy, drop nozzles should be used to direct more of the spray on the weeds.

**Other Spray Equipment:** Do not use selective application equipment such as recirculating sprayers, wiper applicators, or shielded applicators. Application of POAST with control drop applicator (CDA) nozzles is not recommended due to erratic coverage which causes inconsistent weed control.

## CULTIVATION INFORMATION

Do not cultivate within 5 days prior to application of POAST or within 7 days following application.

A timely cultivation after 7 days may aid in providing season-long control. For control of quackgrass, delay

cultivation until 14 to 21 days after application of POAST.

## ADDITION OF OIL CONCENTRATE

A nonphytotoxic oil concentrate (commonly referred to as oil concentrate) should always be added to the spray tank. The oil concentrate must contain either a petroleum or vegetable oil base and must meet the following criteria:

- 1) be nonphytotoxic,
- 2) contain only EPA-exempt ingredients,
- 3) provide good mixing quality in the jar test (see below), and
- 4) be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers which provide good mixing quality. For vegetable oil concentrates, it has been observed that highly refined vegetable oils are more satisfactory than unrefined vegetable oils. For additional information, see **Jar test for estimating suitability of oil concentrates**.

**Rate of Oil Concentrate:**  
2 pints/acre.

## Jar Test for Estimating Suitability of Oil Concentrates

1. **Water supply:** Use only water from intended source and at the source temperature.
2. **Amount of water in jar:** For 20 gals./A spray volume use 3-1/3 cups (833 ml) of water.

For 10 gals./A spray volume use 1-2/3 cups (400 ml) of water.

For 5 gals./A spray volume use 5/8 cup (200 ml) of water.

For other spray volumes, adjust proportionately to above.

3. **Amount of herbicide and oil concentrate to add:** Add herbicide and oil concentrate at the rate of 1 teaspoon (5 ml) for each pint of recommended rate.
4. **Add components in following sequence, gently mixing between component additions:**
  - a. UAN solution or ammonium sulfate (if used)
  - b. Oil concentrate
  - c. POAST®
5. **Cap jar, invert 10 cycles, let stand 15 minutes, evaluate.**
6. **Evaluation:** An ideal tank mix combination will be uniform; thus, the suitability of the oil concentrate is questionable if any of the following are observed.

Free oil at the surface - film or globules.

Flocculation - fine particles which may be suspended in the liquid or found as a precipitated layer at the bottom of the jar.

Clabbering - thickening texture (coagulated) resembling yogurt or a curd-like texture as with cottage cheese.

## MIXING/SPRAYING

Fill tank of a thoroughly clean sprayer half to two-thirds full with clean water. Start agitation and add UAN solution or ammonium sulfate (if used), and oil concentrate; allow to mix thoroughly. Add POAST and remaining volume of water. Maintain constant agitation during application.

## RESTRICTIONS AND LIMITATIONS

- Do not apply within 15 days of harvest.

- Do not apply more than a total of 1.5 pints per acre before harvest.
- Up to 3 additional pints of POAST may be applied after harvest.
- Do not apply more than 4-1/2 pints per season.
- Physical incompatibility, reduced weed control or crop injury may result from mixing POAST with other pesticides (fungicides, herbicides, insecticides or miticides), additives or fertilizers. BASF DOES NOT RECOMMEND THE USE OF POAST TANK MIXES OTHER THAN THOSE LISTED ON BASF LABELS, SUPPLEMENTAL LABELING, OR TECHNICAL BULLETINS. Local agricultural authorities may be a source of information when using other than BASF recommended combinations. Do not apply POAST in combination with another pesticide whose label cautions against use in combination with oil adjuvants.
- Do not apply to grasses under stress, such as stress due to lack of moisture or herbicide injury, as unsatisfactory control may result.
- Do not apply if rainfall is expected within one hour following application as grass control will be unsatisfactory.
- Do not apply POAST through any type of irrigation system.
- Always adjust spray pressure, spray volume and height of spray boom to ensure penetration of plant canopy and thorough coverage of grasses to be controlled.
- Do not apply to drought-stressed grass, or grass which has gone through an extended dry period.
- In irrigated areas it may be necessary to irrigate prior to treatment with POAST herbicide to ensure weeds are growing actively.
- Always add 2 pints of oil concentrate per acre.

## RECOMMENDATION FOR GRASS CONTROL

- Apply POAST with ground equipment to actively growing grasses at sizes indicated in the following table.
- Always follow recommendations given in the Application Information section.



9911

RH02ARB

ANNUAL GRASSES  
(IL, IN, MI, MN, OH and WI)

ANNUAL GRASSES	MAXIMUM HEIGHT OF GRASSES AT TIME OF APPLICATION	RATE OF POAST PER ACRE	ADDITIVE/ OIL CONCENTRATE
Barnyardgrass	8"	1 pint	2 pints
Crabgrass, Large	4"	1 pint	
Foxtail, Giant	8"	1 pint	
Foxtail, Green	8"	1 pint	
Foxtail, Yellow	8"	1 pint	
Panicum, Fall	8"	1 pint	
Sandbur, Field	3"	1-1/2 pints	

## SPOT TREATMENT APPLICATION WITH POAST

For control of grasses when using knapsack sprayers or high volume equipment utilizing handguns or other suitable nozzle arrangement, prepare a solution of **POAST** herbicide plus oil concentrate in water according to the **Spot treatment application table** on the following page. Apply to actively growing grasses before tillering and/or seedhead formation. Apply to the foliage of grasses on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

### SPOT TREATMENT APPLICATION TABLE Annual Grass Control - Spot Application

See annual grasses listed in Broadcast Application Table above.	CONCENTRATION OF SPRAY SOLUTION		
	Poast®		Oil Concentrate
	Grass up to 6" height	Grass height up to 12" height	1%
	1%	1-1/2%	

\* Repeat application as needed.

### Perennial Grass Suppression - Spot Application

GRASSES	MAXIMUM SIZE RANGE	CONCENTRATION OF SPRAY SOLUTION	
		Poast®	Oil Concentrate
Johnsongrass Rhizome	15-20" height	1-1/2%	1%
Quackgrass	6-8" height	1-1/2%	1%

\* Repeat application as needed.

### SOLUTION TABLE

Desired Spray Solution Volume	Amount Of Poast Or Oil Concentrate To Be Added For Solution	
	1%	1-1/2%
1 Gallon	1-1/4 fl. oz.	2 fl. oz.
3 Gallons	3-3/4 fl. oz.	6 fl. oz.
5 Gallons	6-1/4 fl. oz.	10 fl. oz.

1 Tablespoon = 1/2 fl. oz.

11 3 11

## PROCEDURE FOR CLEANING SPRAY EQUIPMENT

Clean sprayer thoroughly prior to application of POAST®, particularly if a herbicide was used which has the potential to injure crops.

Consult the label of previously used herbicides for cleaning instructions. If no instructions are available, the steps listed below are suggested for cleaning of spray equipment prior to or following applications of POAST.

Step #1 Hose down thoroughly the inside as well as the outside of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of this rinse water.

Step #2 Refill tank with water while adding 1 gallon household ammonia or 1 pint household dish washing detergent per 100 gallons of water. Or add a commercial sprayer cleaner according to the manufacturer's directions. Operate the pump to circulate the detergent solution through the sprayer system for 5 to 10 minutes and discharge a small amount of solution through the boom and nozzles. Let the solution stand for 24 hours.

Step #3 Flush the detergent solution out of the spray tank through the boom.

Step #4 Remove the nozzles and screens and flush the system with two tankfuls of water.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of

excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Triple rinse container (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## CONDITIONS OF SALE AND WARRANTY

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. **BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Poast is a registered trademark of BASF AG.

:ksw