

**BASF**

RT date 11-17-93

# Conclude™ G

## herbicide

**Postemergence Herbicide****Active ingredient:**

2-[1-(ethoxyimino)butyl]-5-2-(ethylthio)propyl-3-hydroxy-2-cyclohexen-1-one\*.....18%

**Inert ingredients:**.....82%**Total**.....100.0%

\*Equivalent to 1.5 pounds per gallon

EPA Reg. No. 7969-58

EPA Est. No. 34313-TX-01

**KEEP OUT OF REACH OF CHILDREN.****WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to read it to you in detail.)

**Statement of practical treatment**

**If in eyes:** Immediately wash eyes with running water for 15 minutes. If irritation develops, consult a physician. **If on skin:** Wash affected areas with plenty of soap and water. Remove and launder contaminated clothing before re-use. If irritation develops, consult a physician. **If swallowed:** DO NOT INDUCE VOMITING. Promptly drink a large quantity of milk, egg whites, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol. Never give fluids or induce vomiting if the victim is unconscious or having convulsions. **If inhaled:** Move to fresh air. Aid in breathing, if necessary, and get immediate medical attention.

**Precautionary Statements****Hazards to Humans (And Domestic Animals):**

Causes substantial but temporary eye injury. Harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes.

**Personal Protective Equipment (PPE)****Applicators and other handlers must wear:**

- Coveralls over short-sleeved shirt and pants
- Chemical-resistant gloves such as barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or vitron ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, and loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls Statement:** When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations****Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

**Endangered Species Concerns**

**Notice:** The use of any pesticide in a manner that may kill or otherwise harm an endangered or threatened species or adversely modify their habitat is a violation of federal law.

The use of this product is controlled to prevent death or harm to Solanogloss which occurs in Solano County, California. Before using this product in this county, you must obtain the EPA Endangered Species Bulletin (EPA/ES-85-13) available from either your County Agricultural Extension Agent, the Endangered Species Specialist in the California Department of Fish and Game, or the Regional Office of the U.S. Fish and Wildlife Service (Portland, Oregon) or the U.S. Environmental Protection Agency. This bulletin must be reviewed prior to pesticide use. The use of this product is prohibited in this county unless specified otherwise in the bulletin.

**In Case of Emergency**

In case of large-scale spillage regarding this product, call:

CHEMTREC 800-424-9300

BASF Corporation 800-832-HELP

In case of medical emergency regarding this product, call:

1. Your local doctor for immediate treatment
2. Your local poison control center (hospital)
3. BASF Corporation 800-832-HELP

See inside booklet for complete **Directions For Use** and **Conditions of Sale and Warranty**.

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the **Directions For Use** section for information about this standard.

Net contents 7.5 gallons

BASF Corporation

P.O. box 13528, Research Triangle Park, NC, 27709

**BEST AVAILABLE COPY****Specimen Label****ACCEPTED**

DEC - 9 1993

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 7969-58

## Directions For Use — Tank Mix of Conclude B and Conclude G

(Hereafter referred to as **Conclude**)

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### General Information

**Conclude** is intended for the early postemergence control of a wide spectrum of broadleaf weeds and annual grasses in soybeans (See **Table 1**).

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry to

treated areas during the REI of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, or vitron  $\geq$  14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

### General Use Area

Use only in the following areas: Arkansas, Kentucky, Louisiana, Michigan, Mississippi, Missouri (bootheel counties of Butler, Dunklin, Mississippi, New Madrid, Pemiscott, Ripley, Scott, and Stoddard), and Tennessee.

### Storage and Disposal

Keep from freezing. Store above 32°F.

Do not contaminate water, food or feed by storage or disposal.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law.

**Table 1: Maximum Weed Heights Controlled by Tank Mixing Conclude B (1.5 pints per acre) and Conclude G (1.5 pints per acre) with Crop Oil Concentrate (1 pint per acre)**

Broadleaves:	Max. Weed Height	Grasses:	Max. Weed Height	Perennials: (top growth suppression)	Max. Weed Height
Balloonvine	2"	Barryardgrass	4"	Canada Thistle	6"
Beggarticks	5"	Broadleaf Signalgrass	4"	Johnsongrass (Rhizome)	4"
Bristly Starbur	3"	Crabgrass, Large	4"	Yellow Nutsedge	4"
Carpetweed	2"	Crabgrass, Smooth	4"		
Cocklebur	6"	Foxtail, Giant	4"		
Common Lambsquarters	2"	Foxtail, Green	4"		
Common Purslane	1"	Foxtail, Yellow	4"		
Common Ragweed	3"	Goosegrass	4"		
Crotalaria	6"	Johnsongrass (seedling)	4"		
Croton, Tropic	<2"	Junglegrass	4"		
Woolly	<2"	Panicum, Browntop	4"		
Dayflower	4"	Panicum, Fall	4"		
Eclipta	2"	Panicum, Texas	4"		
Giant Ragweed	6"	Red Spangletop	4"		
Jimsonweed	6"	Witchgrass	4"		
Ladythumb	6"	Woolly Cupgrass	4"		
Marshelder	2"				
Morningglories	2"				
Nightshade, Black	2"				
Pigweed, Redroot	2"				
Smooth	3"				
Prickly Sida/Tea Weed	2"				
Redweed	3"				
Sesbania	6"				
Shepherdspurse	4"				
Smartweed, Pennsylvania	4"				
Spurred Anoda	2"				
Texasweed	5"				
Tropic Croton	2"				
Velvetleaf	5"				
Vernice Mallow	2"				
Wild Buckwheat	3"				
Wild Mustard	4"				
Wild Sunflower	5"				
Wild Poinsetta	4"				

For new germination or perennial regrowth, follow up no sooner than 15 days later with **Basagran**, **Poast Plus**, **Storm**, or **Blazer**. Refer to the respective labels for **Directions For Use**.

**BEST AVAILABLE COPY**

If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Triple rinse container (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### **Mode of Action:**

**Conclude** is effective through postemergence contact and systemic activity. Weeds must be thoroughly covered with spray. Large crop-and-weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage.

#### **Crop Tolerance:**

All soybean varieties are tolerant to **Conclude** at all stages of growth. Leaf speckling may occur but plants generally outgrow this condition within 10 days.

#### **Cultivation:**

Do not cultivate before application or within five days after application of **Conclude**. Cultivation may put weeds under stress and reduce control obtained. A timely cultivation 2-3 weeks after applying **Conclude** may assist weed control in soybeans grown in rows more than 10 inches apart.

#### **Application Rate and Timing**

Apply **Conclude B** and **Conclude G** at the rate of 1.5 pints each per acre at early postemergence when weeds are small and actively growing (generally when soybeans are in the 2nd to 3rd trifoliate leaf stage of growth). Add oil concentrate at a rate of 1 pint per acre. Under excessively dry, wet, or cold conditions, which may reduce herbicidal activity, add oil concentrate at the rate of 2 pints per acre.

#### **Mixing**

Fill tank of a thoroughly clean sprayer half to two-thirds full with clean water. Start agitation and add the recommended amounts of product in the following order: **Conclude B**, oil concentrate, and **Conclude G**. Then add the remaining quantity of water. Maintain constant agitation during application. After mixing, spray within 48 hours.

#### **Ground Application**

Use a minimum of 10-20 gallons of water per broadcast acre at a minimum of 40 psi pressure (measured at the boom, not at the pump or in the line) to ensure adequate spray coverage. Use standard high pressure pesticide hollow cone or flat fan nozzles spaced 18-20 inches apart. Do not use flood, whirl chamber, or controlled droplet applicator nozzles.

#### **Aerial Application**

Use a minimum of 5 gallons of water per acre and a maximum of 40 psi pressure. To obtain uniform coverage and to avoid drift hazards, the following application equipment and practices should be used:

**Nozzle Type:** Use only diaphragm-type nozzles producing cone or fan spray patterns.

**Nozzle Height:** 6-10 feet above crop.

**Nozzle Orientation:** Nozzles must be oriented so as to discharge straight back with the air stream (opposite the direction of travel of the aircraft) and not more than 20° downwind.

Nozzles must be located no farther out than  $\frac{3}{4}$  the distance from the center of the aircraft to the end of the wing or rotor.

Do not apply **Conclude** by aircraft when wind is blowing at a velocity above 10 mph. Coarse sprays (larger droplets) are less likely to drift. Do not apply **Conclude** by air if ornamental or sensitive nontarget crops such as cotton, sugar beets, sunflowers, or okra are within 200 feet downwind.

Applicator must follow the most restrictive use cautions to avoid drift hazard and must follow labeling as well as applicable state and local regulations and ordinances.

#### **Procedure For Cleaning Spray Equipment**

Clean sprayer thoroughly before and after application of **Conclude**, particularly if a herbicide with the potential to injure crops was previously used. Consult the label of the previously used herbicide for cleaning instructions. If no instructions are available, the steps listed below are suggested for thorough cleaning of spray equipment prior to or following application of **Conclude**.

**Step 1:** Hose down thoroughly the inside as well as the outside of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of this rinse water.

**Step 2:** Refill tank with water while adding 1 gallon household ammonia or 1 pint household dishwashing detergent or 1 pound of dishwasher detergent per 100 gallons of water. Or add a commercial sprayer cleaner according to the manufacturer's directions. Operate the pump to circulate the detergent solution through the sprayer system for 5-10 minutes and discharge a small amount of solution through the boom and nozzles. Let the solution stand for 24 hours.

**Step 3:** Flush the detergent solution out of the spray tank through the boom.

**Step 4:** Remove the nozzles and screens and flush the system with two tankfuls of water.

#### **Restrictions and Limitations**

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive label applies when using tank mixes. Only one application of **Conclude** per acre per season should be made.

**BEST AVAILABLE COPY**

as stress due to lack of moisture, previous herbicide injury, mechanical injury or cold temperatures, as unsatisfactory weed control could result.

Do not apply if rainfall or irrigation is expected soon after application.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Conclude** with other pesticides (fungicides, herbicides, insecticides or miticides) additives or fertilizers not recommended on the label. BASF does not recommend the use of **Conclude** in tank mixes other than those listed on BASF labels, supplemental labels, or technical bulletins. Local agricultural authorities may be a source of information when using other BASF recommended combinations.

Do not apply **Conclude** to soybeans that show injury, leaf phytotoxicity, and/or other prior herbicide applications because this injury may be enhanced or prolonged.

Do not apply **Conclude** through any type of irrigation system.

Do not apply to soybeans within 75 days of harvest.

Do not use treated plants for feed or forage.

In case of crop failure, only soybeans or peanuts may be immediately be replanted.

Crop rotation restriction: Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in field treated with **Conclude** for 18 months following treatment.

Do not apply more than 2 pints of **Basagran** or 1 pint of **Blazer** following an application of **Conclude**.

Do not apply more than 1.5 pints of **Storm** following an application of **Conclude**.

Do not apply sequential applications of **Blazer**, **Basagran**, or **Poast Plus** within 15 days following the application of **Conclude**.

After application of **Conclude**, do not apply more than 3.5 pints of **Poast** per acre, per season.

Do not use selective application equipment such as recirculating sprayers, wiper applicators, or shielded applicators when applying **Conclude**.

**Conditions of Sale and Warranty**  
The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

*Basagran is a registered trademark of BASF AG.*

*Poast Plus, Conclude B, and Conclude G are trademarks of BASF Corporation.*

*Storm and Blazer are registered trademarks of BASF Corporation.*

© 1994 BASF Corporation

XX ----- JJ8942 A0990993

**BEST AVAILABLE COPY**

**BASF Corporation**

P.O. Box 13528

Research Triangle Park, NC 27709

**BASF**

## Supplemental Labeling

# Poast<sup>®</sup>

herbicide

ACCEPTED

DEC - 9 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7969-58

RT: 11-17-93

## Tank mix with Storm<sup>®</sup> herbicide for postemergence use in soybeans

Poast EPA Reg. No 7969-58

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

### Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

### General Information

A tank mix of Poast<sup>®</sup> herbicide plus Storm<sup>®</sup> herbicide is intended for the early postemergence control of a wide spectrum of broadleaf weeds and annual grasses in soybeans (See Table 1).

### General Use Area

Use only in the following areas: Arkansas, Kentucky, Louisiana, Michigan, Mississippi, Missouri (bootheel counties of Butler, Dunklin, Mississippi, New Madrid, Pemiscott, Ripley, Scott, and Stoddard), and Tennessee.

### Storage and Disposal

Keep from freezing. Store above 32°F.

Do not contaminate water, food or feed by storage or disposal.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law.

If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Triple rinse container

(or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### Mode of Action:

A tank mix of Poast plus Storm is effective through postemergence contact and systemic activity. Weeds must be thoroughly covered with spray. Large crop-and-weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage.

### Crop Tolerance:

All soybean varieties are tolerant to A tank mix of Poast plus Storm at all stages of growth. Leaf speckling may occur but plants generally outgrow this condition within 10 days.

### Cultivation:

Do not cultivate before application or within five days after application of the tank mix of Poast plus Storm. Cultivation may put weeds under stress and reduce control obtained. A timely cultivation 2-3 weeks after applying the tank mix of Poast plus Storm may assist weed control in soybeans grown in rows more than 10 inches apart.

### Application Rate and Timing

Apply the tank mix of Poast plus Storm at the rate of 1.5 pints each per acre at early postemergence when weeds are small and actively growing (generally when soybeans are in the 2nd to 3rd trifoliate leaf stage of growth). Add oil concentrate at a rate of 1 pint per acre. Under excessively dry, wet, or cold

conditions, which may reduce herbicidal activity, add oil concentrate at the rate of 2 pints per acre.

### Mixing

Fill tank of a thoroughly clean sprayer half to two-thirds full with clean water. Start agitation and add the recommended amounts of product in the following order: Storm, oil concentrate, and Poast. Then add the remaining quantity of water. Maintain constant agitation during application. After mixing, spray within 48 hours.

### Ground Application

Use a minimum of 10-20 gallons of water per broadcast acre at a minimum of 40 psi pressure (measured at the boom, not at the pump or in the line) to ensure adequate spray coverage. Use standard high pressure pesticide hollow cone or flat nozzles spaced 18-20 inches apart. Do not use flood, whirl chamber, or controlled droplet applicator nozzles.

### Aerial Application

Use a minimum of 5 gallons of water per acre and a maximum of 40 psi pressure. To obtain uniform coverage and to avoid drift hazards, the following application equipment practices should be used:

**Nozzle Type:** Use only diaphragm type nozzles producing cone or fan spray patterns.

**Nozzle Height:** 6-10 feet above crop.

**Nozzle Orientation:** Nozzles must be oriented so as to discharge straight back with the air stream (opposite the direction of travel of the aircraft) and not more than 20° downwind.

BEST AVAILABLE COPY

Agricultural Products

# BASF

Nozzles must be located no farther out than  $\frac{3}{4}$  the distance from the center of the aircraft to the end of the wing or rotor.

Do not apply the tank mix of **Poast plus Storm** by aircraft when wind is blowing at a velocity above 10 mph. Coarse sprays (larger droplets) are less likely to drift.

Do not apply the tank mix of **Poast plus Storm** by air if ornamental or sensitive nontarget crops such as cotton, sugar beets, sunflowers, or okra are within 200 feet downwind.

Applicator must follow the most restrictive use cautions to avoid drift hazard and must follow labeling as well as applicable state and local regulations and ordinances.

#### Procedure For Cleaning Spray Equipment

Clean sprayer thoroughly before and after application of the tank mix of **Poast plus Storm**, particularly if a herbicide with the potential to injure crops was previously used. Consult the label of the previously used herbicide for cleaning instructions. If no instructions are available, the steps listed below are suggested for thorough cleaning of spray equipment prior to or following application of the tank mix of **Poast plus Storm**.

**Step 1:** Hose down thoroughly the inside as well as the outside of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of this rinse water.

**Step 2:** Refill tank with water while adding 1 gallon household

ammonia or 1 pint household dishwashing detergent or 1 pound of dishwasher detergent per 100 gallons of water. Or add a commercial sprayer cleaner according to the manufacturer's directions. Operate the pump to circulate the detergent solution through the sprayer system for 5-10 minutes and discharge a small amount of solution through the boom and nozzles. Let the solution stand for 24 hours.

**Step 3:** Flush the detergent solution out of the spray tank through the boom.

**Step 4:** Remove the nozzles and screens and flush the system with two tankfuls of water.

**Table 1: Maximum Weed Heights Controlled by Tank Mixing Poast (1.5 pints per acre) and Storm (1.5 pints per acre) with Crop Oil Concentrate (1 pint per acre)**

Broadleaves:	Max. Weed Height	Grasses:	Max. Weed Height	Perennials: (top growth suppression)	Max. Weed Height
Balloonvine	2"	Barnyardgrass	4"	Canada Thistle	6"
Beggarticks	5"	Broadleaf Signalgrass	4"	Johnsongrass (Rhizome)	4"
Bristly Starbur	3"	Crabgrass, Large	4"	Yellow Nutsedge	4"
Carpetweed	2"	Crabgrass, Smooth	4"		
Cocklebur	6"	Foxtail, Giant	4"		
Common Lambsquarters	2"	Foxtail, Green	4"		
Common Purslane	1"	Foxtail, Yellow	4"		
Common Ragweed	3"	Goosegrass	4"		
Crotalaria	6"	Johnsongrass (seedling)	4"		
Croton, Tropic	<2"	Junglegrass	4"		
Woolly	<2"	Panicum, Browntop	4"		
Dayflower	4"	Panicum, Fall	4"		
Eclipta	2"	Panicum, Texas	4"		
Giant Ragweed	6"	Red Spangletop	4"		
Jimsonweed	6"	Witchgrass	4"		
Ladythumb	6"	Woolly Cupgrass	4"		
Marshelder	2"				
Morningglories	2"				
Nightshade, Black	2"				
Pigweed, Redroot	2"				
Smooth	3"				
Prickly Sida/Tea Weed	2"				
Redweed	3"				
Sesbania	6"				
Sheperdspurse	4"				
Smartweed, Pennsylvania	4"				
Spurred Anoda	2"				
Texasweed	5"				
Tropic Croton	2"				
Velvetleaf	5"				
Venice Mallow	2"				
Wild Buckwheat	3"				
Wild Mustard	4"				
Wild Sunflower	5"				
Wild Poinsetta	4"				

For new germination or perennial regrowth, follow up no sooner than 15 days later with **Basagran**, **Poast Plus**, **Storm**, or **Blazer**. Refer to the respective labels for **Directions For Use**.

**BEST AVAILABLE COPY**

### Restrictions and Limitations

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive label applies when using tank mixes. Only one application of the tank mix of **Poast plus Storm** per acre per season should be made. Do not apply to weeds under stress, such as stress due to lack of moisture, previous herbicide injury, mechanical injury or cold temperatures, as unsatisfactory weed control could result. Do not apply if rainfall or irrigation is expected soon after application. Physical incompatibility, reduced weed control, or crop injury may result from mixing the tank mix of **Poast plus Storm** with other pesticides (fungicides, herbicides, insecticides or miticides) additives or fertilizers not recommended on the label. BASF does not recommend the use of **Poast plus Storm** in tank mixes other than those listed on BASF labels, supplemental labels, or technical bulletins. Local agricultural authorities may be a source of information when using other BASF recommended combinations. Do not apply a tank mix of **Poast plus Storm** to soybeans that show injury, leaf phytotoxicity, and/or other prior herbicide applications because this injury may be enhanced or prolonged.

Do not apply a tank mix of **Poast plus Storm** through any type of irrigation system.

Do not apply to soybeans within 75 days of harvest.

Do not use treated plants for feed or forage.

In case of crop failure, only soybeans or peanuts may be immediately be replanted.

Crop rotation restriction: Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in field treated with a tank mix of **Poast plus Storm** for 18 months following treatment.

Do not apply more than 2 pints of **Basagran** or 1 pint of **Blazer** following an application of a tank mix of **Poast plus Storm**.

Do not apply more than 1.5 pints of **Storm** following an application of **Conclude**.

Do not apply sequential applications of **Blazer**, **Basagran**, or **Poast Plus** within 15 days following the application of a tank mix of **Poast plus Storm**.

After application of a tank mix of **Poast plus Storm**, do not apply more than 3.5 pints of **Poast** per acre, per season.

Do not use selective application equipment such as recirculating sprayers, wiper applicators, or shielded applicators when applying a tank mix of **Poast plus Storm**.

### Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

*Basagran is a registered trademark of BASF AG.*

*Poast Plus, Conclude B, and Conclude G are trademarks of BASF Corporation.*

*Storm and Blazer are registered trademarks of BASF Corporation.*

© 1994 BASF Corporation

XX ----- JJ8942 A0990993

## BEST AVAILABLE COPY

**BASF Corporation**

P.O. Box 13528

Research Triangle Park, NC 27709

# BASF