

PM 25 7969-58

Page 1 of 43

WPS Label



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

FEB 24 1994

ME-14

CHARLOTTE A. SANSON
BASF CORP.
AGRICULTURAL PRODUCTS
BOX 13528
RESEARCH TRIANGLE PARK, NC 27709

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 8/25/93 Response to PR Notice 93-7
EPA Reg. No. 7969-58
BASF POAST HERBICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- **BEFORE** selling or distributing any product bearing the final printed labeling
- AND**
- **WITHIN** one year from date of this acceptance.



Recycled/Recyclable
Printed with SoyCanada Ink on paper that
contains at least 50% recycled fiber

Page 2


Submit the final printed labeling via the U.S. Postal Service to:

**Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001**

Hand or courier deliveries of final printed labeling may be made to:

**Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202**

Sincerely,



**Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)**

Attachment

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

Charlotte A. Sanson
BASF CORP
AGRICULTURAL PRODUCTS
BOX 13528
RESEARCH TRIANGLE PARK NC 27709

Comment for: EPA Reg Nr.7969-58
BASF POAST HERBICIDE

The following specific comments pertain to your WPS
labeling submission concerning the product
cited above:

Delete the crossed-out statements on your proposed label.
They are redundant statements or phrases.

Correct the typographical errors circled on your proposed
label.

BASF

ACCEPTED
with COMMENTS
In EPA Letter Dated

FEB 24 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
7969-58

Poast®

herbicide

Active ingredient:

2-[1-(ethoxymino)butyl-5-[2-(ethylthio)propyl]-3-hydroxy-2-

cyclohexen-1-one18.0%

Inert ingredients:82.0%

Total100.0%

*Equivalent to 1.5 pounds per gallon

EPA Reg. No. 7969-58

KEEP OUT OF REACH OF CHILDREN.

WARNING

VISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to read it to you in detail.)

Statement of practical treatment

If in eyes: Immediately wash eyes with running water for 15 minutes. If irritation develops, consult a physician.

If on skin: Wash affected areas with plenty of soap and water. Remove and launder contaminated clothing before re-use. If irritation develops, consult a physician.

If swallowed: DO NOT INDUCE VOMITING. Dilute with water and get immediate medical attention. Never give fluids or induce vomiting if the victim is unconscious or having convulsions.

If inhaled: Move to fresh air. Aid in breathing, if necessary, and get immediate medical attention.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions For Use for information about this standard.

Net contents 1 gallon

BASF Corporation

P.O. Box 13528, Research Triangle Park, NC, 27709

BEST AVAILABLE COPY

Specimen Label

Table of Contents

5 7 43

Background Information	1
Warning	1
Precautionary Statements	5
Personal protective equipment	5
Engineering controls statement	5
User safety recommendations	5
Environmental hazards	5
Endangered species concerns	5
Storage and disposal	5
All crops—Directions for use	5
Control symptoms	6
Application information	6
Cultivation information	6
Ground application	6
Spray volume	6
Spray pressure	6
Nozzle selection	6
Boom height	6
Band application	6
Tall crop application	6
Air application	6
Special directions	6
Spray volume	6
Spray pressure	6
Nozzle selection	6
Boom height	6
Nozzle orientation	6
Spot or small area treatment	7
Additives	7
Addition of Dash® spray adjuvant or oil concentrate	7
Addition of Urea Ammonium Nitrate	7
Solution or Ammonium Sulfate	7
Rate per acre of additives	7
Mixing/spraying	7
Jar test for estimating	7
suitability of oil concentrate	7
Procedure for cleaning spray equipment	8
General restrictions and limitations—all crops	8
Field crops—Directions for use (Cotton, Flax, Peanuts, Soybeans, Sugar Beets, Sunflowers, Set Aside Conservation Reserve Land.)	9
Crop Specific Restrictions and Limitations	9
Regional use maps	10
Field Crop Rate Charts	
Midwest, South, and Northeast Regions	
Annual Grasses	11
Perennial Grasses	11
High and Rolling Plains of Texas and Western Oklahoma, Western Kansas and Eastern New Mexico	
Annual Grasses	12
Perennial Grasses	12
Western and Mountain States	
Annual Grasses	13
Perennial Grasses	13
Soybean tank mix or sequential application	14
Separate applications preceded or followed by Basagran® herbicide or Basagran + Blazer® herbicide tank mix	14
Tank mix combination with Basagran/Blazer® herbicides	15
Restrictions and limitations	15
(Tank mix of 2,4-D) and Poast herbicide burndown	16
Restrictions and limitations	16
Flax	
General information	17
Rate chart	17
Tank mixes for flax with Buctril and MCPA	17

Forage crops—Directions for use (Alfalfa, Birdfoot Trefoil and Sainfoin)	18
Crop Specific Restrictions and Limitations	18
Regional use maps	19
Use recommendations	20
Forage Crop Rate Charts	
Midwest, South, and Northeast Regions	
Annual Grasses	21
Perennial Grasses	21
High and Rolling Plains of Texas, Western Oklahoma, Western Kansas and Eastern New Mexico	
Annual Grasses	22
Perennial Grasses	22
Western and Mountain States	
Annual Grasses	23
Perennial Grasses	23
Tank mixes with 2,4-DB	24
Restrictions and limitations	24
Grass control in set aside conservation reserve land, fallow acreage	24
Restrictions and limitations	24
Vegetable crops—Directions for use	25
Crop Specific Restrictions and Limitations	25
Regional use maps	26
Vegetable Crop Rate Charts	
Midwest, South, and Northeast Regions	
Annual Grasses	27
Perennial Grasses	27
High and Rolling Plains of Texas, Western Oklahoma, Western Kansas and Eastern New Mexico	
Annual Grasses	28
Perennial Grasses	28
Western and Mountain States	
Annual Grasses	29
Tank mix with Lexone Sencor for potato and tomato	29
Restrictions and limitations	29
Fruit crops—Directions for use	30
Crop Specific Restrictions and Limitations	30
Fruit Crop Rate Charts	
Annual Grasses	31
Perennial Grasses	31
Strawberries—Regional	
Use maps	32
Strawberries Rate Charts	
Midwest, South, and Northeast Regions	
Annual Grasses	33
Perennial Grasses	33
High and Rolling Plains of Texas, Western Oklahoma, Western Kansas and Eastern New Mexico	
Annual Grasses	34
Perennial Grasses	34
Western and Mountain States	
Annual Grasses	35
Perennial Grasses	35

7 3 43

Nonbearing food crops—Directions for use36
Nonbearing Food Crops Rate Charts	
Annual Grasses36
Perennial Grasses36
Crops grown for seed37
Deciduous trees, nonfood crop areas, fallow land for grass control, tall fescue and growth suppression—Directions for use38
Timing and application information for tall fescue growth suppression in nonfood areas39
Spot Treatment Application39
Spot treatment application directions39
Spot Treatment Rate Charts39
Appendix	
Scientific names of grasses listed in this label40
Conditions of sale and warranty40

8 4 4 3

Precautionary Statements HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

Precautionary Statements:

Causes substantial but temporary eye injury. Do not get into eyes or on clothing. Harmful if swallowed.

Personal protective equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate, or viton ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, and loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User safety recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental hazards

Do not apply directly to water or wetlands (swamps, bogs, marshes, or potholes). Do not contaminate water when disposing of equipment washwaters.

Endangered species concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered or threatened species or adversely modify their habitat is a violation of Federal law.

The use of this product is controlled to prevent death or harm to Solano grass which occurs in Solano County, California. Before using this product in this county you must obtain the EPA Endangered Species Bulletin (EPA/ES-85-13) available from either your County Agricultural Extension Agent, the Endangered Species Specialist in the California Department of Fish and Game, or the Regional Offices of the U.S. Fish and Wildlife Service (Portland, Oregon) or the U.S. Environmental Protection Agency (San Francisco, California). **THIS BULLETIN MUST BE REVIEWED PRIOR TO PESTICIDE USE. THE USE OF THIS PRODUCT IS PROHIBITED IN THIS COUNTY UNLESS SPECIFIED OTHERWISE IN THE BULLETIN.**

In case of emergency

In case of large-scale spillage regarding this product: Avoid contact, isolate area and keep out animals and unprotected persons. Confine spill and call:

CHEMTREC 800-424-9300
BASF Corporation 800-832-HELP

In case of medical emergency regarding this product, call:

1. Your local doctor for immediate treatment
2. Your local poison control center (hospital).
3. BASF Corporation 800-832-HELP.

Storage and disposal!

Do not contaminate water, food or feed by storage or disposal.

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or waste is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous waste representative at the nearest EPA Regional Office for guidance.

Triple rinse container (or equivalent). Then offer for recycling or reconditioning, or puncture and

dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Directions for use—all crops

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural use requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40-GFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate, or viton ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

General information

Poast is a selective broad spectrum postemergence herbicide for control of annual and perennial grass weeds. **Poast** does not control sedges or broadleaf weeds.

Essentially, all grass crops such as sorghum, corn, small grains and rice, as well as ornamental grasses such as turf, are susceptible to **Poast**. Avoid all direct or indirect contact with any desired grass crop unless otherwise specified on the label for **Poast**.

Control symptoms

Poast rapidly enters the plant through the foliage and translocates throughout the plant. Control symptoms exhibited by the grass plant progress from a slowing or stopping of growth (generally within two days), to reddening of the foliage and to leaf tip burn. Subsequently, burn-back of the foliage occurs. These symptoms will generally be observed within three weeks depending on environmental conditions.

Application information

Applications can be made as broadcast, band or spot spray application at rates and growth stages listed in weed tables. Do not exceed application rates and use restrictions specified in **Restrictions and limitations**.

Apply **Poast** to actively growing grasses when they are at the proper growth stage as specified in the rate charts.

Do not apply to grasses under stress, such as stress due to lack of moisture, herbicide injury, mechanical injury or cold temperatures, since unsatisfactory control will probably result.

All **Poast** applications to control volunteer cereals (barley, corn, oats, rye, wheat) should be made prior to tillering.

Volunteer cereals which emerge from late spring through early summer (May through July) may be partially or incompletely controlled due to unfavorable conditions at time of application in the Western Region.

Poast is not recommended for spring control of volunteer cereals that emerged the previous fall.

Cultivation information

Do not cultivate within 5 days prior to application of **Poast** or within 7 days following application.

A timely cultivation after 7 days may aid in providing season-long control. For control of quackgrass, a cultivation 14 to 21 days after an initial or sequential application will aid in control.

In irrigated areas it may be necessary to irrigate prior to treatment to ensure weeds are growing actively.

Ground application

Spray volume: Under most conditions a spray volume of 10 gallons per acre is optimal. A minimum volume of 5 gallons and maximum volume of 20 gallons of spray solution per acre for broadcast application may be used. In the Western Region a minimum of 10 gallons per acre is recommended. In the High and Rolling Plains of Texas, Western Oklahoma, Western Kansas and Eastern New Mexico a maximum of 10 gallons per acre is recommended.

Spray pressure: When using standard high pressure hollow cone or flat fan nozzles adjust pressure to a minimum of 40 psi and a maximum of 60 psi measured at the nozzle.

Nozzle selection: Thorough spray coverage of grass foliage is essential. For broadcast application use standard high pressure pesticide nozzles. **Do not use flood or whirl chamber nozzles. Application of Poast herbicide with control drop applicator (CDA) nozzles is not recommended due to erratic coverage which causes inconsistent weed control.**

Boom height: Always adjust spray pressure, spray volume and height of spray boom to ensure penetration of plant canopy and thorough coverage of grasses to be controlled. When tall weeds, such as volunteer corn, are to be controlled, the boom height should be high enough to cover the entire plant. Refer to the nozzle manufacturer's directions for recommended height.

Band application: Banding of **Poast** may be used to control annual grasses. Grasses which are not covered or only partly covered by the spray mixture will not be adequately controlled. When treating taller weeds, such as volunteer corn, the spray boom must be high enough to thoroughly cover the top leaves and whorls of the plant. All recommendations are on a broadcast basis unless otherwise stated. When banding, rates of **Poast**, additives and water should be reduced in proportion to the area sprayed. Banding is not recommended for perennial grasses.

Tall crop application: When a crop, such as cotton, is 24 or more inches in height and the grasses may be below the crop canopy, drop nozzles should be used to insure good coverage of the grass species. Good coverage is essential for maximum control.

Air application

(Special directions): Do not apply **Poast** by aircraft when wind is blowing at a velocity above 10 mph (or above 5 mph in California). Coarse sprays (large droplets) are less likely to drift.

Applicator must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

Spray volume: Thorough spray coverage of grass foliage is essential. Use a minimum of 5 gallons of water per acre. Increase water volume to 10 gallons per acre if grass foliage and/or crop canopy is dense.

Spray pressure: Should not exceed 40 psi pressure.

Nozzle selection: Use only diaphragm nozzles producing cone or fan spray patterns.

Boom height: Do not exceed a maximum height of 10 feet above the crop.

Nozzle orientation: Nozzles must be oriented so as to discharge with the air stream (opposite the direction of travel of the aircraft) at approximately a 45° angle downward. Nozzles must not be located farther out than three-fourths the distance from the center of the aircraft to the end of the wing or rotor.

Spot or small area treatment

When using knapsack sprayers or high volume spray equipment utilizing hand guns or other suitable nozzle arrangements, prepare a 1% solution of **Poast** in water unless otherwise specified under specific crops. **Dash**® spray adjuvant or a recommended oil concentrate must also be used at a concentration of 1% for **Dash** and oil concentrate.

Apply to foliage of grasses on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff. Prepare the desired volume of spray solution by mixing the amount of **Poast** and the amount of **Dash** or oil concentrate in water according to

the table below. In soybeans and cotton, spot or small area treatments should not exceed 1/2 of an acre in size, and no more than 10% of any given acre should be treated.

In soybeans, do not make more than one spot or small area treatment to the same area within the same growing season. Also in soybeans, do not apply both broadcast and spot or small area treatments to the same area within the same growing season.

In cotton, do not make more than two spot or small area treatments in the same area within the same growing season.

For additional information regarding spot treatment application, see page 39.

ammonium sulfate may be better than fertilizer grade. Low quality ammonium sulfate may contain material that will not readily dissolve which could result in nozzle tip plugging. To determine quality, perform a jar test adding 1/2 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to addition to the spray tank. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet lines. **Ensure that ammonium sulfate is completely dissolved before adding other products.**

Rate per acre of additives

Table 1

Desired Spray Solution Volume	Amount to be Added to Obtain a 1% Solution	
	Poast	Dash and Oil Concentrate
1 Gallon	1 1/2 fl. oz.*	1 1/2 fl. oz.
25 Gallons	1 quart	1 quart
50 Gallons	2 quarts	2 quarts
100 Gallons	4 quarts	4 quarts

*2 Tablespoons = 1 fl. oz.

	Ground Application	Air Application
UAN Solution*	1/2-1 gallon	1/2 gallon
Ammonium Sulfate*	2 1/2 lbs.	2 1/2 lbs.
Oil Concentrate	2 pints	2 pints
Dash*	2 pints	2 pints

Additives

Addition of Dash or oil concentrate

Dash may be substituted for an oil concentrate with some exceptions. In some crops and tank mixes **Dash** is not recommended (see **Directions for use** tables in appropriate crop sections).

A nonphytotoxic oil concentrate (commonly referred to as oil concentrate) or **Dash** should always be added to the spray tank as recommended. The oil concentrate must contain either a petroleum or vegetable oil base and must meet the following criteria: 1) be nonphytotoxic, 2) contain only EPA-exempt ingredients, 3) provide good mixing quality in the jar test, and 4) be successful in local experience.

The exact composition of suitable oil concentrates will vary, however, vegetable and petroleum oil concentrates should contain emulsifiers which provide good mixing quality. For vegetable oil concentrates, it has been observed that highly refined vegetable oils are more satisfactory than unrefined vegetable oils. For additional information, see **Jar test for estimating suitability of oil concentrates**.

Addition of Urea Ammonium Nitrate Solution (UAN) or Ammonium Sulfate (AMS)

Addition of UAN Solution or AMS is recommended only for soybeans, alfalfa, flax, sunflowers, peanuts, cotton, sugar beets, and for enhanced activity on certain grass species in potato, beans, and peas. UAN solution is commonly referred to as 28%, 30%, or 32% nitrogen and is a water solution of urea and ammonium nitrate. When ammonium sulfate is used, three quarts of liquid ammonium sulfate (8-8-0 analysis) may be substituted for 2 1/2 lbs. solid ammonium sulfate.

In some areas use of a nitrogen additive has improved control of rhizome johnsongrass. Consult your local BASF representative for recommendations for your area.

Since most nitrogen solutions are mildly corrosive to galvanized, mild steel and brass spray equipment, rinse the entire spray system with water soon after use.

It is important to use high quality ammonium sulfate to avoid plugging of spray nozzles. The ammonium sulfate must be readily soluble in water and contain no insoluble materials. Local sources of high quality fine feed grade

***Dash**, UAN, and ammonium sulfate are not to be used in California. UAN and AMS are not recommended in the Pacific Northwest.

Mixing/spraying

Fill tank of a thoroughly clean sprayer one-half to two-thirds full with clean water. Start agitation and add UAN or ammonium sulfate first. Next add **Dash**® or oil concentrate; allow to mix thoroughly. (**Dash** and ammonium sulfate are not to be used in California.) Add **Poast** and remaining volume of water. **Apply Poast soon after mixing.** Maintain constant agitation during application.

Jar test for estimating suitability of oil concentrate

- 1. Water supply:** Use only water from intended source and at the source temperature.
- 2. Amount of water in jar:** For 20 gals./A spray volume use 3 1/2 cups (800 ml) of water. For 10 gals./A spray volume use 1 1/2 cups (400 ml) of water. For 5 gals./A spray volume use 3/4 cup (200 ml) of water. For other spray volumes, adjust proportionately to above.

3. **Amount of herbicide(s) and oil concentrate to add:** Add herbicide(s) and oil concentrate at the rate of 1 teaspoon (5 ml) for each pint of recommended label rate.

4. **Add components in following sequence, gently mixing between component additions:**

- 1) Water miscible or soluble products (such as Basagran® herbicide, Blazer® herbicide, ammonium sulfate, or UAN solution) when applicable.
- 2) Dash or oil concentrate.
- 3) Poast (and other emulsifiable concentrates when applicable).

5. **Cap jar, invert 10 cycles, let stand for 15 minutes, evaluate.**

6. **Evaluation:** An ideal tank mix will be uniform; thus, the suitability of the oil concentrate is questionable if any of the following are observed:

Free oil at the surface—film or globules.

Flocculation—fine particles which may be suspended in the liquid or found as a precipitated layer at the bottom of the jar.

Clabbering—thickening texture (coagulated) resembling yogurt or a curd-like texture as with cottage cheese.

Procedure for cleaning spray equipment

Clean sprayer thoroughly prior to application of Poast, particularly if a herbicide was used which has the potential to injure crops.

Consult the label of previously used herbicides for cleaning instructions. If no instructions are available, the steps listed below are suggested for cleaning of spray equipment prior to or following applications of Poast.

1. Hose down thoroughly the inside as well as the outside of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of this rinse water.
2. Refill tank with water while adding 1 gallon household ammonia or 1 pint household dish washing detergent per 100 gallons of water. Or add a commercial sprayer cleaner according to the manufacturer's directions. Operate the pump to circulate the detergent solution through the sprayer system for 5 to 10 minutes and discharge a small amount of solution through the boom and nozzles. Let the solution stand for 24 hours.
3. Flush the detergent solution out of the spray tank through the boom.
4. Remove the nozzles and screens and flush the system with two tankfuls of water.

General restrictions and limitations—all crops

Do not apply to grasses under stress, such as stress due to lack of moisture, herbicide injury, mechanical injury or cold temperatures, since unsatisfactory control will probably result.

Do not apply if rainfall is expected within one hour following application as grass control will probably be unsatisfactory

11 3 43
PHYSICAL INCOMPATIBILITY, REDUCED WEED CONTROL OR CROP INJURY MAY RESULT FROM MIXING POAST® HERBICIDE WITH PESTICIDES (Fungicides, Herbicides, Insecticides or Miticides), ADDITIVES, OR FERTILIZERS. BASF DOES NOT RECOMMEND THE USE OF POAST TANK MIXES OTHER THAN THOSE LISTED ON BASF LABELS, SUPPLEMENTAL LABELING, OR TECHNICAL BULLETINS. LOCAL AGRICULTURAL AUTHORITIES MAY BE A SOURCE OF INFORMATION WHEN USING OTHER THAN BASF RECOMMENDED COMBINATIONS. DO NOT APPLY POAST IN COMBINATION WITH OTHER PESTICIDES WHOSE LABELS CAUTION AGAINST THEIR USE IN COMBINATION WITH OIL ADJUVANTS.

Do not apply Poast as a preplant or preemergent treatment prior to corn, milo, millet or sorghum.

Do not apply through any type of irrigation system.

Do not tank mix Poast with Classic® or Scepter herbicides. Classic may cause antagonism when sprayed from 7 days prior to application, to 1 day after application of Poast. This antagonism is more likely to occur in grasses under stress conditions.

Other spray equipment: Do not use selective application equipment such as recirculating sprayers, wiper applicators, or shielded applicators.

Field crops

Cotton, Flax, Peanuts, Soybeans, Sugar Beets, Sunflowers, Set Aside Conservation Reserve Land.

Directions for use

- Apply to actively growing grasses at the sizes indicated.
- Always follow recommendations given in Application Information section (page 5).

- Always adjust spray pressure, spray volume and height of spray boom to ensure penetration of plant canopy and thorough coverage of grasses to be controlled.
- Do not apply to drought-stressed grass or grass which has gone through an extended dry period.
- In irrigated areas it may be necessary to irrigate prior to treatment with Poast to ensure weeds are growing actively.

- Labeled crops at all stages of growth are tolerant to Poast.
- Always add 1 quart Dash[®] spray adjuvant or oil concentrate per acre.
- For maximum use rate and minimum time from last application to harvest, consult Table 2).

**Table 2—Field Crops
Crop Specific Restrictions and Limitations for Poast**

Crop	Minimum Time from Application to Harvest (days)	Maximum Rate per Acre per Application (pints)	Maximum Rate per Acre per Season (pints)	Livestock Grazing or Feeding	Aircraft Application	Comments
Cotton	40	2½	7½	No**	Yes	Spot or small area treatments should not exceed 1/10 of an acre in size, and no more than 10% of any given acre should be treated. Do not make more than two spot or small area treatments in the same area within the same growing season.
Flax	75	1½	4	Yes**	Yes*	When tank mixing, follow restrictions and limitations on Buctril or MCPA label, the most restrictive label applies. See label for other information.
Peanut	40	2	2½	No**	Yes	
Set Aside Conservation Reserve Land	n/a	2½	7½	Alfalfa (see limitations on page 24)	Yes	Do not plant any other crop to be harvested for 120 days after application unless Poast is registered for use in that crop.
Soybean	90	2	5	Only seed and hay	Yes	See tank mix section for use with Basagran [®] herbicide, Blazer [®] herbicide, or 2,4-DB. Burndown application: Poast may be applied before, during or after planting. Spot or small area treatments should not exceed 1/10 of an acre in size, and no more than 10% of any given acre should be treated. Do not make more than one spot or small area treatment in the same area within the same growing season. Do not apply both broadcast and spot or small area treatments to the same area within the same growing season.
Sugar Beets	100 (if tops are fed)	2½	5	Yes**	Yes	
Sunflower	70	2½	2½	No**	Yes	Commercially released varieties of sunflower are tolerant to Poast at all stages of growth; however, leaf speckling has been occasionally observed on sunflower with no corresponding reduction in vigor or growth. Poast is not recommended for use on sunflower inbred lines grown for seed because crop safety of these lines has not adequately been established.

*Aircraft application is not a registered use in California. However, application by aircraft equipment may be allowed under State Special Local Need regulation as provided under section 24(c) of FIFRA; inquire with state authorities regarding currently allowed uses.


**Processed pulp and molasses may be fed from sugar beets. Processed meal may be fed from cotton, flax, peanut, soybean, sunflower (also soap stock.)

For additional Restrictions and Limitations see pages 8, 15, 16, 24 and 29.


Regional use maps

All rate and time of application recommendations are based on growing region. Refer to the maps below. Follow the Rate and Time of Application tables for your region only.

Midwest, South, and Northeast and all other regions not listed below (see page 11)




High and Rolling Plains of Texas, Western Oklahoma, Western Kansas and Eastern New Mexico (see page 12)



Description: An area east of the Continental Divide in New Mexico excluding the counties of Dona Ana, Luna, Sierra, Socorro and Valencia. Western Texas, Oklahoma and Kansas—West of a line running north from Del Rio to Gainesville, TX and extending along Interstate 35 to the Oklahoma-Kansas border. Then west along border to Highway 83 and then north to the Kansas-Nebraska border.

BEST AVAILABLE COPY

Western and Mountain States (see page 13)



Description: West of line following the Continental Divide, commencing at the U.S.-Canada border and terminating at the U.S.-Mexico border and also including the counties of Dona Ana, Luna, Sierra, Socorro, and Valencia in New Mexico. Includes Hawaii and Alaska.

14 7 53



Table 3
Field Crops—Annual Grasses
 (Cotton, peanuts, soybeans, sugar beets, sunflowers)
 Midwest, South and Northeast Regions

Grass	Rate and Maximum Height at Application					
	Special Early		Standard		Rescue***	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Barnyardgrass	4	¾*	8	1	12	1-1½
Crabgrass, Large	—	—	6	1	8	1½
Smooth	—	—	6	1	8	1½
Cupgrass, Woolly	—	—	8	1	—	—
Foxtails, Giant	4	¾	8	1	16	1½
Green	4	¾	8	1	16	1½
Yellow	—	—	8	1	16	1½
Goosegrass	3	¾	6	1	8	1½
Itchgrass	—	—	4	2	—	—
Johnsongrass (seedling)	—	—	8	1	15	1½
Junglerice	—	—	8	1	—	—
Millet, Wild Proso	10	½	10	½	24	1
Oats, Wild	—	—	4	1	—	—
Panicum, Browntop	—	—	8	1	—	—
Fall	4	¾	8	1	12	1½
Texas	4	¾	8	1	12	1
Red Rice	—	—	4	2	—	—
Ryegrass, Annual	—	—	8	1	—	—
Sandbur, Field	—	—	3	1¼	—	—
Shattercane/Wildcane	—	—	18	1	—	—
Signalgrass, Broadleaf	4	¾	8	1	12	1½
Sprangletop	—	—	8	1	—	—
Volunteer** Barley	—	—	4	1½	—	—
Corn	12	¾	20	1	—	—
Oats	—	—	4	1½	—	—
Rye	—	—	4	1½	—	—
Wheat	—	—	4	1½	—	—
Witchgrass	—	—	8	1	—	—

*In the following states use 1 pts: AL, AR, FL, GA, LA, MS, NC, SC, TN, TX, VA.

**See page 6 Application Information on volunteer cereals.

***Rescue treatment for controlling selected annual grasses

For best results, always apply Poast® herbicide to annual grasses at the growth stage as specified in the above table (Annual Grasses — Standard Recommendations). However, if Poast cannot be applied at the recommended time, larger annual grasses can be controlled with a later application by increasing the rate of Poast. Apply to actively growing grasses at the rates and sizes indicated above.

For crabgrass and all volunteer cereals the addition of ½–1 gallon UAN or 2½ lbs. AMS is recommended.

Table 4
Field Crops—Perennial Grasses
 (Cotton, peanuts, soybeans, sugar beets, sunflowers)
 Midwest, South and Northeast Regions

Grass	Rate and Maximum Height at Application			
	Standard Initial Application		Sequential Application	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Bermudagrass	6" Stolon	1½	4" Stolon	1
Johnsongrass (Rhizome)	25	1	12	1
Johnsongrass (No-Till)	20	1	12	1
Muhly, Wirestem	6	1¼	6	1¼
Quackgrass	8	1½	8	1

For quackgrass control, the addition of ½–1 gallon UAN or 2½ lbs. AMS is recommended.

BEST AVAILABLE COPY

15 743

Table 5
Field Crops—Annual Grasses
 (Cotton, peanuts, soybeans, sugar beets, sunflowers)
 High and Rolling Plains of Texas, Western Oklahoma,
 Western Kansas and Eastern New Mexico



Rate and Maximum Height at Application				
Grass	Standard		Rescue**	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Barnyardgrass	8	1½	16	2
Crabgrass, Smooth	4		—	—
Crabgrass, Large	4		—	—
Foxtails, Giant	8		—	—
Foxtails, Green	8		—	—
Foxtails, Yellow	8		—	—
Goosegrass	4		—	—
Johnsongrass (seedling)	8		—	—
Jungleice	8		—	—
Paricum, Browntop	8		—	—
Paricum, Fall	8		—	—
Paricum, Texas	8		—	—
Shattercane/Wildcane	18	1½	—	—
Signalgrass, Broadleaf	8		—	—
Sprangletop, Red	8		—	—
Volunteer* Barley	4		—	—
Volunteer* Corn	20		—	—
Volunteer* Oats	4		—	—
Volunteer* Rye	4		—	—
Volunteer* Wheat	4		—	—
Wild Proso Millet	10	1	—	—
Witchgrass	8	1½	—	—

*See page 6—Application Information on volunteer cereals.

**Rescue treatment for controlling selected annual grasses

For best results, always apply Poast® herbicide to annual grasses at the growth stage as specified in the above table (Annual Grasses—Standard Recommendations). However, if Poast cannot be applied at the recommended time, larger annual grasses can be controlled with a later application by increasing the rate of Poast. Apply to actively growing grasses at the rates and sizes indicated above.

Table 6
Field Crops—Perennial Grasses
 (Cotton, peanuts, soybeans, sugar beets, sunflowers)
 High and Rolling Plains of Texas, Western Oklahoma,
 Western Kansas and Eastern New Mexico

Rate and Maximum Height at Application				
Grass	Standard Initial Application		Sequential Application	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Bermudagrass	6" Stolon	2	4" Stolon	1½
Johnsongrass (Rhizome)	10	1½	8	1

BEST AVAILABLE COPY

16 7 4 3



Table 7
Field Crops—Annual Grasses
(Cotton, soybeans, sugar beets, sunflowers)
Western and Mountain States

Rate and Maximum Height at Application				
Grass	Standard		Rescue**	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Barnyardgrass	8	1½	16	2
Crabgrass, Smooth	4		—	—
Large	4		—	—
Cupgrass, Southwestern	8		—	—
Foxtails, Giant	8		—	—
Green	8		—	—
Yellow	8		—	—
Goosegrass	4		—	—
Johnsongrass (seedling)	8		—	—
Junglerice	8		—	—
Oats, Wild***	4		—	—
Panicum, Fall	4		—	—
Ryegrass, Annual	8		—	—
Shattercane/Wildcane	18		—	—
Volunteer* Barley	4	2	—	—
Corn	12	1½	—	—
Oats	4	2	—	—
Rye	4	2	—	—
Wheat	4	2	—	—
Wild Proso Millet	10	1	—	—
Witchgrass	8	1½	—	—

*See page 6 Application information on volunteer cereals.

**Rescue treatment for controlling selected annual grasses

For best results, always apply Poast to annual grasses at the growth stage as specified in the above table (Annual Grasses—Standard Recommendations). However, if Poast cannot be applied at the recommended time, larger annual grasses can be controlled with a later application by increasing the rate of Poast. Apply to actively growing grasses at the rates and sizes indicated above.

***For use in ID, OR, WA only

Table 8
Field Crops—Perennial Grasses
(Cotton, soybeans*, sugar beets, sunflowers)
Western and Mountain States

Rate and Maximum Height at Application				
Grass	Standard Initial Application		Sequential Application	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Bermudagrass	6" Stolon	2½	4" Stolon	1½
Johnsongrass (Rhizome)	10	2½	8	1½
Quackgrass	8	2½	8	1½
Ryegrass, Perennial	8	1½	8	1½

*The maximum allowable Poast dosage in soybeans is 2 pints/A per application. The maximum seasonal dosage is 5 pints/A.

BEST AVAILABLE COPY

Soybean tank mix or sequential application

General information

Poast®, **Basagran®** and **Blazer®** herbicides may be tank mixed for postemergence control of broadleaf and grass weeds. Weeds must be actively growing and at the recommended growth stages.

Separate applications should be made if: a) all weeds to be controlled are not at the correct growth stage for treatment at the same time, or b) grasses to be controlled include rhizome johnsongrass, quackgrass, bermudagrass, wirestem muhly, volunteer corn, shattercane, volunteer cereals, wild oats, red rice or itchgrass. (See rate tables on page 15).

Ground application

For the tank mixes of **Poast**, use 20 gallons of total spray solution per acre (broadcast basis) and a minimum of 40 psi pressure. Use standard high pressure, hollow cone or flat fan nozzles spaced 20 inches apart. Do not use flood or whirl chamber nozzles.

Air application

Poast + Basagran
Use a minimum of 5 gallons of total spray solution per acre.

Poast + Basagran and Poast + Blazer
Use a minimum of 10 gallons of total spray solution per acre.

Mixing

Fill spray tank half full with water, and add the recommended amount of product in the following order:

- A) **Poast + Basagran**
Add **Basagran**, UAN or ammonium sulfate, **Dash®** spray adjuvant or oil concentrate, **Poast**—while the agitator is running. Add the remaining quantity of water.
- B) **Poast + Basagran + Blazer**
Add **Basagran**, **Blazer**, oil concentrate, **Poast**—while the agitator is running. Add the remaining quantity of water.
- C) **Poast + Blazer**
Add **Blazer**, oil concentrate, **Poast**—while the agitator is running. Add the remaining quantity of water.

Soybeans—separate applications of Poast, preceded or followed by Basagran or Basagran + Blazer Tank Mix*:

Applications of **Poast** can be preceded or followed by **Basagran** and/or **Blazer** to obtain broad spectrum control of weeds listed on the respective product labels (refer to this label and the labels for **Basagran** and **Blazer**). Also refer to these product labels for timing, rate and other information for ground and aerial applications.

For best results when making separate applications a minimum period of time is recommended between applications, depending upon their order according to **Table 9**.

Table 9
Sequential Applications

Order of Application		Minimum Time Between Applications
First Product(s) Applied	Second Product(s) Applied	
Basagran	Poast	24 Hours
Basagran + Blazer	Poast	7 Days
Poast	Blazer** or Basagran or Basagran + Blazer	24 Hours
Blazer	Poast	7 Days

*Tank mixes not applicable in California

BEST AVAILABLE COPY

Table 10
Poast Tank Mix Combinations

18 7 43

Basagran (1-2 pts./A) + Poast			Blazer (1/2-1 pt./A) + Poast		Basagran + Blazer + Poast	
Grass	Max. Size (inches)	Poast Rate/A (pints)	Max. Size (inches)	Poast Rate/A (pints)	Max. Size (inches)	Poast Rate/A (pints)
Barnyardgrass	8	1 1/2	8	1 1/2	8	1 1/2
Crabgrass, Large	6	1 1/2	6	1 1/2	6	1 1/2
Smooth	6	1 1/2	6	1 1/2	6	1 1/2
Cupgrass, Woolly	8	1	8	1	8	1
Foxtail, Giant	8	1 1/2	8	1 1/2	8	1 1/2
Green	8	1 1/2	8	1 1/2	8	1 1/2
Yellow	8	1 1/2	8	1 1/2	8	1 1/2
Goosegrass	6	1 1/2	6	1 1/2	6	1 1/2
Johnsongrass (seedling)	8	1 1/2	8	1 1/2	8	1 1/2
Jungle rice	8	1 1/2	8	1/2	8	1
Millet, Wild Proso	10	3/4	10	1/2	10	3/4
Panicum, Browntop			8	1 1/2		
Fall			8	1 1/2	8	1
Texas	8	1	8	1 1/2	8	1 1/2
Signalgrass, Broadleaf	8	1 1/2	8	1 1/2	8	1 1/2
Sprangletop, Red	8	1 1/2	8	1 1/2	8	1 1/2
Volunteer Corn	12	1	—	—	—	—
Witchgrass	8	1	8	1 1/2	8	1 1/2
Additive Rate per Acre: Dash 2 pts. + UAN 1/2-1 gal. or Oil concentrate 2 pts. + UAN 1/2-1 gal.			Additive Rate per Acre: Oil concentrate 2 pts.		Additive Rate per Acre: Oil concentrate 2 pts.	

Restrictions and limitations
(partial list)

Read and follow the **Restrictions and limitations** on the labels for **Poast**, **Basagran**, and **Blazer**. The most restrictive labeling applies in tank mixes.

Do not add UAN solution or ammonium sulfate to a tank mix of **Poast + Basagran + Blazer + oil** concentrate.

Above **Poast** tank mixes are not applicable in California.

Poast® herbicide burndown
Poast + 2,4-D Low Volatile Ester
 (LVE) for use as a burndown prior
 to planting soybeans.

Selection of 2,4-D (LVE)
formulation

Use only low volatile ester formu-
 lations of 2,4-D such as 2,4-D
 isooctyl ester. Note that the recom-
 mended rate of 2,4-D is calculated
 on an acid equivalent (a. e.) basis.
 Make adjustments for the concen-
 tration of 2,4-D formulation used.
 Since the exact composition of suit-
 able products will vary, it is advised
 to conduct the Jar test for esti-
 mating suitability of oil concen-
 trates and 2,4-D (LVE) formulation
 used.

Restrictions and limitations
(partial list)

Do not plant soybeans until 3
 months after treatment or until the
 2,4-D (LVE) has disappeared from
 the soil.

Do not apply if rainfall is expected
 within 6 hours following application,
 as weed control will probably be
 unsatisfactory.

Since all crops such as sorghum,
 corn, small grains, cotton, soy-
 beans, sugar beets, trees, shrubs,
 as well as ornamental grasses such
 as turf are extremely susceptible to
 Poast plus 2,4-D (LVE) tank mix,
 avoid all direct or indirect post-
 emergence contact with
 any desired plant.

Do not spray if the wind is blowing
 toward desired sensitive plants, or
 at anytime when the wind exceeds
 6 miles per hour (refer to 2,4-D
 (LVE) label).

Observe all restrictions and limita-
 tions specified on labels for 2,4-D
 (LVE) and Poast. The most restric-
 tive labeling applies in tank mixes.

This tank mix does not control
 sedges or provide season-long
 control of hard-to-kill perennial
 weeds.

Do not apply this tank mix during or
 following planting or after soybean
 emergence; severe soybean injury
 will result.

Table 11

Poast Burndown*
Crops: Soybeans

Rate and Maximum Height at Application			
Weed Species	Max. Ht. (inches)	Poast** Rate/A (pints)	2,4-D*** Lbs. a.e. (lbs.)
Barnyardgrass	3	1/2	1/2
Crabgrass, Large Smooth			
Cupgrass, Woolly			
Foxtails, Giant Green Yellow			
Johnsongrass, seedling			
Fall, Panicum			
Signalgrass, Broadleaf			
Wild Proso Millet	4		
Witchgrass	3		
*For annual grass only—Poast may be applied before, during, or after planting in accordance with the Directions for use. Apply to actively growing grasses up to the maximum indicated in the rate table for field crops. **Always add Dash® spray adjuvant at 1 pint/A or oil concentrate. ***See 2,4-D label for specific broadleaf weed information.			

BEST AVAILABLE COPY



Reg # 7969-58

PM-25

ju
1922

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

FEB 21 1994

CHARLOTTE A. SANSON
BASF CORP.
AGRICULTURAL PRODUCTS
BOX 13528
RESEARCH TRIANGLE PARK, NC 27709

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 8/25/93 Response to PR Notice 93-7
EPA Reg. No. 7969-58
BASF POAST HERBICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola ink on paper that
contains at least 50% recycled fiber

Page 2


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

39 22

BASF

ACCEPTED
with COMMENTS
in EPA Letter Dated

ES 24 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

7969-58

Poast[®]

herbicide

Active ingredient:

2-[1-(ethoxymino)butyl-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one	18.0%
Inert ingredients	82.0%
Total	100.0%

*Equivalent to 1.5 pounds per gallon

EPA Reg. No. 7969-58

KEEP OUT OF REACH OF CHILDREN.

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find someone to read it to you in detail.)

Statement of practical treatment

If in eyes: Immediately wash eyes with running water for 15 minutes. If irritation develops, consult a physician.

If on skin: Wash affected areas with plenty of soap and water. Remove and launder contaminated clothing before re-use. If irritation develops, consult a physician.

If swallowed: DO NOT INDUCE VOMITING. Dilute with water and get immediate medical attention. Never give fluids or induce vomiting if the victim is unconscious or having convulsions.

If inhaled: Move to fresh air. Aid in breathing, if necessary, and get immediate medical attention.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions For Use for information about this standard.

Net contents 1 gallon

BASF Corporation

P.O. Box 13528, Research Triangle Park, NC, 27709

BEST AVAILABLE COPY

Specimen Label

49 22

Forage crops—Directions for use (Alfalfa, Birdfoot Trefoil and Sainfoin)	18
Crop Specific Restrictions and Limitations	18
Regional use maps	19
Use recommendations	20
Forage Crop Rate Charts	
Midwest, South, and Northeast Regions	
Annual Grasses	21
Perennial Grasses	21
High and Rolling Plains of Texas, Western Oklahoma, Western Kansas and Eastern New Mexico	
Annual Grasses	22
Perennial Grasses	22
Western and Mountain States	
Annual Grasses	23
Perennial Grasses	23
Tank mixes with 2,4-DB	24
Restrictions and limitations	24
Grass control in set aside conservation reserve land, fallow acreage	24
Restrictions and limitations	24
Vegetable crops—Directions for use	25
Crop Specific Restrictions and Limitations	25
Regional use maps	26
Vegetable Crop Rate Charts	
Midwest, South, and Northeast Regions	
Annual Grasses	27
Perennial Grasses	27
High and Rolling Plains of Texas, Western Oklahoma, Western Kansas and Eastern New Mexico	
Annual Grasses	28
Perennial Grasses	28
Western and Mountain States	
Annual Grasses	29
Tank mix with Lexone/Sencor for potato and tomato	29
Restrictions and limitations	29
Fruit crops—Directions for use	30
Crop Specific Restrictions and Limitations	30
Fruit Crop Rate Charts	
Annual Grasses	31
Perennial Grasses	31
Strawberries—Regional	
Use maps	32
Strawberries Rate Charts	
Midwest, South, and Northeast Regions	
Annual Grasses	33
Perennial Grasses	33
High and Rolling Plains of Texas, Western Oklahoma, Western Kansas and Eastern New Mexico	
Annual Grasses	34
Perennial Grasses	34
Western and Mountain States	
Annual Grasses	35
Perennial Grasses	35

BEST AVAILABLE COPY

Precautionary Statements HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

Precautionary Statements:

Causes substantial but temporary eye injury. Do not get into eyes or on clothing. Harmful if swallowed.

Personal protective equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate, or viton ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, and loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User safety

recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental hazards

Do not apply directly to water or wetlands (swamps, bogs, marshes, or potholes). Do not contaminate water when disposing of equipment washwaters.

Endangered species concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered or threatened species or adversely modify their habitat is a violation of Federal law. The use of this product is controlled to prevent death or harm to Solano grass which occurs in Solano County, California. Before using this product in this county you must obtain the EPA Endangered Species Bulletin (EPA/ES-85-13) available from either your County Agricultural Extension Agent, the Endangered Species Specialist in the California Department of Fish and Game, or the Regional Offices of the U.S. Fish and Wildlife Service (Portland, Oregon) or the U.S. Environmental Protection Agency (San Francisco, California). **THIS BULLETIN MUST BE REVIEWED PRIOR TO PESTICIDE USE. THE USE OF THIS PRODUCT IS PROHIBITED IN THIS COUNTY UNLESS SPECIFIED OTHERWISE IN THE BULLETIN.**

In case of emergency

In case of large-scale spillage regarding this product:

Avoid contact, isolate area and keep out animals and unprotected persons. Confine spill and call:
CHEMTREC 800-424-9300
BASF Corporation 800-832-HELP

In case of medical emergency regarding this product, call:

1. Your local doctor for immediate treatment.
2. Your local poison control center (hospital).
3. BASF Corporation 800-832-HELP.

Storage and disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Triple rinse container (or equivalent). Then offer for recycling or reconditioning, or puncture and

dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Directions for use—all crops

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural use requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate, or viton ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

BEST AVAILABLE COPY

Spot or small area treatment

When using knapsack sprayers or high volume spray equipment utilizing hand guns or other suitable nozzle arrangements, prepare a 1% solution of **Poast** in water unless otherwise specified under specific crops. **Dash**® spray adjuvant or a recommended oil concentrate must also be used at a concentration of 1% for **Dash** and oil concentrate.

Apply to foliage of grasses on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff. Prepare the desired volume of spray solution by mixing the amount of **Poast** and the amount of **Dash** or oil concentrate in water according to

the table below. In soybeans and cotton, spot or small area treatments should not exceed $\frac{1}{2}$ of an acre in size, and no more than 10% of any given acre should be treated.

In soybeans, do not make more than one spot or small area treatment to the same area within the same growing season. Also in soybeans, do not apply both broadcast and spot or small area treatments to the same area within the same growing season.

In cotton, do not make more than two spot or small area treatments in the same area within the same growing season.

For additional information regarding spot treatment application, see page 39.

ammonium sulfate may be better than fertilizer grade. Low quality ammonium sulfate may contain material that will not readily dissolve which could result in nozzle tip plugging. To determine quality, perform a jar test adding $\frac{1}{2}$ cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to addition to the spray tank. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet lines. **Ensure that ammonium sulfate is completely dissolved before adding other products.**

Rate per acre of additives

Table 1

Desired Spray Solution Volume	Amount to be Added to Obtain a 1% Solution	
	Poast	Dash and Oil Concentrate
1 Gallon	1 $\frac{1}{4}$ fl. oz.*	1 $\frac{1}{4}$ fl. oz.
25 Gallons	1 quart	1 quart
50 Gallons	2 quarts	2 quarts
100 Gallons	4 quarts	4 quarts

*2 Tablespoons = 1 fl. oz.

Additives

Addition of Dash or oil concentrate

Dash may be substituted for an oil concentrate with some exceptions. In some crops and tank mixes

Dash is not recommended (see **Directions for use** tables in appropriate crop sections).

A nonphytotoxic oil concentrate (commonly referred to as oil concentrate) or **Dash** should always be added to the spray tank as recommended. The oil concentrate must contain either a petroleum or vegetable oil base and must meet the following criteria: 1) be nonphytotoxic, 2) contain only EPA-exempt ingredients, 3) provide good mixing quality in the jar test, and 4) be successful in local experience.

The exact composition of suitable oil concentrates will vary, however, vegetable and petroleum oil concentrates should contain emulsifiers which provide good mixing quality. For vegetable oil concentrates, it has been observed that highly refined vegetable oils are more satisfactory than unrefined vegetable oils. For additional information, see **Jar test for estimating suitability of oil concentrates**.

Addition of Urea Ammonium Nitrate Solution (UAN) or Ammonium Sulfate (AMS)

Addition of UAN Solution or AMS is recommended only for soybeans, alfalfa, flax, sunflowers, peanuts, cotton, sugar beets, and for enhanced activity on certain grass species in potato, beans, and peas. UAN solution is commonly referred to as 28%, 30%, or 32% nitrogen and is a water solution of urea and ammonium nitrate. When ammonium sulfate is used, three quarts of liquid ammonium sulfate (8-8-0 analysis) may be substituted for 2 $\frac{1}{2}$ lbs. solid ammonium sulfate.

In some areas use of a nitrogen additive has improved control of rhizome johnsongrass. Consult your local BASF representative for recommendations for your area.

Since most nitrogen solutions are mildly corrosive to galvanized, mild steel and brass spray equipment, rinse the entire spray system with water soon after use.

It is important to use high quality ammonium sulfate to avoid plugging of spray nozzles. The ammonium sulfate must be readily soluble in water and contain no insoluble materials. Local sources of high quality fine feed grade

	Ground Application	Air Application
UAN Solution*	$\frac{1}{2}$ -1 gallon	$\frac{1}{2}$ gallon
Ammonium Sulfate*	2 $\frac{1}{2}$ lbs.	2 $\frac{1}{2}$ lbs.
Oil Concentrate	2 pints	2 pints
Dash *	2 pints	2 pints

***Dash**, UAN, and ammonium sulfate are not to be used in California. UAN and AMS are not recommended in the Pacific Northwest.

Mixing/spraying

Fill tank of a thoroughly clean sprayer one-half to two-thirds full with clean water. Start agitation and add UAN or ammonium sulfate first. Next add **Dash**® or oil concentrate; allow to mix thoroughly. (**Dash** and ammonium sulfate are not to be used in California.) Add **Poast** and remaining volume of water. **Apply Poast soon after mixing.** Maintain constant agitation during application.

Jar test for estimating suitability of oil concentrate

1. **Water supply:** Use only water from intended source and at the source temperature.
2. **Amount of water in jar:**
For 20 gals./A spray volume use 3 $\frac{1}{2}$ cups (800 ml) of water.
For 10 gals./A spray volume use 1 $\frac{1}{2}$ cups (400 ml) of water.
For 5 gals./A spray volume use $\frac{1}{2}$ cup (200 ml) of water.
For other spray volumes, adjust proportionately to above.

BEST AVAILABLE COPY

73-22

Field crops

Cotton, Flax, Peanuts, Soybeans, Sugar Beets, Sunflowers, Set Aside Conservation Reserve Land.

Directions for use

- Apply to actively growing grasses at the sizes indicated.
- Always follow recommendations given in **Application Information** section (page 5).

- Always adjust spray pressure, spray volume and height of spray boom to ensure penetration of plant canopy and thorough coverage of grasses to be controlled.
- Do not apply to drought-stressed grass or grass which has gone through an extended dry period.
- In irrigated areas it may be necessary to irrigate prior to treatment with Poast to ensure weeds are growing actively.

- Labeled crops at all stages of growth are tolerant to Poast.
- Always add 1 quart Dash[®] spray adjuvant or oil concentrate per acre.
- For maximum use rate and minimum time from last application to harvest, consult Table 2).

Table 2—Field Crops
Crop Specific Restrictions and Limitations for Poast

Crop	Minimum Time from Application to Harvest (days)	Maximum Rate per Acre per Application (pints)	Maximum Rate per Acre per Season (pints)	Livestock Grazing or Feeding	Aircraft Application	Comments
Cotton	40	2½	7½	No**	Yes	Spot or small area treatments should not exceed 1/10 of an acre in size, and no more than 10% of any given acre should be treated. Do not make more than two spot or small area treatments in the same area within the same growing season.
Flax	75	1½	4	Yes**	Yes*	When tank mixing, follow restrictions and limitations on Butril or MCPA label, the most restrictive label applies. See label for other information.
Peanut	40	2	2½	No**	Yes	
Set Aside Conservation Reserve Land	n/a	2½	7½	Alfalfa (see limitations on page 24)	Yes	Do not plant any other crop to be harvested for 120 days after application unless Poast is registered for use in that crop.
Soybean	90	2	5	Only seed and hay	Yes	See tank mix section for use with Basagran [®] herbicide, Blazer [®] herbicide, or 2,4-DB. Burndown application: Poast may be applied before, during or after planting. Spot or small area treatments should not exceed 1/10 of an acre in size, and no more than 10% of any given acre should be treated. Do not make more than one spot or small area treatment in the same area within the same growing season. Do not apply both broadcast and spot or small area treatments to the same area within the same growing season.
Sugar Beets	100 (if tops are fed)	2½	5	Yes**	Yes	
Sunflower	70	2½	2½	No**	Yes	Commercially released varieties of sunflower are tolerant to Poast at all stages of growth; however, leaf speckling has been occasionally observed on sunflower with no corresponding reduction in vigor or growth. Poast is not recommended for use on sunflower inbred lines grown for seed because crop safety of these lines has not adequately been established.

*Aircraft application is not a registered use in California. However, application by aircraft equipment may be allowed under State Special Local Need regulation as provided under section 24(c) of FIFRA; inquire with state authorities regarding currently allowed uses.

**Processed pulp and molasses may be fed from sugar beets. Processed meal may be fed from cotton, flax, peanut, soybean, sunflower (also soap stock.)

For additional Restrictions and Limitations see pages 8, 15, 16, 24 and 29.

Table 3
Field Crops—Annual Grasses
 (Cotton, peanuts, soybeans, sugar beets, sunflowers)
 Midwest, South and Northeast Regions



Grass	Rate and Maximum Height at Application					
	Special Early		Standard		Rescue***	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Barnyardgrass	4	¾*	8	1	12	1-1½
Crabgrass, Large	—	—	6	1	8	1½
Smooth	—	—	6	1	8	1½
Cupgrass, Woolly	—	—	8	1	—	—
Foxtails, Giant	4	¾	8	1	16	1½
Green	4	¾	8	1	16	1½
Yellow	—	—	8	1	16	1½
Goosegrass	3	¾	6	1	8	1½
Itchgrass	—	—	4	2	—	—
Johnsongrass (seedling)	—	—	8	1	16	1½
Junglerice	—	—	8	1	—	—
Millet, Wild Proso	10	½	10	½	24	1
Oats, Wild	—	—	4	1	—	—
Panicum, Browntop	—	—	8	1	—	—
Fall	4	¾	8	1	12	1½
Texas	4	¾	8	1	12	1
Red Rice	—	—	4	2	—	—
Ryegrass, Annual	—	—	8	1	—	—
Sandbur, Field	—	—	3	1¼	—	—
Shattercane/Wildcane	—	—	18	1	—	—
Signalgrass, Broadleaf	4	¾	8	1	12	1½
Sprangletop	—	—	8	1	—	—
Volunteer** Barley	—	—	4	1½	—	—
Corn	12	¾	20	1	—	—
Oats	—	—	4	1½	—	—
Rye	—	—	4	1½	—	—
Wheat	—	—	4	1½	—	—
Witchgrass	—	—	8	1	—	—

*In the following states use 1 pts: AL, AR, FL, GA, LA, MS, NC, SC, TN, TX, VA.
 **See page 6 Application Information on volunteer cereals.
 ***Rescue treatment for controlling selected annual grasses
 For best results, always apply Poast® herbicide to annual grasses at the growth stage as specified in the above table (Annual Grasses - Standard Recommendations). However, if Poast cannot be applied at the recommended time, larger annual grasses can be controlled with a later application by increasing the rate of Poast. Apply to actively growing grasses at the rates and sizes indicated above.
 For crabgrass and all volunteer cereals the addition of ½-1 gallon UAN or 2½ lbs. AMS is recommended.

Table 4
Field Crops—Perennial Grasses
 (Cotton, peanuts, soybeans, sugar beets, sunflowers)
 Midwest, South and Northeast Regions

Grass	Rate and Maximum Height at Application			
	Standard Initial Application		Sequential Application	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Bermudagrass	6" Stolon	1½	4" Stolon	1
Johnsongrass (Rhizome)	25	1	12	1
Johnsongrass (No-Till)	20	1	12	1
Muhly, Wirestem	6	1¼	6	1¼
Quackgrass	8	1½	8	1

For quackgrass control, the addition of ½-1 gallon UAN or 2½ lbs. AMS is recommended.

BEST AVAILABLE

Table 7
Field Crops—Annual Grasses
 (Cotton, soybeans, sugar beets, sunflowers)
 Western and Mountain States

Rate and Maximum Height at Application				
Grass	Standard		Rescue**	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Barnyardgrass	8	1½	16	2
Crabgrass, Smooth	4		—	—
Large	4		—	—
Cupgrass, Southwestern	8		—	—
Foxtails, Giant	8		—	—
Green	8		—	—
Yellow	8		—	—
Goosegrass	4		—	—
Johnsongrass (seedling)	8		—	—
Junglerice	8		—	—
Oats, Wild***	4		—	—
Panicum, Fall	4		—	—
Ryegrass, Annual	8		—	—
Shatter cane/Wildcane	18		—	—
Volunteer* Barley	4	2	—	—
Corn	12	1½	—	—
Oats	4	2	—	—
Rye	4	2	—	—
Wheat	4	2	—	—
Wild Proso Millet	10	1	—	—
Witchgrass	8	1½	—	—

*See page 6 Application information on volunteer cereals.
 **Rescue treatment for controlling selected annual grasses
 For best results, always apply Poast to annual grasses at the growth stage as specified in the above table (Annual Grasses—Standard Recommendations). However, if Poast cannot be applied at the recommended time, larger annual grasses can be controlled with a later application by increasing the rate of Poast. Apply to actively growing grasses at the rates and sizes indicated above.
 ***For use in ID, OR, WA only.

Table 8
Field Crops—Perennial Grasses
 (Cotton, soybeans*, sugar beets, sunflowers)
 Western and Mountain States

Rate and Maximum Height at Application				
Grass	Standard Initial Application		Sequential Application	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Bermudagrass	6" Stolon	2½	4" Stolon	1½
Johnsongrass (Rhizome)	10	2½	8	1½
Quackgrass	8	2½	8	1½
Ryegrass, Perennial	8	1½	8	1½

*The maximum allowable Poast dosage in soybeans is 2 pints/A per application. The maximum seasonal dosage is 5 pints/A.

BEST AVAILABLE COPY

Table 10
Poast Tank Mix Combinations

10 7 22

Basagran (1-2 pts./A) + Poast			Blazer (½-1 pt./A) + Poast		Basagran + Blazer + Poast	
Grass	Max. Size (inches)	Poast Rate/A (pints)	Max. Size (inches)	Poast Rate/A (pints)	Max. Size (inches)	Poast Rate/A (pints)
Barnyardgrass	8	1½	8	1½	8	1½
Crabgrass, Large	6	1½	6	1½	6	1½
Smooth	6	1½	6	1½	6	1½
Cupgrass, Woolly	8	1	8	1	8	1
Foxtail, Giant	8	1½	8	1½	8	1½
Green	8	1½	8	1½	8	1½
Yellow	8	1½	8	1½	8	1½
Goosegrass	6	1½	6	1½	6	1½
Johnsongrass (seedling)	8	1½	8	1½	8	1½
Junglerice	8	1½	8	½	8	1
Millet, Wild Proso	10	¾	10	½	10	¾
Panicum, Browntop			8	1½		
Fall			8	1½	8	1
Texas	8	1	8	1½	8	1½
Signalgrass, Broadleaf	8	1½	8	1½	8	1½
Sprangletop, Red	8	1½	8	1½	8	1½
Volunteer Corn	12	1	—	—	—	—
Witchgrass	8	1	8	1½	8	1½
Additive Rate per Acre: Dash 2 pts. + UAN ½-1 gal. or Oil concentrate 2 pts. + UAN ½-1 gal.			Additive Rate per Acre: Oil concentrate 2 pts.		Additive Rate per Acre: Oil concentrate 2 pts.	

Restrictions and limitations (partial list)

Read and follow the **Restrictions and limitations** on the labels for **Poast**, **Basagran**, and **Blazer**. The most restrictive labeling applies in tank mixes.

Do not add UAN solution or ammonium sulfate to a tank mix of **Poast + Basagran + Blazer + oil** concentrate.

Above **Poast** tank mixes are not applicable in California.

Flax

General Information

Flax competes poorly with weeds. It is important to control grass weeds before the flax stand is re-

duced and the crop vigor suffers. Where flax stands are poor or when flax is growing slowly, new grass may germinate following an application of Poast. Apply Poast to

actively growing grasses at the sizes indicated in the following table. For other Restrictions and limitations see Table 2.

11
7
22

Table 12

Flax—Annual Grasses

Grass	Rate and Maximum Height at Application					
	Special Early		Standard		Rescue	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Barnyardgrass	—	—	4	1	8	1½
Cupgrass, Woolly	—	—	4	1	—	—
Foxtails, Giant*	<1½	<½	4	1	8	1½
Green	<1½	<½	4	1	8	1½
Yellow	<1½	<½	4	1	8	1½
Oats, Wild	—	—	4	1	1	1½
Panicum, Fall	—	—	4	1	—	—
Shattercane/Wildcane	—	—	8	1	—	—
Volunteer**	—	—	—	—	—	—
Barley	—	—	6	1½	—	—
Corn	—	—	8	1	—	—
Oats	—	—	6	1½	—	—
Rye	—	—	6	1½	—	—
Wheat	—	—	6	1½	—	—
Wild Proso Millet	—	—	10	½	—	—
Witchgrass	—	—	4	1	—	—

*When using the Special Early rate, the foxtail species should not have started to tiller.
 **All Poast applications to control volunteer cereals should be made prior to tillering.

Tank mixes for flax

Tank mix of Poast with Buctril® and MCPA herbicides for grass and broadleaf weed control

Use a tank mix of Poast plus MCPA or Poast plus Buctril for the control of mixed populations of grasses and broadleaf weeds listed as susceptible on the respective product labels. Prepare the tank mix by adding water soluble forms of herbicides (such as MCPA amine) to half the final water volume, then oil concentrate or Dash, then Poast, then emulsifiable herbicides (such

as Buctril), and bring the mixture to the final volume. Agitation must be continuous from the time of mixing through spraying. Include Buctril or MCPA with Poast according to the rates recommended on the respective product labels, up to a maximum of 1 pint of Buctril equivalent per acre or up to a maximum of ¼ lb. MCPA acid equivalent per acre.


Do not delay spraying broadleaf weeds even though grassy weeds are not in correct stage for treatment. Buctril or MCPA

applied with Poast may cause leaf burn, retarded growth and delayed maturity of the crop. Some reduced grassy control may be experienced with the above tank mixes.


Do not add ammonium sulfate or UAN solution to a tank mix of Poast plus Buctril or Poast plus MCPA. Follow all restrictions detailed on the MCPA or Buctril labels that apply to use in flax. The most restrictive labeling must apply to a tank mix.

Regional use maps
All application recommendations
are based on growing region. Fol-
low the recommendations for
grass control for your region only.

Midwest, South and Northeast (and all regions not listed below) (see page 21)




High and Rolling Plains of Texas, Western Oklahoma, Western Kansas and Eastern New Mexico
(see page 22)



Description: An area east of the Continental Divide in New Mexico excluding the counties of Dona Ana, Luna, Sierra, Socorro and Valencia. West Texas, Oklahoma and Kansas—West of a line running north from Del Rio to Gainesville, TX and extending along Interstate 35 to the Oklahoma-Kansas border then west along border to Highway 83 and then north to the Kansas-Nebraska border.


Western and Mountain States (see page 23)



Description: West of a line following the Continental Divide, commencing at the U.S.-Canada border and terminating at the U.S.-Mexico border and also including the counties of Dona Ana, Luna, Sierra, Socorro, and Valencia in New Mexico. Includes Hawaii and Alaska.

BEST AVAILABLE COPY

Table 14
Forage Crops—Annual Grasses
(Alfalfa, Birdsfoot Trefoil and Sainfoin)
Midwest, South and Northeast Regions



Rate and Maximum Height at Application				
Grass	Special Early		Standard	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Barnyardgrass	4	¾	8	1
Crabgrass, Large	—	—	4	1
Smooth	—	—	4	1
Cupgrass, Woolly	—	—	8	1
Foxtails, Giant	4	¾	8	1
Green	4	¾	8	1
Yellow	—	—	8	1
Goosegrass	3	¾	4	1
Itchgrass	—	—	4	2
Johnsongrass (seedling)	—	—	8	1
Junglerice	—	—	8	1
Oats, Wild	—	—	4	1
Tame	—	—	8	¾
Panicum, Browntop	—	—	8	1
Fall	4	¾	8	1
Texas	4	¾	8	1
Red Rice	—	—	4	2
Ryegrass, Annual	—	—	8	1
Sandbur, Field	—	—	3	1½
Shattercane/Wildcane	—	—	18	1
Signalgrass, Broadleaf	4	¾	8	1
Volunteer** Barley	—	—	4	1½
Corn	12	¾	20	1
Oats	—	—	4	1½
Rye	—	—	4	1½
Wheat	—	—	4	1½
Wild Proso Millet	10	½	10	1
Witchgrass	—	—	8	1

*In the following states use 1 pt: AL, AR, FL, GA, LA, MS, NC, SC, TN, TX, VA.
 **See page 6—Application information on volunteer cereals.
 For crabgrass, wild oats and all volunteer cereals, the addition of ½–1 gallon UAN or 2½ lbs. AMS is recommended

Table 15
Forage Crops—Perennial Grasses
(Alfalfa, Birdsfoot Trefoil and Sainfoin)
Midwest, South and Northeast Regions

Rate and Maximum Height at Application				
Grass	Initial Application		Sequential Applications	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Bermudagrass	6" stolon	2½	4" stolon	2½
Johnsongrass (Rhizome)	25	2½	12	2½
Quackgrass	8	2½	8	2½
Ryegrass, Perennial	8	2	8	2
Wirestem Muhly	6	1½	6	1½

For quackgrass control, the addition of ½–1 gallon UAN or 2½ lbs. AMS is recommended. For additional information, see page 7

BEST AVAILABLE COPY

Table 18
Forage Crop—Annual Grasses
(Alfalfa, Birdsfoot Trefoil and Sainfoin)
Western and Mountain States

Rate and Maximum Height at Application				
Grass	Standard		Rescue***	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Barnyardgrass	8	1½	—	—
Crabgrass, Large*	4	1½	16	2
Smooth	4	1½	—	—
Cupgrass, Southwestern	8	1½	—	—
Foxtails, Giant	8	1½	—	—
Green	8	1½	—	—
Yellow	8	1½	—	—
Goosegrass	4	1½	—	—
Johnsongrass (seedling)	8	1½	—	—
Junglerice	8	1½	—	—
Oats, Wild	4	1½	—	—
Panicum, Fall	8	1½	—	—
Ryegrass, Annual	8	1½	—	—
Shattercane/Wildcane	18	1½	—	—
Volunteer** Barley	4	2	—	—
Corn	4	2	—	—
Oats	4	2	—	—
Rye	4	2	—	—
Wheat	4	2	—	—
Wild Proso Millet	10	1	—	—
Witchgrass	8	1½	—	—

*Apply before boot stage.
 **See page 6—Application information on volunteer cereals.
 ***Rescue treatment for controlling selected annual grasses. For best results, always apply Poast® herbicide to annual grasses at the growth stage specified above (Annual Grasses—Standard Recommendations). However, if Poast cannot be applied at the recommended time, larger annual grasses can be controlled with a later application by increasing the rate of Poast. Apply to actively growing grasses at the rates and sizes indicated above.

Table 19
Forage Crops—Perennial Grasses
(Alfalfa, Birdsfoot Trefoil and Sainfoin)
Western and Mountain States

Rate and Maximum Height at Application				
Grass	Initial Application		Sequential Application	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Bermudagrass	6" Stolon	2½	4" Stolon	2½
Johnsongrass (Rhizome)	10	2½	8	2½
Quackgrass	8	2½	8	2½
Ryegrass, Perennial	8	2	8	2

Vegetable crops

Artichoke
Beans (dry & succulent)
Broccoli
Brussel Sprouts
Cabbage
Cabbage (bok choy, napa)
Chinese Broccoli
Cantaloupe

Cauliflower
Celery
Collard
Cucumber
Eggplant
Garlic
Kale
Kohlrabi

Leek
Lentil
Lettuce (head & leaf)
Muskmelon
Mustard Greens
Onion (dry bulb & green bunching)
Peas (dry & succulent)
Peppers

Potato
Pumpkin
Rape Greens
Shallot
Spinach
Squash
Tomato
Watermelon

Directions for use

- Apply to actively growing grasses at the sizes indicated.
- Always follow recommendations given in **Application Information** page 6.
- Always adjust spray pressure, spray volume and height of spray boom to ensure penetration of

plant canopy and thorough coverage of grasses to be controlled.

- Do not apply to drought-stressed grass or grass which has gone through an extended dry period.
- In irrigated areas it may be necessary to irrigate prior to treatment with **Poast** to ensure weeds

are growing actively.

- Labeled crops at all stages of growth are tolerant to **Poast**.
- **Always add 1 quart oil concentrate per acre.**
- For maximum use rate and minimum time from last application to harvest consult **Table 20**.

Table 20—Vegetables
Crop Specific Restrictions and Limitations for Poast

Crop	Minimum Time from Application to Harvest (days)	Maximum Rate per Acre per Application (pints)	Maximum Rate per Acre per Season (pints)	Livestock Grazing or Feeding	Aircraft Application	Comments
Artichoke	7	2½	5	No	Yes*	California Only
Beans, dry succulent	30 15	2½ 2½	4 4	Yes Yes	Yes* Yes*	
Bulb vegetables, garlic leek onion	30	1½	4½	No	Yes	
Broccoli	30	1½	3	No	Yes*	
Cabbage	30	1½	3	No	Yes*	
Cantaloupe	14	1½	3	No	Yes*	
Cauliflower	30	1½	3	No	Yes*	
Celery	30	1½	3	No	Yes*	
Cucumber	14	1½	3	No	Yes*	
Eggplant	20	1½	4½	No	Yes	
Lentil***	50	2½	4	No	Yes*	
Lettuce, Leaf Head	15 30	1½ 1½	3 3	No No	Yes* Yes*	
Muskmelon	14	1½	3	No	Yes*	
Peas, dry succulent	30 15	2½ 2½	4 4	Yes Yes	Yes* Yes*	
Peppers	20	1½	4½	No	Yes	
Potato	30	2½	5	No**	Yes	
Pumpkin	14	1½	3	No	Yes*	
Spinach	15	1½	3	No	Yes*	
Squash	14	1½	3	No	Yes*	
Tomato	20	1½	4½	No**	Yes	
Watermelon	14	1½	3	No	Yes*	

*Aircraft application is not a registered use in California. However, application by aircraft equipment may be allowed under State Special Local Need regulation as provided under section 24(c) of FIFRA; inquire with state authorities regarding currently allowed uses.

**Potato and tomato waste may be fed to animals.

*****Poast** is not currently registered in California for use in lentils.

For additional **Restrictions and Limitations** see pages 8 and 29.

Caution:

Poast plus oil concentrate should be used with caution under the following conditions, due to potential leaf injury.

- When the temperature exceeds 90°F and the relative humidity is 60% or greater,

OR

- Anytime the temperature exceeds 100°F, regardless of the humidity.

Table 21
Vegetables Crops—Annual Grasses
 (For maximum allowable use rate, refer to Table 20)
Midwest, South and Northeast Regions



Grass	Rate and Maximum Height at Application					
	Special Early		Standard		Rescue	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Barryardgrass	4	¾****	8	1	12	1½
Crabgrass, Large	—	—	10	1*	8	1½
Smooth	—	—	6	1	8	1½
Cupgrass, Woolly	—	—	8	1	—	—
Foxtails, Giant	4	¾	8	1	16	1½
Green	4	¾	8	1	16	1½
Yellow	—	—	8	1	16	1½
Goosegrass	3	¾	6	1	8	1½
Itchgrass	—	—	4	2	—	—
Johnsongrass (seedling)	—	—	8	1	16	1½
Junglerice	—	—	8	1	—	—
Oats, Wild	—	—	4	1½**	—	—
Panicum, Browntop	—	—	8	1	—	—
Fall	4	¾	8	1	12	1½
Texas	4	¾	8	1	12	1½
Red Rice	—	—	4	2	—	—
Ryegrass, Annual	—	—	8	1	—	—
Sandbur, Field (Midwest only)	—	—	3	1¼	—	—
Shattercane/Wildcane	—	—	18	1	—	—
Signalgrass, Broadleaf	4	¾	8	1	12	1½
Sprangletop, Red	—	—	8	1	—	—
Volunteer*** Barley	—	—	4	1½*	—	—
Corn	12	¾	20	1**	—	—
Oats	—	—	4	1½*	—	—
Rye	—	—	4	1½*	—	—
Wheat	—	—	4	1½*	—	—
Wild Proso Millet	10	½	10	½	24	1
Witchgrass	—	—	8	1	—	—

*Plus UAN or Ammonium Sulfate in legumes (beans & peas) only.
 **Plus UAN or Ammonium Sulfate in potato and legumes (bean & peas) only.
 ***See page 6—Application Information on volunteer cereals.
 ****In the following states use 1 pt. (AL, AR, FL, GA, LA, MS, NC, SC, TN, TX, VA).

Table 22
Vegetable Crops—Perennial Grasses
 (For maximum allowable use rate, refer to Table 20)
Midwest, South and Northeast Regions

Grass	Rate and Maximum Height at Application			
	Initial Application		Sequential Application	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Bermudagrass	6" Stolon	1½	4" Stolon	1
Johnsongrass (Rhizome)**	25	1	12	1*
Muhly, Wirestem	6	1½	6	1½
Quackgrass***	8	1½*	8	1*
Ryegrass, Perennial	8	1	8	1

*Plus UAN or ammonium sulfate for johnsongrass (potato only), for quackgrass (potato and legumes only).
 **When using 10 to 20 gallons of spray per acre, use 1½ pints of Poast® herbicide in the initial application.
 ***A cultivation 14 to 21 days after the last application will aid in control.

Special Use—Potatoes/Maine

In case of heavy infestations of quackgrass, apply 2½ pints per acre followed by 1½ pints per acre sequential if needed.

Table 25

Vegetable Crops—Annual Grasses

(For maximum allowable use rate, refer to Table 20)

Western and Mountain States

Rate and Maximum Height at Application		
Grass	Max. Ht. (inches)	Rate/A (pints)
Barnyardgrass	8	1½
Crabgrass, Large	4	1½*
Smooth	4	1½
Cupgrass, Southwestern	8	1½
Woolly	8	
Foxtails, Giant	8	
Green	8	
Yellow	8	
Goosegrass	4	
Johnsongrass (seedling)	8	
Junglerice	8	
Oats, Wild*	4	
Panicum, Fall	8	
Texas	8	
Ryegrass, Annual	8	
Shattercane/Wildcane	18	
Signalgrass, Broadleaf	8	
Volunteer Corn	12	
Wild Proso Millet	10	1
Witchgrass	8	1½

*Idaho, Oregon, and Washington only.

Tank mix of Poast® herbicide for annual grass and broadleaf weed control in potato and tomato*

Use a tank mix of Poast plus Lexone/Sencor for the control of mixed populations of annual grasses and broadleaf weeds listed as susceptible on the two product labels.

Rates for Poast are the same as those listed for annual grasses in the Vegetable crops section of this label. Always add oil concentrate at the rate of 2 pints per acre. Rates for Lexone/Sencor DF are as follows:

*This tank mix is not applicable in California.

Crop	Pounds Product per Acre	
	Broadcast	Directed
Potato	½ to ¾	—
Tomato	½ to ½	¾ to 1½

Add components in the following sequence: 1) Sencor, 2) oil concentrate, 3) Poast.

Restrictions and limitations (partial list)

Observe all precautionary statements and limitations on the labels of both products. The most restrictive labeling applies to tank mixes.

Do not apply Poast and Lexone/Sencor as a tank mix unless all environmental restrictions on the Sencor label can be followed.

Do not add UAN solution or ammonium sulfate to a Poast plus Lexone/Sencor tank mix.

Do not treat transplanted tomatoes within 14 days of transplanting. Tomatoes must have recovered

from transplant shock and new growth evident. Do not treat seeded tomatoes until plants have reached the 5 to 6 leaf stage.

Apply only to russeted or white-skinned varieties of potato that are not early maturing.

Do not apply this tank mix in any type of irrigation system.

Do not use this tank mix if all weeds to be controlled are not at the correct growth stage for treatment at the same time.

Do not use this tank mix if grasses to be controlled include rhizome johnsongrass, quackgrass, bermudagrass, wirestem muhly,

volunteer corn or cereal, shattercane, red rice or itchgrass.

Do not apply tank mix if crop shows injury (leaf phytotoxicity and/or plant stunting) produced by any other herbicide treatment as injury may be enhanced and/or prolonged.

For potatoes, do not apply the tank mix within 60 days of harvest.

For tomatoes, do not apply the tank mix within 20 days of harvest.

Apply only if there has been at least three successive days of sunny weather prior to application, or crop injury may occur.

Table 27

**Fruit Crops (Except Strawberries)—Annual Grasses
All Regions**

Grass	Rate and Maximum Height at Application			
	Standard		Rescue	
	Max. Ht. (inches)	Rate/A* (pints)	Max. Ht. (inches)	Rate/A* (pints)
Barnyardgrass Crabgrass, Large Smooth Cupgrass, Woolly Foxtails, Giant Green Yellow Goosegrass Johnsongrass (seedling) Junglerice Lovegrass Orchardgrass, Seedling Panicum, Fall Texas Shattercane/Wildcane Signalgrass, Broadleaf Sprangletop, Red** Tall Fescue (seedling) Volunteer*** Barley Corn Oats Rye Wheat Wild Proso Millet Witchgrass	6	1½	12	2½
*Repeat application as needed. Do not apply more than 5 pints per acre per season for blueberries, grapes, and raspberries. Do not apply more than 7½ pints per acre per season for apple, crabapple, pear, and quince. Do not apply more than 10 pints per acre per season for citrus. **Not recommended in CA and AZ. ***See page 6—Application information on volunteer cereals.				

Table 28

**Fruit Crops (Except Strawberries)—Perennial Grasses
All Regions**

Rate and Maximum Height at Application		
Grass	Initial Application	
	Max. Ht. (inches)	Rate/A (pints)*
Bermudagrass	6" Stolon	2½
Johnsongrass (Rhizome)	20	
Quackgrass	8	
Ryegrass, Perennial	6	
*Repeat application as needed. Do not apply more than 5 pints per acre per season for blueberries, grapes, and raspberries. Do not apply more than 7½ pints per acre per season for apple, crabapple, pear, and quince. Do not apply more than 10 pints per acre per season for citrus.		

Spot treatment application

For control or suppression of grasses when using knapsack sprayers or high volume equipment (hand guns or other suitable nozzle arrangements), prepare a solution of **Poast** plus oil concentrate in water according to the table. The best spray application will be a fine spray which will cover but not drench the leaves and run off. By keeping the spray gallonage low, a relatively concentrated solution (1-1½) of **Poast** is used. The best performance is obtained when the spray gallonage is maintained at 10 gallons per acre, and the spray gallonage should not exceed 20 gallons per acre.

Strawberries

17 9 22

Note to strawberry growers:

Do not tankmix or sequentially apply Poast® herbicide plus oil concentrate within one week before or after application of Tenoran® herbicide as strawberry injury may occur.

Application of Poast plus oil concentrate applied up to 6 weeks after a Sinbar® herbicide application can occasionally cause strawberry leaf injury. It is believed to be variety related. Growers should determine injury potential by treating a small area first then waiting a week before treating the rest of the strawberry field with Poast plus oil concentrate.

Table 29

Strawberries—Annual Grasses Midwest, South and Northeast Regions

Rate and Maximum Height at Application				
Grass	Standard		Rescue	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Barnyardgrass	8	1½	12	2
Crabgrass, Large	4	1½	8	2
Smooth	4	1½	8	2
Cupgrass, Woolly	8	1½	—	—
Foxtails, Giant	8	1½	16	2
Green	8	1½	16	2
Yellow	8	1½	16	2
Goosegrass	4	1½	8	2
Itchgrass	4	2½	—	—
Johnsongrass (seedling)	8	1½	16	2
Junglerice	8	1½	—	—
Millet, Wild Proso	10	¾	24	1
Oats, Wild	4	2	—	—
Panicum, Browntop	8	1½	—	—
Fall	8	1½	12	2
Texas	8	1½	12	2
Red Rice	4	2½	—	—
Ryegrass, Annual	8	1½	—	—
Shattercane/Wildcane	18	1½	—	—
Signalgrass, Broadleaf	8	1½	12	2
Sprangletop, Red	8	1½	—	—
Volunteer* Barley	6	2	—	—
Corn	20	1½	—	—
Oats	6	2	—	—
Rye	6	2	—	—
Wheat	6	2	—	—
Witchgrass	8	1½	—	—

*Poast is not recommended for spring control of volunteer cereals that emerged the previous fall.




Table 30

Strawberries—Perennial Grasses Midwest, South and Northeast Regions

Rate and Maximum Height at Application				
Grass	Initial Application		Sequential Application	
	Max. Ht. (inches)	Rate/A (pints)	Max. Ht. (inches)	Rate/A (pints)
Bermudagrass	6" Stolon	1½	4" Stolon	1
Johnsongrass (Rhizome)	10	1½	8	1
Muhly, Wresterm	6	1½	6	1
Quackgrass	8	2½	—	—
Ryegrass, Perennial	8	1½	8	1

NOTE: A cultivation between 14 to 21 days after application will aid in control. Depending on environmental conditions and crop cultural system, season-long control may not always be obtained. However, competition from quackgrass will be reduced.

Table 33
Strawberries—Annual Grasses
Western and Mountain States

Rate and Maximum Height at Application		
Grass	Standard	
	Max. Ht. (inches)	Rate/A (pints)
Bermudagrass	8	2
Crabgrass, Large Smooth	4 4	
Cupgrass, Southwestern	8	
Foxtails, Giant Green Yellow	8 8 8	
Goosegrass	4	
Johnsongrass (seedling)	8	
Junglerice	8	
Panicum, Fall Texas	8 8	
Shattercane/Wildcane	18	
Signalgrass, Br...	8	
Volunteer* Barley Corn Oats Rye Wheat	4 12 4 4 4	2½
Witchgrass	8	2

*Volunteer cereals which emerge from late spring through early summer (May through July) may be partially or incompletely controlled due to unfavorable conditions at time of application.



Table 34
Strawberries—Perennial Grasses
Western and Mountain States

Rate and Maximum Height at Application		
Grass	Single Application*	
	Max. Ht. (inches)	Rate/A (pints)*
Bermudagrass	6" Stolon	2½
Johnsongrass	10	2½
Quackgrass	8	2½

*A single application may not provide complete control of perennial grasses. Do not use more than 2½ pints per acre per year for strawberries.

Crops grown for seed
Poast is recommended for use on all crops on this label when they are grown for seed production. Follow the use recommendations as stated on this label for each crop. Slight modifications in application methods may be required for certain seed crops due to crop canopy or different cultural methods from

the corresponding food crop. Contact BASF or local authorities before modifying application methods to confirm they are not in conflict with labeling.

Poast is also registered on the following crops but only when they are grown for seed. The information provided below is only to be used as a guide. Refer to the respective SLN**** for specific use requirements.

21922
7

Table 37

Seed Crop	Weed	Height (inches)	Rate/A (pints)
Carrot* (ID, WA only)	Barnyardgrass	3-6	1½
		6-12	2½
Fine Fescue** (OR only)	Ryegrass, Annual	4-8	1½
	Brome, Downy	2-6	2½
	German Velvetgrass	2-4	2-2½
	Bentgrass, Colonial	2-4	1½-2½
	Bentgrass, Highland	2-4	1½-2½
Clover*** (CA only)	Watergrass (Barnyardgrass)		1½-2
	Ryegrass		1½-2

***SLN # ID880005 and WA 880022 (use in carrots for seed)**

- Read and follow the general recommendations under the **All crops and vegetable crops** sections.
- Use 5-20 gallons of water per acre at 40-60 psi.
- Do not apply more than 5 pints of Poast per acre to carrots in one season.

****SLN #OR830002 (use in fine fescue for seed)**

- Read and follow the general recommendations under all the **All crops** section.
- Treat only Creeping Red, Chewing and Hard fine fescue types
- Make applications to semi-dormant fine fescue in late fall (generally November 1–March 15) after maximum grass weed germination.
- Use higher rates of Poast for well-established weeds.
- If regrowth occurs or new plants emerge make a second applica-

tion at the same Poast rate and weed size listed above.

- Use a minimum of 10 gallons of water per acre at 40 psi and increase to 20 gallons and 60 psi if foliage is dense.
- Poast does not control annual bluegrass or rattail fescue.
- DO NOT graze treated fields and DO NOT feed treated fescue screenings or hay to livestock.
- DO NOT apply Poast to tall fescue because injury will occur.
- DO NOT apply Poast to fine fescue by air.

*****SLN # CA900053 (use in clover for seed)**

- Read and follow the general recommendations under all the **All crops and Forage crops** sections.
- Apply a minimum of 10 gallons of water per acre by ground and a minimum of 5 gallons by air.
- If additional flushes of annual grasses emerge after the first application, make additional applications at the same rate.

- DO NOT apply more than 7½ pints per acre per season.
- DO NOT allow clover crops treated with Poast to be grazed or treated field residues, seed millings or seeds to be used for feed or food.
- Specific reporting requirements must be followed to meet California Department of Food and Agriculture standards. DO NOT make any applications of this product until you have obtained and read a copy of SLN # CA900053 and complied with these requirements.

******SLN REGISTRATIONS ARE VALID UNTIL WITHDRAWN, SUSPENDED OR CANCELED BY THE STATE, EPA, THE 24C REGISTRANT OR BASF.**

SLN LABELS MUST BE IN POSSESSION OF THE USER AT THE TIME OF POAST APPLICATION.

Timing and application for tall fescue growth suppression in nonfood areas

22922

Use only in the states of: AL, GA, KY, NC, SC, TN, VA, WV.

- Apply to actively growing tall fescue before extensive tillering and/or seedhead formation.
- Follow water volume and spray pressure recommendations.
- Apply to tall fescue at the sizes indicated below.
- In irrigated areas it may be necessary to irrigate prior to treatment with Poast to ensure weeds are growing actively.

Timing

Apply Poast to actively growing tall fescue after it has had 4 to 6 inches of new growth, before the

emergence of seedheads and before conifer bud break. Application from July 1 to mid-August may be less effective, especially if day temperatures reach 90°F. Tall fescue must be one year old before the first application of Poast.

Do not apply to grasses under stress, such as stress due to lack of moisture, herbicide injury, or cold temperatures, since unsatisfactory suppression may result.

Adequate coverage of the leaf surface is necessary for absorption of this herbicide; thus, for optimum control, do not mow tall fescue turf for thirty days before or fourteen days after application of Poast.

Rate

Apply Poast at 1 to 1¼ pints per acre. For greater fescue suppression up to 2½ pints per acre of Poast can be used. Because of environmental differences at application and growth differences of tall fescue, control of tall fescue may exceed or fall short of that desired. Users of Poast are advised to begin use of Poast at a minimum recommended rate and adjust rates as local conditions and experience dictate. Additional applications may be made if extended growth suppression is desired.

Spot treatment application with Poast

For control of grasses when using knapsack sprayers or high volume equipment utilizing handguns or other suitable nozzle arrangement, prepare a solution of Poast plus oil concentrate in water according to the table below. Apply to actively growing grasses before tillering and/or seedhead formation. Apply to the foliage of grasses on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

Table 40
Spot Treatment Application Table
Annual Grass Control

Grasses	Concentration in Spray Solution**		
	Poast*		Oil Concentrate
	Grass up to 6" Height	Grass up to 12" Height	
See annual grasses listed in Broadcast Application tables under specific crop.	1%	1½%	1%

*Repeat application as needed.
**Refer to Table 42 (Solution Table) for preparation of desired solution volume.

*Repeat application as needed.

**Refer to Table 42 (Solution Table) for preparation of desired solution volume.

Table 41
Perennial Grass Suppression

Grasses	Maximum Size Range	Concentration in Spray Solution**	
		Poast*	Oil Concentrate
Bermudagrass (Wiregrass)	Up to 6" Height	1½%	1%
Johnsongrass (Rhizome)	15-20" Height	1½%	1%
Quackgrass	6-8" Height	1½%	1%
Wirestem Muhly	Up to 6" Runners	1½%	1%

*Repeat application as needed

**Refer to Table 42 (Solution Table) for preparation of desired volume.

Table 42
Solution Table

Desired Spray Solution Volume	Amount of Poast or Oil Concentrate to be Added for Solution	
	1	1½%
1 Gallon	1¼ fl. oz.	2 fl. oz.
3 Gallons	3¾ fl. oz.	6 fl. oz.
5 Gallons	6¼ fl. oz.	10 fl. oz.

1 Tablespoon = ½ fl. oz.

Timing and application for tall fescue growth suppression in nonfood areas

Use only in the states of: AL, GA, KY, NC, SC, TN, VA, WV.

- Apply to actively growing tall fescue before extensive tillering and/or seedhead formation.
- Follow water volume and spray pressure recommendations.
- Apply to tall fescue at the sizes indicated below.
- In irrigated areas it may be necessary to irrigate prior to treatment with Poast to ensure weeds are growing actively.

Timing

Apply Poast to actively growing tall fescue after it has had 4 to 6 inches of new growth, before the

emergence of seedheads and before conifer bud break. Application from July 1 to mid-August may be less effective, especially if day temperatures reach 90°F. Tall fescue must be one year old before the first application of Poast.

Do not apply to grasses under stress, such as stress due to lack of moisture, herbicide injury, or cold temperatures, since unsatisfactory suppression may result.

Adequate coverage of the leaf surface is necessary for absorption of this herbicide; thus, for optimum control, do not mow tall fescue turf for thirty days before or fourteen days after application of Poast.

Rate

Apply Poast at 1 to 1½ pints per acre. For greater fescue suppression up to 2½ pints per acre of Poast can be used. Because of environmental differences at application and growth differences of tall fescue, control of tall fescue may exceed or fall short of that desired. Users of Poast are advised to begin use of Poast at a minimum recommended rate and adjust rates as local conditions and experience dictate. Additional applications may be made if extended growth suppression is desired.

Spot treatment application with Poast

For control of grasses when using knapsack sprayers or high volume equipment utilizing handguns or other suitable nozzle arrangement, prepare a solution of Poast plus oil concentrate in water according to the table below. Apply to actively growing grasses before tillering and/or seedhead formation. Apply to the foliage of grasses on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

Table 40

**Spot Treatment Application Table
Annual Grass Control**

Grasses	Concentration in Spray Solution**		
	Poast*		Oil Concentrate
	Grass up to 6" Height	Grass up to 12" Height	
See annual grasses listed in Broadcast Application tables under specific crop.	1%	1½%	1%
*Repeat application as needed. **Refer to Table 42 (Solution Table) for preparation of desired solution volume.			

Table 41

Perennial Grass Suppression

Grasses	Maximum Size Range	Concentration in Spray Solution**	
		Poast*	Oil Concentrate
Bermudagrass (Wiregrass)	Up to 6" Height	1½%	1%
Johnsongrass (Rhizome)	15-20" Height	1½%	1%
Quackgrass	6-8" Height	1½%	1%
Western Muhly	Up to 6" Runners	1½%	1%
*Repeat application as needed. **Refer to Table 42 (Solution Table) for preparation of desired volume.			

Table 42

Solution Table

Desired Spray Solution Volume	Amount of Poast or Oil Concentrate to be Added for Solution	
	1	1½%
1 Gallon	1¼ fl. oz.	2 fl. oz.
3 Gallons	3¾ fl. oz.	6 fl. oz.
5 Gallons	6¼ fl. oz.	10 fl. oz.
1 Tablespoon = ½ fl. oz.		

Appendix

The following are scientific names for the weeds listed in this label. For specific recommendations on

control of these weeds, refer to the major and/or tank mix sections.

Grasses

Common Name	Scientific Name
Barnyardgrass	<i>Echinochloa crus-gali</i>
Bermudagrass	<i>Cynodon dactylon</i>
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>
Crabgrass, Large	<i>Digitaria sanguinalis</i>
Smooth	<i>Digitaria ischaemum</i>
Cupgrass, Southwestern	<i>Eriochloa gracilis</i>
Woolly	<i>Eriochloa villosa</i>
Foxtails, Giant	<i>Setaria faberi</i>
Green	<i>Setaria viridis</i>
Yellow	<i>Setaria glauca</i>
Goosegrass	<i>Eleusine indica</i>
Itchgrass	<i>Rottboellia exaltata</i>
Johnsongrass	<i>Sorghum halepense</i>
Junglerice	<i>Echinochloa colonum</i>
Lovegrass (see Stinkgrass)	
Orchardgrass	<i>Dactylis glomerata</i>
Pigeongrass (see Foxtails)	
Panicum, Browntop	<i>Panicum fasciculatum</i>
Fall	<i>Panicum dichotomiflorum</i>
Texas	<i>Panicum texanum</i>
Quackgrass	<i>Agropyron repens</i>
Red Rice	<i>Oryza sativa</i>
Ryegrass, Annual	<i>Lolium multiflorum</i>
Perennial	<i>Lolium perenne</i>
Sandbur, Field	<i>Cenchrus incertus</i>
Shattercane/Wildcane	<i>Sorghum bicolor</i>
Sprangletop, Red	<i>Leptochloa filiformis</i>
Stinkgrass	<i>Eragrostis ciliaris</i>
Tall Fescue	<i>Festuca arundinacea</i>
Tame Oats	<i>Avena sativa</i>
Volunteer Barley	<i>Hordeum vulgare</i>
Corn	<i>Zea mays</i>
Oats	<i>Avena sativa</i>
Rye	<i>Secale Cereale</i>
Wheat	<i>Triticum aestivum</i>
Watergrass (see Barnyardgrass)	
Wild Oats	<i>Avena fatua</i>
Wild Proso Millet	<i>Panicum miliaceum</i>
Wiregrass (see Bermudagrass)	
Wirestem Muhly	<i>Muhlenbergia frondosa</i>
Witchgrass	<i>Panicum capillare</i>

Conditions of sale and warranty

The Directions for use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably

fit for the purposes referred to in the Direction for use, subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Poast and Basagran is a registered trademark of BASF AG.

Dash and Blazer are registered trademarks of BASF Corporation.

Classic and Sinbar are registered trademarks of E. I. DuPont de Nemours and Company.

Buctril is a registered trademark of Rhone-Poulenc.

Lexone is a registered trademark of DuPont.

Sencor is a registered trademark of Bayer AG.

Tenoran is a registered trademark of Ciba-Geigy Corp.

© 1991, 1992, 1993 BASF Corporation

PO 1 042 JJ8840 A0730793

BASF Corporation
PO Box 13528
Research Triangle Park, NC 27709