

PM 25 11/11

SUGARBEET LABEL AMENDMENT

Recommendations for use in alfalfa, cotton, flax, peanuts, sugar beets, and sunflowers.

General use recommendations—all regions

- Apply to actively growing grasses at the sizes indicated below.
- Always follow recommendations given in **Application Information Section**.
- Always adjust spray pressure, spray volume and height of spray boom to ensure penetration of plant canopy and thorough coverage of grasses to be controlled.
- Do not apply to drought-stressed grass or grass which has gone through an extended dry period.
- In irrigated areas it may be necessary to irrigate prior to treatment with Poast to ensure weeds are growing actively
- **Always add 2 pints Dash or oil concentrate per acre.**

Restrictions and limitations—all regions

Alfalfa

- Do not apply Poast within 7 days of grazing, feeding, or cutting for (undried) forage, or within 20 days of cutting alfalfa for (dry) hay.
- Do not apply more than a total of 5 pints of Poast per acre in one season
- Poast may be applied to alfalfa by ground equipment only
- Do not apply Poast and 2,4-DB as a tank mix unless the 60-day feeding, grazing and harvesting restrictions on the 2,4-DB label can be observed

Cotton

- Do not apply Poast within 40 days of harvest.
- Do not apply more than a total of 7½ pints of Poast per acre in one season
- Do not graze treated cotton fields and do not feed treated forage to livestock.

Flax

- Do not apply Poast within 75 days of harvest.
- Do not apply more than a total of 4 pints per acre in one season to flax
- Under weather conditions of high humidity and/or temperature, some flowers may be damaged

*Dash is not to be used in California


Regional use map

All recommendations are based on growing region. Refer to the map below. Follow the recommendations for grass control for your region only.

Midwest, South and Northeast (see pages 20-27)

High and Rolling Plains of Texas, Oklahoma and Eastern New Mexico (see pages 28-33)

Western and Mountain States (see pages 34-39)



- or dislodged by applications of Poast and oil concentrate
- Do not graze or feed treated flax forage to livestock.
- Read and follow restrictions and limitations on bromoxynil or MCPA labels when tank mixing with Poast.
- Do not apply Poast by air equipment to flax.

Peanuts

- Do not apply Poast within 40 days of harvest.
- Do not apply more than a total of 2½ pints of Poast per acre in one season.
- Do not feed treated peanut forage or hay to livestock.

Sugar Beets

- Do not apply more than a total of 5 pints of Poast per acre in one season.
- **Do not apply POAST within 60 days of harvest to Sugarbeets.**

Sunflower

- Do not apply Poast within 70 days of harvest
- Do not apply more than a total of 2½ pints of Poast per acre in one season.
- Do not feed treated sunflower forage to livestock.
- Commercially released varieties of sunflower are tolerant to Poast at all stages of growth, however, leaf speckling has been occasionally observed on sunflower with no corresponding reduction in vigor or growth. Poast is not recommended for use on sunflower inbred lines grown for seed because crop safety of these lines has not been adequately established