

PM 25 7969-58 2/12/94  
**BASF**

Page 1 of 20  
Copy 2.2

LH Date:12/11/98

NVA 98-4-25-0100

# Poast<sup>®</sup>

## herbicide

**Active Ingredient:**

Sethoxydim: 2-[1-(ethoxyimino)butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one\* .....18.0%

**Inert Ingredients:** .....82.0%

**Total** .....100.0%

\*Equivalent to 1.5 pounds of sethoxydim per gallon


EPA Reg. No. 7969-58

EPA Est. No. 34313-TX-01

**KEEP OUT OF REACH OF CHILDREN.**

### WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)



See the attached booklet for complete **Precautionary Statements, Statement of Practical Treatment, Directions For Use, and Conditions of Sale and Warranty.**

**Net contents:**

## I. Precautionary Statements

Hazard to Humans and Domestic Animals

**WARNING.** Causes substantial but temporary eye injury. Do not get into eyes or on clothing. Harmful if swallowed.

### Statement of Practical Treatment

**If in eyes:** Immediately wash eyes with running water for 15 minutes. If irritation develops, consult a physician.

**If on skin:** Wash affected areas with soap and water. If irritation develops, consult a physician.

**If swallowed:** Do not induce vomiting. Dilute with water and get immediate medical attention. Never give fluids or induce vomiting if the victim is unconscious or having convulsions.

**If inhaled:** Move to fresh air. Aid in breathing if necessary, and get immediate medical attention.

### Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. For more options, refer to category **G** on an EPA chemical resistance category selection chart.

#### Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as barrier laminate or viton  $\geq 14$  mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, and loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## User Safety Recommendations

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

This product is toxic to aquatic organisms. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

## Endangered Species Concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of federal law.

## II. Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** are to be followed. This labeling must be in the user's possession during application.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, or viton  $\geq 14$  mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Do not store below 32° F or above 100° F. Store in a dry place away from heat or open flame. Avoid contamination of feed or foodstuffs.

**Pesticide Disposal:** Pesticide wastes are toxic. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### Container Disposal:

• **Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### Bulk/Mini-bulk Containers:

Refillable/reusable containers should be returned to the point of purchase for cleaning and refilling.

## Five Steps to Repackaging Minibulk Containers

While it is not required by the EPA currently, BASF has developed this mini-bulk checklist to aid in your inspection (Steps 1-5). BASF recommends that you complete these checklists every time a mini-bulk container is refilled and keep them on file.

### Step 1. Inspect all mini-bulk tanks before filling with bulk products.

- \_\_\_\_\_ • Is the tank empty and clean (according to the current approved mini-bulk tank cleanout procedure), or does it contain only residue of the same BASF product with which it is about to be filled?
- \_\_\_\_\_ • Does the tank have a capacity greater than 55 gallons?
- \_\_\_\_\_ • Has the tank been inspected to be sure it is free of any punctures or structural defects?

### Step 2. Inspect all tank valves, hoses, pumps, meter, and seals before filling with bulk product.

- \_\_\_\_\_ • Are all fittings free of visual signs of leaking or heavy wear?
- \_\_\_\_\_ • Are all fittings and plumbing clean (according to the currently approved mini-bulk tank cleanout procedure)?

### Step 3. Label all tanks properly before filling with bulk product.

- \_\_\_\_\_ • Is there an up-to-date legible product label including **Directions For Use** displayed on the tank?
- \_\_\_\_\_ • Did you write the net contents (in gallons) on the product label after every refill?
- \_\_\_\_\_ • Did you write your EPA Establishment Number on the product label?

### Step 4. Properly secure the tank to any vehicle prior to transportation.

### Step 5. Maintain a file of the following items at the location where repackaging occurs.

- \_\_\_\_\_ • A copy of the manufacturer's cleanout procedure.
- \_\_\_\_\_ • A signed copy of the manufacturer's repackaging label authorization for this retail location.

Please post these instructions in an area visible by all employees and be sure to follow them prior to filling any mini-bulk tank. Leave all product and bar code labels in place. Product labels must remain in place to comply with Department of Transportation regulations. Return container promptly to distributor.

## In Case of Emergency

In case of large-scale spillage regarding this product, call:

CHEMTREC 800-424-9300  
BASF Corporation 800-832-HELP

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment.
- Your local poison control center (hospital).
- BASF Corporation (800-832-HELP).

## III. General Information

**Poast® herbicide** is a selective, broad spectrum, postemergence herbicide for control of annual and perennial grass weeds. **Poast** does not control sedges or broadleaf weeds. Essentially, all grass crops, such as sorghum, corn, small grains, and rice, as well as ornamental grasses, such as turf, are susceptible to **Poast**.

### Mode of Action

**Poast** rapidly enters the target weed through its foliage and translocates throughout the plant. The effects range from slowing or stopping growth (generally within 2 days), to foliage reddening and leaf tip burn. Subsequently, foliage burnback may occur. These symptoms will generally be observed within 3 weeks depending on environmental conditions.

### Crop Tolerance

All labeled crops are tolerant to **Poast** at all stages of growth.

### Herbicide Resistance

Repeated use of **Poast** (or similar postemergence grass herbicides with the same mode of action) may lead to the selection of naturally occurring biotypes with resistance to these products. If poor performance cannot be attributed to adverse weather conditions or improper application methods, a resistant biotype may be present. Consult your local representative or agricultural advisor for assistance.

### Irrigation

In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth.

### Cultivation

Do not cultivate within 5 days before or 7 days after applying **Poast**. Cultivating 7-14 days after treatment may help provide season-long control.

### Cleaning Spray Equipment

Clean spray equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product, particularly if a product with the potential to injure crops was used.

4720

## IV. Application Instructions

Applications can be made to actively growing weeds as aerial, broadcast, band, or spot spray applications at the rates and growth stages listed in **Tables 3, 4 and 5**, unless instructed differently in section **IX**.

**Crop-Specific Information.** The most effective control will result from making postemergent applications of **Poast** early, when weeds are small. Delaying application permits weeds to exceed the maximum size stated and will prevent adequate control.

Apply **Poast** to the foliage of grasses on a spray-to-wet basis uniformly and completely because large leaf canopies shelter smaller weeds and can prevent adequate spray coverage. Do not spray to the point of runoff.

Do not apply when conditions favor drift from target area or when windspeed is greater than 10 mph.

All **Poast**<sup>®</sup> herbicide applications to control volunteer cereals (barley, corn, oats, rye, and wheat) should be made before tillering. Volunteer cereals that emerged the previous fall may not be adequately controlled with **Poast** applications for spring control.

In the West Region, (see regional descriptions in **Tables 3 and 4**) volunteer cereals that emerge from late spring through early summer (May through July) may be partially or incompletely controlled because of unfavorable conditions at application time.

### Air Application

**Water Volume:** Use a minimum of 5 gallons of water per acre. Increase water volume to at least 10 gallons of water per acre if grass foliage or crop canopy is dense.

**Spray Pressure:** Use up to 40 psi.

**Application Equipment:** Use only diaphragm-type nozzles that produce fan spray patterns.

### Special Directions for Aerial Application

To obtain uniform coverage and to avoid drift hazards, follow these guidelines:

- Do not apply **Poast** by aircraft when wind is blowing more than 10 mph. Use coarse sprays (larger droplets) as they are less likely to drift.
- Do not apply **Poast** by air if sensitive species are within 200 feet downwind.

The applicator must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

**Table 1. Spot Treatment Dilution**

Spray Solution Volume	Amount of Product to be Added			
	Poast (1%)	Poast (1.5%)	Oil Concentrates (1%)	Dash <sup>®</sup> HC/Sundance <sup>®</sup> HC (0.5%)
1 gallon	1.3 fl. oz.	1.9 fl. oz.	1.3 fl. oz.	0.6 fl. oz.
3 gallons	3.8 fl. oz.	5.8 fl. oz.	3.8 fl. oz.	1.9 fl. oz.
5 gallons	6.4 fl. oz.	9.6 fl. oz.	6.4 fl. oz.	3.2 fl. oz.
25 gallons	2 pints	3 pints	2 pints	1 pint
50 gallons	4 pints	6 pints	4 pints	2 pints
100 gallons	8 pints	12 pints	8 pints	4 pints

2 tablespoons = 1 fluid ounce

**Table 2. Spot Treatment Application Rates**

Grass (see <b>Tables 3-4</b> for the complete list of grasses controlled)	Concentration in Spray Solution <sup>1</sup>		
	Poast	Crop Oil Concentrates/ Methylated Seed Oil	Dash HC/ Sundance HC
Annual grasses up to 6" height	1%	1%	0.5%
Annual grasses up to 12" height	1.5%	1%	0.5%
Perennial grasses <sup>2</sup>	1.5%	1%	1%

<sup>1</sup> Refer to **Table 1 (Spot Treatment Dilution)** for preparing the desired solution volume.

<sup>2</sup> Repeat application as needed.

### Ground Application (Banding)

**Poast** may be applied by banding to control annual grasses. Banding is not recommended for perennial grasses.

Follow **Ground Application (Broadcast)** instructions for band applications. When applying **Poast** by banding, determine the amount of herbicide and water volume needed using the following formula:

$$\frac{\text{Bandwidth in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast rate}}{\text{per acre}} = \frac{\text{Banding herbicide}}{\text{rate per acre}}$$

$$\frac{\text{Bandwidth in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast volume per acre}}{\text{volume per acre}} = \frac{\text{Banding water}}{\text{volume per acre}}$$

### Ground Application (Broadcast)

**Water Volume:** Use 5-20 gallons of spray solution. In the West Region, (see regional descriptions in **Tables 3 and 4**), do not use less than 10 gallons of spray solution per acre. In the High and Rolling Plains Region, do not use more than 10 gallons of spray solution per acre.

**Spray Pressure:** Use 40-60 psi (measured at the boom, not at the pump or in the line). When crop and weed foliage is dense, use a maximum of 20 gallons of water and 60 psi.

**Application Equipment:** Use standard high-pressure pesticide flat fan or hollow cone nozzles spaced up to 20 inches apart. Do not use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles as erratic coverage can cause inconsistent weed control. When tall weeds such as volunteer corn are to be controlled, the boom should be high enough to cover the entire plant. Refer to the nozzle manufacturer's directions for recommended height. When a crop such as cotton is 24 inches or taller and the grasses are below the crop canopy, drop nozzles should be used to ensure good coverage of the grass species.

Do not use selective application equipment such as recirculating sprayers or wiper applicators.

### Rescue Treatment for Controlling Selected Annual Grasses

If **Poast** cannot be applied at the recommended time, larger annual grasses may be controlled with a later application by increasing the rate of **Poast**. Do not exceed the maximum rate per acre, per season, for specific crops (see **Table 5 and 7**).

**Table 3. Standard Application Rates and Timing — Annual Grasses**

All application rate and timing recommendations are based on growing region. Therefore, refer to the maps below and descriptions below to ensure application accuracy. Follow the **Application Rate and Timing** tables for your region only. Refer to **Table 7** for the maximum allowable use rates for specific crop and use sites.

Annual Grass	Midwest, South, and Northeast		High and Rolling Plains		West	
	Maximum Height	Rate Per Acre (pints)	Maximum Height	Rate Per Acre (pints)	Maximum Height	Rate Per Acre (pints)
Barnyardgrass	8"	1	8"	1.5	8"	1.5
Crabgrass, Large <sup>1</sup>	6"	1	4"	1.5	4"	1.5
, Smooth <sup>1</sup>	6"	1	4"	1.5	4"	1.5
Cupgrass, Southwestern	—	—	—	—	8"	1.5
, Woolly	8"	1	—	—	—	—
Fescue, Tall (seedling)	6"	1.5	—	—	—	—
Foxtail, Giant	8"	1	8"	1.5	8"	1.5
, Green	8"	1	8"	1.5	8"	1.5
, Yellow	8"	1	8"	1.5	8"	1.5
Goosegrass	6"	1	4"	1.5	4"	1.5
Itchgrass	4"	2	—	—	—	—
Johnsongrass (seedling)	8"	1	8"	1.5	8"	1.5
Junglerice	8"	1	8"	1.5	8"	1.5
Lovegrass	6"	1.5	—	—	—	—
Millet, Wild Proso	10"	0.5	10"	1	10"	1
Oats, Tame	6"	1.5	—	—	—	—
, Wild <sup>1</sup>	4"	1	—	—	4"	1.5
Orchardgrass (seedling)	6"	1.5	—	—	—	—
Panicum, Browntop	8"	1	8"	1.5	—	—
, Fall	8"	1	8"	1.5	4"	1.5
, Texas	8"	1	8"	1.5	—	—
Red Rice <sup>1</sup>	4"	2	—	—	—	—
Ryegrass, Annual	8"	1	—	—	8"	1.5
Sandbur, Field	3"	1.25	—	—	—	—
Shattercane/Wildcane <sup>1</sup>	18"	1	18"	1.5	18"	1.5
Signalgrass, Broadleaf	8"	1	8"	1.5	—	—
Sprangletop, Red	8"	1	8"	1.5	—	—
Stinkgrass	6"	1.5	—	—	—	—
Volunteer <sup>2</sup> Barley <sup>1</sup>	4"	1.5	4"	2	4"	2
Corn <sup>1</sup>	20"	1	20"	1.5	12"	1.5
Oats <sup>1</sup>	4"	1.5	4"	2	4"	2
Rye <sup>1</sup>	4"	1.5	4"	2	4"	2
Wheat <sup>1</sup>	4"	1.5	4"	2	4"	2
Witchgrass <sup>1</sup>	8"	1	8"	1.5	8"	1.5

<sup>1</sup> Add nitrogen to the crop oil concentrate to improve grass control on indicated species.




<sup>2</sup> Apply **Poast® herbicide** before tillering.

### Spot or Small Area Application

Do not make spot treatments in addition to broadcast or band treatments. When using knapsack sprayers or high-volume spray equipment with hand guns or other suitable nozzle arrangements, prepare a 1-1.5% solution of **Poast® herbicide** in water unless otherwise specified under specific crops. Use a concentration of 0.5% for **Dash® HC**, **Sundance® HC spray** **adjuvants**, or 1% for oil concentrate. Prepare the desired volume of spray solution by mixing the amount of **Poast** and the amount of **Dash HC**, **Sundance HC** or oil concentrate in water according to **Table 1**.

**Table 4. Standard Application Rates and Timing — Perennial Grasses<sup>1</sup>**

All application rate and timing recommendations are based on growing region. Therefore, refer to the maps below and descriptions below to ensure application accuracy. Follow the **Application Rate and Timing** tables for your region only. Refer to **Table 7** for the maximum allowable use rates for specific crop and use sites.

Perennial Grass	Midwest, South, and Northeast		High and Rolling Plains		West	
						
Standard Initial Application	Maximum Height	Rate Per Acre (pints)	Maximum Height	Rate Per Acre (pints)	Maximum Height	Rate Per Acre (pints)
Bermudagrass	6" stolon	1.5	6" stolon	2 <sup>2</sup>	6" stolon	2.5
Johnsongrass (Rhizome)	25"	1.5	10"	1.5 <sup>2</sup>	10"	2.5
Johnsongrass (No-Till)	20"	1.5	—	—	—	—
Muhly, Wirestem	6"	1.25	—	—	—	—
Quackgrass <sup>1</sup>	8"	1.5	—	—	8"	2.5
Ryegrass, Perennial	8"	1.5	—	—	8"	1.5
Sequential Application	Maximum Height	Rate Per Acre (pints)	Maximum Height	Rate Per Acre (pints)	Maximum Height	Rate Per Acre (pints)
Bermudagrass	4" stolon	1	4" stolon	1.5 <sup>2</sup>	4" stolon	1.5
Johnsongrass (Rhizome)	12"	1	8"	1 <sup>2</sup>	8"	1.5
Johnsongrass (No-Till)	12"	1	—	—	—	—
Muhly, Wirestem	6"	1.25	—	—	—	—
Quackgrass <sup>1</sup>	8"	1	—	—	8"	1.5
Ryegrass, Perennial	8"	1.5	—	—	8"	1.5

<sup>1</sup> Add nitrogen to the crop oil concentrate to improve grass control on indicated species. Cultivate 7-14 days after an initial or sequential application to aid control.

<sup>2</sup> Use 2.5 pints per acre for the following forage crops: alfalfa, clover, birdsfoot trefoil, sainfoin.


**Regional Descriptions**

**High and Rolling Plains:** An area east of the Continental Divide in New Mexico excluding the counties of Dona Ana, Luna, Sierra, Socorro and Valencia. Western Texas, Oklahoma and Kansas; west of a line running north from Del Rio to Gainesville, Texas, and extending along Interstate 35 to the Oklahoma-Kansas border, then west along border to Highway 83 and then north to the Kansas-Nebraska border.

**West:** West of a line following the Continental Divide, commencing at the U.S.-Canada border and terminating at the U.S.-Mexico border and also including the counties of Dona Ana, Luna, Sierra, Socorro, and Valencia in New Mexico. Includes Hawaii and Alaska.

**Midwest, South, and Northeast:** all other regions not listed above.

**Table 5. Special Application Rates and Timing for Midwest, South and Northeast**

Annual Grass				
	Special Early Maximum Height	Early Rate Per Acre (Pints)	Rescue Maximum Height	Rescue Rate Per Acre (Pints)
Barnyardgrass	4"	0.75 <sup>1</sup>	12"	1.5
Crabgrass, Large	—	—	8"	1.5
, Smooth	—	—	8"	1.5
Foxtail, Giant <sup>2</sup>	4"	0.75	16"	1.5
, Green <sup>2</sup>	4"	0.75	16"	1.5
, Yellow <sup>2</sup>	—	—	16"	1.5
Goosegrass	3"	0.75	8"	1.5
Johnsongrass (seedling)	—	—	16"	1.5
Millet, Wild Proso	10"	0.50	24"	1.0
Panicum, Fall	4"	0.75	12"	1.5
, Texas	4"	0.75	12"	1.0
Signalgrass, Broadleaf	4"	0.75	12"	1.5
Volunteer Corn <sup>3</sup>	12"	0.75	—	—

<sup>1</sup> In the following states use 1.0 pint: AL, AR, FL, GA, LA, MS, NC, SC, TN, TX, and VA.

<sup>2</sup> For flax, use 0.5 pint per acre when foxtails are less than 1.5" high. When using the special early rate, the foxtail species should not have started to tiller.

<sup>3</sup> Add nitrogen to the crop oil concentrate to improve grass control on indicated species. UAN and AMS are not recommended in the Pacific Northwest and are not registered in California.

## V. Additives

To achieve consistent weed control, always use one of the following additives: **Dash® HC spray adjuvant**, **Sundance® HC spray adjuvant**, methylated/modified seed oil, or crop oil concentrates. In addition, urea ammonium nitrate or ammonium sulfate is recommended for use on alfalfa, beans, cotton, flax, peanuts, peas, potatoes, soybeans, **Poast Protected™ field corn**, sugarbeets, and sunflowers to enhance activity on certain grass species. (See **Table 6. Additive Rates Per Acre** for more information.) Because most nitrogen solutions are mildly corrosive to galvanized, mild steel, and brass spray equipment, rinse the entire spray system with water soon after use. UAN and AMS are not recommended in the Pacific Northwest and are not registered in California.

**Table 6. Additive Rates Per Acre**

Additive	Ground Application	Aerial Application
AMS	2.5 pounds	2.5 pounds
Dash HC/Sundance HC	1 pint	1 pint
Crop Oil Concentrates	2 pints	2 pints
Methylated Seed Oils/MSO	1.5 pints	1.5 pints
UAN Solution	4-8 pints	4 pints

Consult a BASF representative or local agricultural authority for more information on the use of additives.

### **Dash HC, Sundance HC, Crop Oil Concentrates, or Methylated Seed Oils**

A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:

- be nonphytotoxic,
- contain only EPA-exempt ingredients,
- provide good mixing quality in the jar test, and
- be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. For more information, see **Compatibility Test for Mix Components**. For most crops, **Dash HC** or **Sundance HC** may be substituted for crop oil concentrate or methylated seed oil; however, for some crops and tank mixes, **Dash HC, Sundance HC** and **MSO** are not recommended. (See section **IX. Crop-Specific Information** for more information.)

### **Urea Ammonium Nitrate (UAN)**

Commonly referred to as 28%, 30% or 32% nitrogen solution, UAN may be used in addition to **Dash HC, Sundance HC**, or crop oil concentrate to improve weed control. Do not use UAN in California.

### **Ammonium Sulfate (AMS)**

When AMS is used, 3 quarts of liquid AMS (8-8-0 analysis) may be substituted for 2.5 pounds of dry AMS. If the AMS is added directly to the spray tank, add slowly while agitating. Adding the mix too quickly may clog outlet lines. Be sure the AMS is completely dissolved before adding any other products. Do not use AMS in California.

### **Compatibility Test for Mix Components**

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

- 1) **Water**. For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
- 2) **Water-dispersible products**: (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions). Cap the jar and invert 10 cycles.

- 3) **Water-soluble products** Cap the jar and invert 10 cycles.
- 4) **Emulsifiable concentrates** (such as **Poast**, **Dash HC**, **Sundance HC**, oil concentrate, or methylated seed oil when applicable). Cap the jar and invert 10 cycles.
- 5) **Water-soluble additives** (such as **AMS** or **UAN** when applicable). Cap the jar and invert 10 cycles.
- 7) Let the solution stand for 15 minutes.
- 8) **Evaluate** the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. Do not use any spray solution that could clog spray nozzles.

## VI. Mixing Order

- 1) **Water**: Begin by agitating a thoroughly clean sprayer tank half full of clean water.
- 2) **Water-dispersible products** (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions).
- 3) **Water-soluble products**
- 4) **Emulsifiable concentrates** (such as **Poast**, **Dash HC**, **Sundance HC**, oil concentrate, or methylated seed oil when applicable).
- 5) **Water-soluble additives** (**AMS** or **UAN** when applicable).
- 6) Remaining quantity water.

Maintain constant agitation during application. For more information, refer to section **VII. Tank Mixing Application**.

Read and follow the applicable **Restrictions and Limitations and Directions For Use** on all products involved in tank mixing. Refer to section **IX. Crop-Specific Information** for more details. The most restrictive labeling applies to tank mixes. Separate applications should be made if all target weeds are not at the correct growth stage for treatment at the same time.

## VII. General Tank Mixing Information

Tank mixing **Poast® herbicide** with some postemergence broadleaf herbicides has shown some reduction or failure to control some grasses that would otherwise be controlled and therefore may require a higher rate of **Poast**. However, do not exceed the maximum rate per application as listed in **Table 7**. If regrowth occurs or an additional flush of new grasses emerges, reapply **Poast** according to recommended rates in **Tables 3 and 4**.

### **Tank Mix Partners**

The following herbicides may be tank mixed with **Poast** according to the instructions in the respective product labels. Refer to section **IX. Crop-Specific Information** for more information.

- |                 |                    |
|-----------------|--------------------|
| 1. Atrazine     | 17. Laddok® S-12   |
| 2. Basagran®    | 18. Lexone®        |
| 3. Betamix®     | 19. MCPA           |
| 4. Blazer®      | 20. Pursuit®       |
| 5. Buctril®     | 21. Reflex®        |
| 6. Clarity      | 22. Reliance® STS  |
| 7. Classic®     | 23. Resource®      |
| 8. Cobra®       | 24. Sercor® DF     |
| 9. Dual®        | 25. Staple®        |
| 10. Dual II®    | 26. Stellar®       |
| 11. First Rate® | 27. Storm®         |
| 12. Flexstar®   | 28. Surpass®       |
| 13. Frontier®   | 29. Synchrony® STS |
| 14. Galaxy®     | 30. 2,4-D amine    |
| 15. Guardsman®  | 31. 2,4-DB         |
| 16. Harness®    | 32. 2,4-D (LVE)    |

### VIII. General Restrictions and Limitations — All Crops

- **Maximum seasonal use rate:** See **Table 7** for crop-specific maximum seasonal use rates.
- **Preharvest Interval:** See **Table 7** for crop-specific preharvest intervals.
- **Restricted Entry Interval (REI): 12 hours.**
- Avoid all direct or indirect contact with any desired grass crop unless otherwise recommended on the **Poast® herbicide** label.
- **Stress:** Do not apply to grasses or crops under stress such as stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control may result. Irrigation may be needed.
- Do not apply to crops that show **injury** (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced or prolonged.
- Do not apply as a **preplant** or **preemergent treatment** before planting corn, milo, millet, or sorghum.
- Do not use UAN or AMS in California.
- Do not use **selective application equipment** such as recirculating sprayers, wiper applicators, or **shielded applicators**.
- **Rainfast Period:** **Poast** is rainfast 1 hour after application.
- Do not apply through any type of **irrigation** equipment.
- Physical incompatibility, reduced weed control, or crop injury may result from mixing **Poast** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. BASF does not recommend using tank mixes other than those listed on BASF labeling. Local agricultural authorities may be a source of information when using other than BASF recommended tank mixes.



9 7 20

**Table 7. Crop-Specific Restrictions and Limitations for Poast® Herbicide**

Crop	Minimum Time From Application to Harvest (PHI)	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application	Tank Mix Partner
Alfalfa, birdsfoot trefoil, and sainfoin <sup>1</sup>	14 days before cutting for (dry) hay	2.5 pints	6.5 pints	Yes	Yes	31
Alfalfa, birdsfoot trefoil, and sainfoin (Undried) <sup>1</sup>	7 days before grazing, feeding, or cutting for (undried) forage	2.5 pints	6.5 pints	Yes	Yes	31
Apricots	25 days	2.5 pints	5.0 pints	n/a	Yes	
Artichokes <sup>2</sup> (CA only)	7 days	2.5 pints	5.0 pints	No	Yes	
Asparagus	1 day	2.5 pints	5.0 pints	No	Yes	
Avocadoes (nonbearing)	1 year	2.5 pints	7.5 pints	n/a	Yes	
Beans, Dry <sup>2</sup> , Succulent <sup>2</sup>	30 days 15 days	2.5 pints 2.5 pints	4.0 pints 4.0 pints	Yes Yes	Yes Yes	2, 13 (dry only)
Blackberries	45 days	2.5 pints	5.0 pints	No	Yes <sup>3</sup>	
Blueberries <sup>3</sup>	30 days	2.5 pints	5.0 pints	No	Yes	
<u>Brassica<sup>1</sup> including:</u> Broccoli (including Chinese & Raab), Brussels Sprouts, Cabbage (Bok Choy, Chinese Mustard, Napa), Cauliflower, Collards, Kale, Kohlrabi, Mustard Greens, Rape Greens	30 days	1.5 pints	3.0 pints	No	Yes	
<u>Bulb Vegetables<sup>1</sup> including:</u> Garlic, Leeks, Onions (Dry Bulb & Green), Shallots	30 days	1.5 pints	4.5 pints	No	Yes	
Canola/Crambe/Rapeseed <sup>3</sup>	60 days	2.5 pints	5.0 pints	No <sup>4</sup>	Yes	
Carrots	30 days	2.5 pints	5.0 pints	No	Yes	
Celery <sup>2</sup>	30 days	1.5 pints	3.0 pints	No	Yes	
Cherries (Sweet and Sour)	25 days	2.5 pints	5.0 pints	n/a	Yes	
Citrus	15 days	2.5 pints	10.0 pints	No <sup>5</sup>	No	
Clover	7 days before grazing, feeding, or cutting for (undried) forage	2.5 pints	6.5 pints	Yes	Yes	
Clover hay	20 days before grazing, feeding, or cutting for (dry) hay	2.5 pints	6.5 pints	Yes	Yes	
Corn <sup>1</sup> (Poast-protected™ field corn only)	60 days (grain or fodder) 45 days (forage and silage)	1.5 pints	3.0 pints	Yes	Yes	1,2,6,9, 10,13,15, 16,17, 28, 32
Cotton	40 days	2.5 pints	7.5 pints	No <sup>4</sup>	Yes	5 <sup>20</sup> , 25
Cranberries <sup>3</sup>	60 days	2.5 pints	5.0 pints	No	Yes	
<u>Cucurbits including:</u> Cantaloupes (all), Cucumbers, Gherkins, Honeydew Melons, Muskmelons (all), Pumpkins, Squash (all), Watermelons	14 days	1.5 pints	3.0 pints	No	Yes	
Dates (nonbearing)	1 year	2.5 pints	7.5 pints	n/a	Yes	
Deciduous Trees, Non-food Crop Areas, Fallow Land <sup>1</sup>	n/a	2.5 pints	n/a	No	Yes	
Endive	15 days	1.5 pints	3.0 pints	No	Yes	
Fescue, Tall <sup>1,6</sup>	n/a	2.5 pints	n/a	No	Yes	
Figs (nonbearing)	1 year	2.5 pints	7.5 pints	n/a	Yes	
Flax <sup>1,3</sup>	75 days	1.5 pints	4.0 pints	Yes <sup>3,4</sup>	Yes	5, 19
<u>Fruiting Vegetables<sup>1</sup> including:</u> Eggplants, Ground-cherries, Pepinos, Peppers (all), Tomatillos, Tomatoes	20 days	1.5 pints	4.5 pints	No <sup>7</sup>	Yes	18, 24 (tomatoes only)
Grapes	50 days	2.5 pints	5.0 pints	No <sup>8</sup>	Yes	
Lentils <sup>2,3</sup>	50 days	2.5 pints	4.0 pints	No	Yes	

Table 7. Crop Specific Restrictions and Limitations for Poast® Herbicide (continued)

Crop	Minimum Time From Application to Harvest (PHI)	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application	Tank Mix Partner
Lettuce, Leaf <sup>2</sup> Head <sup>2</sup>	15 days 30 days	1.5 pints 1.5 pints	3.0 pints 3.0 pints	No No	Yes Yes	
Mint	20 days	2.5 pints	5.0 pints	No	Yes	2, 5
Nectarines	25 days	2.5 pints	5.0 pints	n/a	Yes	
Olives (nonbearing)	1 year	2.5 pints	7.5 pints	n/a	Yes	
Orchard floor middles <sup>1,3</sup>	n/a	0.5 pint	0.5 pint	n/a	No	30
Peaches	25 days	2.5 pints	5.0 pints	n/a	Yes	
Peanuts	40 days	1.5 pints	2.5 pints	No <sup>4</sup>	Yes	2, 4, 31
Peas, Dry <sup>6</sup> Succulent	30 days 15 days	2.5 pints 2.5 pints	4.0 pints 4.0 pints	Yes Yes	Yes Yes	
Pistachios (nonbearing)	1 year	2.5 pints	7.5 pints	n/a	Yes	
Plums (nonbearing)	1 year	2.5 pints	7.5 pints	n/a	Yes	
Pome Fruits including: Apples, Crabapples, Pears, and Quince	14 days	2.5 pints	7.5 pints	No <sup>9</sup>	No	
Pomegranates (nonbearing)	1 year	2.5 pints	7.5 pints	n/a	Yes	
Potatoes <sup>1</sup> , Field <sup>2</sup> Sweet <sup>10</sup> (East U.S.) Sweet (West U.S.)	30 days 30 days 60 days	2.5 pints 2.5 pints 1.5 pints	5.0 pints 5.0 pints 5.0 pints	No <sup>7</sup> No <sup>7</sup> No <sup>7</sup>	Yes Yes Yes	18, 24
Prunes (nonbearing)	1 year	2.5 pints	7.5 pints	n/a	Yes	
Raspberries	45 days	2.5 pints	5.0 pints	No	Yes <sup>3</sup>	
Rhubarb <sup>2</sup>	30 days <sup>11</sup>	1.5 pints	3.0 pints	No	No	
Set Aside Conservation Land <sup>12</sup>	n/a	2.5 pints	7.5 pints <sup>1</sup>	n/a <sup>1</sup>	Yes	
Soybeans <sup>1, 13</sup>	75 days	2.5 pints <sup>14</sup>	5.0 pints	Only seed and hay <sup>4</sup>	Yes	2,4,7,8,11, 12,13,14,20 21,22,23, 26,27,29,32
Spinach <sup>2</sup>	15 days	1.5 pints	3.0 pints	No	Yes	
Strawberries <sup>1,15</sup>	7 days	2.5 pints	2.5 pints	No	Yes <sup>3</sup>	
Sugar Beets <sup>1</sup>	60 days	2.5 pints	5.0 pints	Yes <sup>16</sup>	Yes	3
Sunflowers <sup>1</sup>	70 days	2.5 pints	2.5 pints	No <sup>5</sup>	Yes	
Tobacco Seedbeds <sup>13, 17</sup>	n/a	1.0 pint	1.0 pint	No	No	
Tree Nuts <sup>1,16</sup>	15 days	2.5 pints	10.0 pints	No <sup>19</sup>	No	

Tank mix partners are as follows:

1. Atrazine
2. Basagran®
3. Betamix®/ Betanex®
4. Blazer®
5. Buctril®
6. Clarity
7. Classic®
8. Cobra®
9. Dual®
10. Dual II®

11. First Rate®
12. Flexstar®
13. Frontier®
14. Galaxy®
15. Guardsman®
16. Harness®
17. Laddok® S-12
18. Lexone®
19. MCPA
20. Pursuit®
21. Reflex®

22. Reliance STS
23. Resource
24. Sencor® DF
25. Staple®
26. Stellar®
27. Storm®
28. Surpass®
29. Synchrony STS
30. 2,4-D amine
31. 2,4-DB
32. 2,4-D (LVE)

<sup>1</sup> See Crop-Specific Information (pages 12-15) for more details.<sup>2</sup> Use crop oil concentrate or crop oil concentrate plus UAN or AMS according to temperature and humidity restrictions (see Vegetable Crops, page 14).<sup>3</sup> Not registered in California.<sup>4</sup> Processed meal may be fed from canola/crambe/rapeseed, cotton, flax, peanuts, soybeans, and sunflowers (also soaps/stock).<sup>5</sup> Pulp and waste may be fed to livestock.<sup>6</sup> For use in Alabama, Georgia, Kentucky, North Carolina, South Carolina, Tennessee, Virginia, and West Virginia only.<sup>7</sup> Potato and tomato waste may be fed to animals.<sup>8</sup> Pomace and raisin waste may be fed to animals.<sup>9</sup> Pressed or processed apple waste may be fed to animals.<sup>10</sup> Eastern U.S. includes AL, FL, GA, LA, MS, NC, SC, TN, TX and VA. Western U.S. includes AZ, CA, ID, NV, OR and WA.<sup>11</sup> Rhubarb grown only in IL, IN, MI, MN, and WI may be harvested up to 15 day PHI.<sup>12</sup> East of the Rocky Mountains only.<sup>13</sup> Use 2,4-D (LVE) for preplant burndown only.<sup>14</sup> The maximum rate per application in soybeans in California is 2.0 pints per acre.<sup>15</sup> Not registered in Florida.<sup>16</sup> Processed pulp and molasses may be fed to animals. Do not use sugar beet tops or roots for livestock grazing or feeding within 100 days after treatment.<sup>17</sup> Do not apply Poast in transplanted tobacco.<sup>18</sup> Tree nuts do not include pistachios.<sup>19</sup> Almond hulls may be fed to animals.<sup>20</sup> For use with Buctril resistant cotton only.

## IX. Crop-Specific Information

### Crops Grown For Seed

**Poast® herbicide** is recommended for use on all crops on this label when they are grown for seed production. Use the **Poast** rates given for each food crop listed in other sections on this label. Slight modifications in application methods may be required for certain seed crops due to crop canopy or different cultural methods from the corresponding food crop. Contact BASF or local authorities before modifying application methods to confirm that they do not conflict with labeling.

### Field Crops

Always add 1 pint of **Dash® HC**, **Sundance® HC** spray adjuvant, or 2 pints of oil concentrate per acre. Add 4-8 pints of UAN or 2.5 pounds of AMS to control crabgrass and all volunteer cereals. (UAN and AMS are not registered in California).

### Corn

**Only Poast Protected™ field corn hybrids are tolerant to Poast applications. Severe crop injury will occur to corn hybrids not designated as Poast Protected corn.**

Over-the-top applications of **Poast** in **Poast Protected** field corn may be made until the onset of pollen shed provided the appropriate preharvest intervals are met. Do not apply **Poast** after pollination occurs.

### Flax

#### **Poast + Buctril or MCPA or Bronate**

**Poast:** up to 1.5 pints per acre

**Buctril:** up to 1 pint equivalent per acre

**MCPA:** up to 0.25 pound acid equivalent per acre

**Bronate:** 0.9 pints per acre

**Buctril, MCPA or Bronate** applied with **Poast** may cause leaf burn, retarded growth, and delayed maturity of the crop. Some reduced grass control may be experienced with the above tank mixes. Conduct the **Compatibility Test for Mix Components**. See section VI. **Mixing Order** for details.

#### **Tank Mixing Restrictions (partial list)**

Do not delay spraying broadleaf weeds even though grassy weeds are not in the correct stage for treatment.

Do not add AMS or UAN solution to a tank mix of **Poast + Buctril or MCPA or Bonate**.

### Soybeans

**Poast + Basagran® + Blazer®, Galaxy, or Storm**  
(Not for use in California.)

When applying a tank mix by air, use a minimum of 10 gallons of total spray solution per acre. For these tank mixes, use a maximum of 1 pint of oil concentrate per acre. Use a maximum of 2 quarts of UAN per acre if velvetleaf is present. AMS at 1-2 pounds may be substituted for UAN. Conduct the **Compatibility Test for Mix Components**. See section VI. **Mixing Order** for details.

#### **Burndown:**

#### **Poast + 2,4-D LVE**

**Poast:** 0.5 pint per acre

**2,4-D LVE:** up to 1 pound a.i. per acre

Use only low volatile ester formulations of 2,4-D such as 2,4-D isooctyl ester. Note that the recommended

rate of 2,4-D (LVE) is calculated on an acid equivalent (a.e.) basis. Adjust the rates based on the concentration of 2,4-D (LVE) formulation used. Conduct the **Compatibility Test for Mix Components**. See section VI. **Mixing Order** for details.

#### **Tank Mix Specific Restrictions**

Do not plant soybeans until 7 days after treatment when using up to 0.5 pound per acre 2,4-D (LVE) or until 30 days after treatment when using up to 1.0 pound a.i. per acre 2,4-D (LVE). Make only one application of this tank mix per growing season.

Do not feed hay, forage, or fodder. Restrict livestock from grazing treated fields or cover crops.

Do not apply if rainfall is expected within 6 hours following application as weed control will probably be unsatisfactory.

Because all crops, such as sorghum, corn, small grains, cotton, soybeans, sugar beets, trees, shrubs, and ornamental grasses, such as turf, are extremely susceptible to **Poast** plus 2,4-D (LVE) tank mix, avoid all direct or indirect **postemergence** contact with any desired plant.

Do not spray if the wind is blowing toward desired sensitive plants, or at anytime when the wind exceeds 6 mph (refer to 2,4-D (LVE) label).

This tank mix does not control sedges or provide season-long control of hard-to-kill perennial weeds. Do not apply this tank mix during or following planting or after soybean emergence as severe soybean injury will result.

Do not use MSO with any tank mix combination.

### Sugar Beet

#### **Poast + Betamix/Betanex**

**Poast:** up to 1.5 pints per acre

**Betamix/Betanex:** up to 7.5 pints per acre

(Not for use in California)

A **Poast** and **Betamix/Betanex** tank mix can be applied when the specified annual grasses are less than 2 inches in length. Grasses of this size generally occur at the second application of the split treatment of **Betamix/Betanex**.

Conduct the **Compatibility Test for Mix Components**. See section VI. **Mixing Order** for details.

#### **Tank Mixing Restrictions (partial list)**

Do not apply this tank mix within 75 days of harvest.

The use of UAN solution or AMS with a **Poast + Betamix/Betanex** tank mix is not recommended.

Consult a BASF representative or local agricultural authority for more information.

Do not use this tank mix if grasses to be controlled include rhizome Johnsongrass, quackgrass, Bermudagrass, wirestem muhly, volunteer corn, shattercane, red rice or itchgrass.

### Sunflowers

Commercially released varieties of sunflower are tolerant to **Poast** at all stages of growth; however, leaf speckling has been occasionally observed on sunflowers with no corresponding reduction in vigor or growth. **Poast** is not recommended for use on sunflower inbred lines grown for seed because crop safety of these lines has not been adequately established.

### Tobacco

Apply **Poast** only at the seedbed stage of growth.

## Forage Crops

### Alfalfa, Birdsfoot Trefoil, Clover, Sainfoin

**Poast® herbicide** may be applied to seedling or established alfalfa and clover grown for hay, silage, green chop, direct grazing, or for seed.

**Mowing:** The best control of annual grasses can be achieved by applying **Poast** before grass weeds are mowed. Once a grass is mowed it becomes tougher to control, as much of the leaf surface may be removed, putting the grass under stress. In areas without a killing frost, some annuals can over-winter after having been mowed a number of times. These grasses can form large crowns and contain many viable buds. A large crown, even if it is an annual grass, may require repeated applications of **Poast** for partial or complete control.

#### Tank Mixing in Alfalfa and Birdsfoot Trefoil Only

##### **Poast + 2,4-DB**

**Poast:** 2.5 pints per acre

**2,4-DB:** up to 0.75 pound a.i. per acre

Some leaf yellowing and burning of the alfalfa may occur with this tank mix. Using 2,4-DB ester formulations may increase the severity of leaf injury. Additionally, in established alfalfa, 2,4-DB alone may cause twisting of stems and malformation of leaves. (Refer to 2,4-DB label.) Alfalfa plants will generally outgrow these temporary leaf injuries. See section VI. **Mixing Order** for details.

#### Tank Mix Specific Restrictions

Do not add UAN solution or AMS to a tank mix of **Poast + 2,4-DB**.

Do not use this tank mix unless the 60-day feeding, grazing, and harvesting restrictions on the 2,4-DB label can be observed.

Do not use this tank mix in the High and Rolling Plains of Texas, Western Oklahoma, Western Kansas, and Eastern New Mexico.

### **IRRIGATED ALFALFA, CLOVER, BIRDSFOOT TREFOIL, AND SAINFOIN:**

Irrigation practices can be very critical to the successful use of **Poast** and may be necessary to start grass weeds growing again. Generally, applications 2-4 days after an irrigation are most effective because:

- grasses resume active growth,
- grasses have less chance to grow too large,
- by waiting later, the clover or alfalfa begins to canopy and interferes with spray coverage.

Irrigation shortly after application (2 days) can be effective, but more consistent grass control is obtained when the irrigation is made before the application.

#### Annual Grass Control

Apply **Poast** at the grass sizes and rates indicated in **Tables 3 and 4**. If a grass has been cut, apply **Poast** after the regrowth reaches the minimum height (so there will be enough leaf area for absorption) and before it exceeds the maximum height indicated. Apply before the clover or alfalfa canopies cover the grasses and interfere with the spray coverage. Also, applications after a clover or alfalfa cutting may need to be timed to follow an irrigation or rainfall which will allow the grasses to regrow to a treatable size.

Some annual grasses are spring- and summer-germinating plants, while others are fall-germinating plants, and the time they are actively growing and most susceptible to **Poast** may vary from area to area. Also, some annuals germinate over a long time, and because control of small grasses is desired, applications after each weed flush may be needed. As a general guideline, spray spring- and summer-germinating grasses as early in the season as possible. The optimum application timing may occur very early in the spring after initial green-up. Spray fall-germinating weeds in the fall soon after they begin growing but before any killing frosts. Late fall applications may be less effective due to environmental changes, such as frosts or the onset of flowering.

### Interseeded Oats

Oats interseeded with clover, alfalfa, birdsfoot trefoil, and sainfoin may be killed by applying **Poast**. Their removal allows the seedling crops to grow with less competition. This application should be made before the oats get too large. Application made in the boot stage or later will not be as effective as when applied onto young oats.

#### Perennial Grass Control

**Poast** effectively controls or suppresses perennial grasses, such as Bermudagrass, johnsongrass, quackgrass, wirestem muhly, and perennial ryegrass. However, their growth characteristics are such that they are more difficult to control than annual grasses, especially in a perennial crop such as established alfalfa or clover. A program of repeated applications is usually necessary for best results.

The most economical way of controlling perennial grasses is to do so in the year of stand establishment before rhizomes or stolons become large and difficult to kill. The field should be disked before seeding to thoroughly fragment rhizomes or stolons.

In summer and fall seedings, cool season grasses (quackgrass, wirestem muhly, and perennial ryegrass) can become very competitive under cool fall conditions. Fall applications of **Poast® herbicide** will reduce late season grass growth and limit the ability of grasses to accumulate nutrient reserves in roots and rhizomes.

In established stands, it is important to begin applying in the spring when conditions favor active growth and before storage tissues have increased their nutrient reserves. Additional applications should be made on any grass regrowth in later cuttings.

### Set Aside Conversation Reserve Land, Fallow Acreage

**Broadleaf Cover Crops:** The growth of broadleaf cover crops such as alfalfa, clover, lespedeza, trefoils, and vetches will not be affected by **Poast**.

**Grass Cover Crops:** Most seeded grass crops such as oats, sudangrass, tall fescue, orchardgrass, bromegrasses, ryegrass, or timothy will be injured or killed by **Poast**, therefore, do not use **Poast** if injury to these grass cover crops is undesirable. Seeded grass cover crops may be injured or killed.

#### Restrictions and Limitations (partial list)

Do not harvest or graze cover crops other than alfalfa, clover, birdsfoot trefoil, or sainfoin treated with **Poast**.

13 7 20

Do not plant any other crop to be harvested for 120 days after application, unless **Poast**® herbicide is registered for use in that crop.

This use is applicable only for the Midwest, South, and Northeast areas (see maps in **Table 3**).

For alfalfa cover crops, do not apply **Poast** within 7 days of grazing, feeding, or cutting for (undried) forage, or within 14 days of cutting alfalfa for (dry) hay.

For alfalfa cover crops, do not apply more than a total of 6.5 pints of **Poast** per acre in one season.

## Fruit and Nut Crops

### Strawberries

A single application may not provide complete control of perennial grasses. The application rate for **Poast** on strawberries may be increased if the application rate does not exceed 2.5 pints per acre, per season.

**Poast** is not recommended for spring control of volunteer cereals that emerged the previous fall.

**Note:** Cultivate 14-21 days after application to aid control. Depending on environmental conditions and crop cultural system, season-long control may not always be obtained. However, competition from quackgrass will be reduced.

Not for use on strawberries in Florida.

### Tree Nuts

**Poast** may be used for grass control and suppression in bearing or nonbearing tree nuts. (Pistachios are not classified as tree nuts.) Tree nuts are very tolerant to **Poast** and **Poast** may be applied over the top of small, nonbearing trees or as a directed spray on larger trees.

Do not apply **Poast** with another pesticide whose label cautions against use with oil adjuvants.

## Interseeded Cover Crops

### Poast Activity on the Cover Crop

Grass cover crops controlled or suppressed by this use include wheat, oats, and barley, or any grass crop for which **Poast** is labeled. **Poast** will selectively control grass cover crops in seedling nongrass or broadleaf field, forage, or vegetable crops without injury. In addition, **Poast** will control any annual grasses that have emerged since planting. The slow-dying grass will provide a protective mulch for the primary crop seedlings for up to 3 weeks after applying **Poast**. This period will allow the crop to develop enough to become more tolerant to damage from wind-blown soil particles.

Apply **Poast** to cereals that are 3-4" in height (before tillering). Do not allow cereals to exceed this height as excessive competition and lack of control may occur.

## Nonbearing Crops and Noncrop Areas

For nonbearing crops, always add 1 quart of oil concentrate per acre.

## Deciduous Trees, Nonfood Crop Areas, Fallow Land

### DECIDUOUS TREES

#### NONFOOD CROP AREAS, FALLOW LAND:

**Poast** may be used in noncrop areas including rights-of-ways, roadsides and other paved areas, along fences and hedgerows, public buildings, recreation areas, industrial sites, storage yards, airports, electric transformer stations, pipeline pumping stations, sewage disposal areas, on potting and top soils, uncultivated agricultural areas, and general indoor or outdoor sites. **Poast** is not recommended for use on red sprangletop in California, Arizona, or western New Mexico.

**Notice to user:** Due to variability within species and in application techniques, neither the manufacturer nor the seller has determined whether or not **Poast** can be safely used on all varieties and species of nonbearing food crops, and other nonfood crops under all conditions. Therefore, determine if **Poast** can be used safely before broad use in the following manner:

On a small test area, apply the recommended rate of **Poast** on nonbearing or nonfood crop species or varieties under the conditions expected to be encountered. Any adverse conditions should be visible within 7 days.

#### TALL FESCUE GROWTH SUPPRESSION:

(Alabama, Georgia, Kentucky, North Carolina, South Carolina, Tennessee, Virginia, and West Virginia only)

Apply **Poast** to actively growing tall fescue after it has 4-6 inches of new growth, before the emergence of seedheads and before conifer bud break. Applications made from July 1 to mid-August may be less effective, especially if day temperatures reach 90° F. Tall fescue must be 1-year old before the first application of **Poast**.

Adequate coverage of the leaf surface is necessary for absorption of this herbicide. Thus, for optimum control, do not mow tall fescue turf for 30 days before or 14 days after applying **Poast**® herbicide.

**Rate:** Apply 1-1.25 pints of **Poast** per acre. For greater fescue suppression, up to 2.5 pints of **Poast** per acre can be used. Because of environmental differences at application, and growth differences of tall fescue, tall fescue control may exceed or fall short of that desired. Begin treating crops with **Poast** at the minimum recommended rate and adjust rates as local conditions and experience dictate. Additional applications may be made if extended growth suppression is desired. Tall fescue can also be treated with **Poast** by spot application.

#### ORCHARD FLOOR MIDDLES:

**Growth Management In Orchard Floor Middles**  
(Not registered for use in California)

##### Poast + 2,4-D

**Poast:** 0.5 pint per acre  
**2,4-D:** 2 pints per acre

**Poast** and 2,4-D dimethylamine can be used in a tank mix for growth management in orchard floor middles to reduce the number of mechanical mowings needed during a season. **Poast** and 2,4-D dimethylamine can be safely applied for growth management in the following cool season grasses and mixtures: Kentucky

bluegrass, perennial ryegrass, and tall fescue. Some degree of discoloration of the turf may occur. However, the turf will regrow and green up as effects of the treatment wear off. Make one application per season from the following options:

- **Poast**® herbicide and 2,4-D dimethylamine can be applied during the spring or summer when growth management is desired. Do not apply during bloom or within 3 days of a mowing.
- An optimal timing for application is after sod green up in the spring (before any mowing) or 3 days after the initial mowing of the season is made.
- A prebloom treatment is recommended as any broadleaf weeds such as dandelions can be controlled before they hamper fruit pollination. This treatment will provide 5-8 weeks of growth management depending on the sod makeup (e.g., grass species, amount of broadleaf weeds present, etc.), environmental conditions and the desired maintenance height of the middles.

See section **VI. Mixing Order** for details.

#### **Tank Mix Specific Restrictions**

Make no more than 1 application of this tank mix per growing season.

Do not apply if rainfall or irrigation is expected within 6 hours after application as growth management effects will probably be unsatisfactory.

Do not apply to a grass sod that is less than 2 years old.

Do not apply to newly established orchards. Trees must be at least 1 year old and in vigorous condition.

Do not apply this tank mix within 14 days of harvest of apples and pears.

Do not apply this tank mix to nonbearing stonefruits within one year of harvest.

### **Vegetable Crops**

Allow a minimum of 14 days between sequential applications.

Always add 2 pints of oil concentrate per acre.

However, under the following conditions, **Poast** plus oil concentrate should be used with caution due to potential leaf injury: when the temperature exceeds 90° F and the relative humidity is 60% or greater, or anytime the temperature exceeds 100° F, regardless of the humidity.

Do not use **Dash HC**® nor **Sundance**® HC spray adjuvants or methylated seed oils

### **Potatoes and Tomatoes**

In case of heavy infestations of quackgrass, use 2.5 pints of **Poast** per acre followed by 1.5 pints per acre sequentially if needed

Apply a tank mix of **Poast + Lexone DF** or **Sencor DF** to control mixed populations of annual grasses and broadleaf weeds listed as susceptible on the two product labels. See section **VI. Mixing Order** for details.

### **Poast + Lexone® DF or Sencor® DF herbicides** (Not applicable in California.)

**Poast:** refer to Table 6  
**Lexone DF or Sencor DF:**

- **for potatoes:** 8-10 ounces per acre (broadcast)
- **for tomatoes:** 5-8 ounces per acre (broadcast), 8-12 ounces per acre (directed spray).

#### **Tank Mix Specific Restrictions**

No tank mixes other than **Lexone**® herbicide or **Sencor**® herbicide are to be applied with **Poast**. Apply only if there have been at least 3 successive days of sunny weather before application or crop injury may occur.

Do not add UAN solution or AMS to a **Poast + Lexone** or **Sencor DF** tank mix.

Do not use this tank mix if grasses to be controlled include rhizome johnsongrass, quackgrass, Bermudagrass, wirestem muhly, volunteer corn or cereal, shattercane, red rice, or itchgrass.

Apply only to russetted or white-skinned varieties of potato that are not early maturing.

Do not apply this tank mix within 60 days of potato harvest.

Do not treat transplanted tomatoes within 14 days of transplanting. Tomatoes must have recovered from transplant shock and new growth must be evident.

Do not treat seeded tomatoes until plants have reached the 5-6 leaf stage.

## Returnable Container Operating Instructions.

### Prodigy™ System Operating Procedure

**Attention!** The Prodigy System is a pressurized delivery system. Do not attempt to open the container. Transfer product only by following these steps:

1. Install a male dry lock connector to the spray tank.
2. Uncoil the hose from the rack and connect the female dry lock connector (at the end of the hose attached to the tank) with the male dry lock connector installed on the spray tank.
3. Turn on the nitrogen gas supply.
4. Push down on the activation handle in the front near the meter until the handle is locked in the lower position allowing the manifold to fill with product and become pressurized. Some tanks do not have a handle; move on to the next step.
5. Turn the meter on by pressing the "ON/TOTAL" button.
6. Press "RESET" button to set current total to "0.00" if desired.
7. Turn the yellow product delivery valve counterclockwise (to horizontal) until the desired amount of product, as indicated on the measuring meter, has been discharged into the spray tank.
8. Turn the yellow product delivery valve clockwise (to vertical) to stop the discharge of product into your spray tank.
9. Lift the activation handle to the unlocked position (in front near the meter) to stop liquid and pressurization from flowing into the manifold. Some tanks do not have a handle; move on to the next step.
10. Turn off the nitrogen gas valve when the Prodigy System is not in use.
11. Hose draining: Starting at the yellow handle on the Prodigy Tank, grasp the hose and walk toward the receiving tank holding the hose level or higher than the dry lock connection allowing all of the product to drain out of the hose.
12. Disconnect the female dry lock connector on the tank hose from the male dry lock connector on the spray tank.
13. Recoil the hose onto the hose rack.
14. Be sure to turn off the nitrogen gas valve on the nitrogen cylinder when the Prodigy System operation is completed, or when the tank is empty, or when the tank is ready to be returned to the point of purchase.

Leave all product and bar code labels in place. Product labels must remain in place to comply with Department of Transportation regulations.

### Return Container Promptly to Distributor

The Prodigy System containers are tracked with bar codes and serial numbers. Distributors are responsible for the containers assigned to them. Return this container to the distributor from which it was purchased. Notify the distributor if the container cannot be returned by a specific time. The distributor is responsible for returning the container to BASF. The distributor will be charged for any container not returned within 30 days.

### 15-Gallon Returnable Container Operating Procedure


**Attention!** The 15-gallon container is a closed system. Do not try to remove the valve. The coupler required for product removal is available from your distributor. Do not use any other type of coupler. The coupler and probe are designed for one-way operation only. Never try to pump materials back into the container.

### Connection Steps


To engage and activate coupler:

1. Twist the containment cap counterclockwise breaking the tamper-evident seal.
2. Remove the cap from the container to expose the extractor valve.
3. Be sure the coupler handle is in the upward position.
4. Securely attach a hose or pump to the threaded connection. Be sure the air inlet has an air filter cap over the inlet or an air pressurization line screwed tightly into the inlet.
5. Place the coupler over the extractor valve and turn the coupler clockwise until it stops.
6. To secure the coupler, press the coupler handle downward completely until it is locked. (The handle cannot be locked if the coupler is incorrectly connected to the extractor valve. Do not force the handle. Start from Step 5 again.)
7. When the coupler handle is locked, the coupler is engaged and the system is open. You are now ready to begin pumping or the pressurization operation.

### Connection



### Connection




### Disconnection Steps

To remove coupler from container:

1. Lift the handle upward to stop the flow. Do not rotate the coupler.
  2. Vent the pressure by pulling the pressure release pin on the side of the coupler.
  3. Keep the handle in the upward position and turn the coupler counterclockwise.
  4. Remove the coupler by pulling it straight up. The coupler is now disconnected from the extractor valve.
  5. Wipe off the extractor valve with a cloth and replace the containment cap on the extractor valve after use or during any form of transportation.
  6. Flush the system with water or air.
  7. Wipe off the coupler with a cloth and store the coupler in a clean place.
  8. Properly dispose of cleaning towels or rinsate.
- Clean the outside of the container with soap and water before returning the container to the distributor. Leave all product and bar code labels in place. Product labels must remain in place to comply with Department of Transportation regulations.

### Disconnection





## Weeds listed in this label:

Common Name	Scientific Name
Barnyardgrass (Watergrass)	<i>Echinochloa crus-galli</i>
Bermudagrass (Wiregrass)	<i>Cynodon dactylon</i>
Crabgrass, Large	<i>Digitaria sanguinalis</i>
, Smooth	<i>Digitaria ischaemum</i>
Cupgrass, Southwestern	<i>Eriochloa gracilis</i>
, Woolly	<i>Eriochloa villosa</i>
Fescue, Tall	<i>Festuca arundinacea</i>
Foxtail, Giant (Pigeongrass)	<i>Setaria faberi</i>
, Green	<i>Setaria viridis</i>
, Yellow	<i>Setaria glauca</i>
Goosegrass	<i>Eleusine indica</i>
Itchgrass	<i>Rottboellia exaltata</i>
Johnsongrass	<i>Sorghum halepense</i>
Junglerice	<i>Echinochloa colonum</i>
Millet, Wild Proso	<i>Panicum miliaceum</i>
Muhly, Wirestem	<i>Muhlenbergia frondosa</i>
Oats, Tame	<i>Avena sativa</i>
, Wild	<i>Avena fatua</i>
Orchardgrass	<i>Dactylis glomerata</i>
Panicum, Browntop	<i>Panicum fasciculatu</i>
, Fall	<i>Panicum dichotomiflorum</i>
, Texas	<i>Panicum texanum</i>
Quackgrass	<i>Agropyron repens</i>
Red Rice	<i>Oryza sativa</i>
Ryegrass, Annual	<i>Lolium multiflorum</i>
, Perennial	<i>Lolium perenne</i>
Sandbur, Field	<i>Cenchrus incertus</i>
Shattercane/Wildcane	<i>Sorghum bicolor</i>
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>
Sprangletop, Red	<i>Leptochloa filiformis</i>
Volunteer Barley	<i>Hordeum vulgare</i>
Corn	<i>Zea mays</i>
Oats	<i>Avena sativa</i>
Rye	<i>Secale Cereale</i>
Wheat	<i>Triticum aestivum</i>
Witchgrass	<i>Panicum capillare</i>

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF Corporation ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

*Basagran, Galaxy, Laddok, and Poast* are registered trademarks of BASF AG.

*Blazer, Clarity, Dash, Frontier, Guardsman, and Storm* are registered trademarks and *Poast Protected* is a trademark of BASF Corporation.

*Betamix and Betanex* are registered trademarks of Schering Ag.

*Buctril and Bronate* are registered trademarks of Rhône-Poulenc AG Company.

*Classic, Lexone, Reliance, Staple, and Synchrony* are registered trademarks of E.I. DuPont de Nemours and Company.

*Cobra* is a registered trademark of Valent USA Corporation.

*Flexstar, Reflex, and Surpass* are registered trademarks of Zeneca Inc.

*Dual and Dual II* are registered trademarks of Novartis Corporation.

*FirstRate* is a registered trademark of DowElanco.

*Harness* is a registered trademark of Monsanto Company.

*Pursuit* is a registered trademarks of American Cyanamid Company.

*Resource and Stellar* are registered trademarks of Valent U.S.A. Corporation.

*Sencor* is a registered trademark of Bayer AG.

*Sundance* is a registered trademark of Rosen's, Inc.

© 1998 BASF Corporation

All rights reserved.

NVA 98-4-25 (10)

BASF Corporation  
P.O. Box 13528  
Research Triangle Park, NC 27709

**BASF**

# Poast<sup>®</sup>

herbicide

**For use in leafy, root and tuberous vegetable and caneberry crops and globe artichoke**

EPA Registration Number 7969-58

All applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of application.

## Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

## Application Instructions

**Poast<sup>®</sup> herbicide** may be applied to the crops listed in **Table 1**. Apply to actively growing grasses. In irrigated areas, it may be necessary to irrigate prior to treatment with **Poast** to ensure weeds are growing actively. Always adjust spray pressure, spray volume and height of spray boom to ensure penetration of plant canopy and thorough coverage of grasses to be controlled. For maximum use rate per application and per season and for the minimum time for application to harvest, see **Table 1**.

## Additives

Always add two pints of oil concentrate per acre.

## Restrictions and Limitations

Avoid all direct or indirect contact with any desired grass crop unless otherwise recommended on the **Poast** label.


Do not apply to grasses or crops under stress such as stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control may result. Irrigation may be needed.

Do not apply to crops that show injury (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced or prolonged.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Poast** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. BASF does not recommend using tank mixes other than those listed on BASF labeling. Local agricultural authorities may be a source of information when using other than BASF recommended tank mixes.

*Poast is a registered trademark of BASF AG.*

© 1998 BASF Corporation  
All rights reserved.  
NVA 98-4-25-0187



**Table 1. Crop-Specific Restrictions and Limitations for Poast® Herbicide**

Crop	Minimum Time From Application To Harvest (PHI)	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application
<u>Head &amp; Petiole Type Vegetables</u> Cardoon, Celery, Celery (Chinese), Celtuce, Fennel (Florence), Lettuce (Head), Radicchio, Rhubarb <sup>1,2</sup> , Swiss Chard	30* days	1.5 pints	3.0 pints	No	Yes
<u>Leafy Vegetables</u> Amaranth, Arugula, Chervil, Chrysanthemum (Edible, Garland), Cilantro, Corn Salad, Cress (Garden, Upland), Dandelion, Dock, Endive (Escarole), Lettuce (Leaf), Orach, Parsley, Purslane (Garden, Winter), Spinach, Spinach (New Zealand, Vine)	15 days				
<u>Root &amp; Tuberous Corms</u> Arracacha, Arrowroot, Artichoke (Chinese, Jerusalem), Canna (Edible), Cassava (Bitter, Sweet), Chayote Root, Chufa, Dasheen(Taro), Ginger, Leren, Potato, Sweet Potato <sup>3</sup> , Tanier, Turmeric, Yam Bean, Yam (True)	30 days	2.5 pints	5.0 pints	No <sup>4</sup>	Yes
Beet (Garden)	60 days				
<u>Caneberries</u> (All varieties and/or hybrids of these) Blackberry, Raspberry (Red, Black), Loganberry, Youngberry	45 days	2.5 pints	5.0 pints	No	Yes <sup>5</sup>
Globe Artichoke	7 days	2.5 pints	5.0 pints	No	Yes

<sup>1</sup> Rhubarb grown only in IL, IN, MI, MN, and WI may be harvested 15 days after treatment.

<sup>2</sup> Aircraft application not registered.

<sup>3</sup> In the West, the PHI of sweet potato is 60 days. Western U.S. includes AZ, CA, ID, NV, OR and WA.

<sup>4</sup> Potato waste may be fed to animals.

<sup>5</sup> Not registered in California.

SAS Corporation  
P.O. Box 13528  
Research Triangle Park, NC 27709

**SAS**

## Supplemental Label

# Poast<sup>®</sup>

herbicide

## For Use as a Preplant Burndown Treatment in Corn

EPA Reg. No 7969-58

All applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of application.

### Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Application Instructions

Follow the application timings for weed sizes as specified on each product label. Follow the spray volume and pressure recommendations as described on the **Poast<sup>®</sup> herbicide** label.

Apply **Poast** at 1.0 pint per acre at 30 or more days prior to planting corn to actively growing grasses. Use additives according to the **Spray Additives** section of this label.

Read and follow the applicable **Restrictions and Limitations** and **Directions For Use** on all products involved in tank mixing. Refer to specific labels for the spectrum of susceptible weeds. The most restrictive labeling applies to tank mixes.

### **Poast + Marksman<sup>®</sup> herbicide**

**Poast** 1.0 pint per acre

**Marksman** 3.5 pints per acre

Apply at 30 or more days prior to planting corn to actively growing weeds and grasses. This tank-mix provides the combined benefits of early burndown and 1.0 pound active ingredient of atrazine preemergence. Use additives according to the **Spray Additives** section of this label.

### **Poast + Clarity<sup>®</sup> herbicide**


**Poast** 1.0 pint per acre

**Clarity** 1.0 pint per acre

Apply at 30 or more days prior to planting corn to actively growing weeds and grasses. Use additives according to the **Spray Additives** section of this label.

### Spray Additives

Use 2.0 - 4.0 pints of UAN or 1.0 - 2.0 pounds of spray grade ammonium sulfate with crop oil concentrate or methylated seed oil at 1.0 - 2.0 pints per acre. Also refer to the respective labels for specific recommendations on additives.



Clarity and Marksman are registered trademarks of BASF Corporation.  
Poast is a registered trademark of BASF AG.

© 1998 BASF Corporation  
All rights reserved.  
NVA 98-4-25-0179

BASF Corporation  
P.O. Box 13528  
Research Triangle Park, NC 27709

# BASF