



The Chemical Company

Poast®

herbicide

Supplemental Labeling

For use in Borage, Buckwheat, Dill, Okra and Root Vegetables

EPA Registration Number 7969-58

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of application.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

I. General Information

Poast® herbicide may be applied to the crops listed in **Table 1**. Apply to actively growing grasses. In irrigated areas, it may be necessary to irrigate prior to treatment with Poast to ensure weeds are growing actively. See **Poast** label for complete list of annual and perennial grasses controlled with use of this herbicide.

Always adjust spray pressure, spray volume and height of spray boom to ensure thorough coverage of grasses to be controlled.

For maximum use rate per application and per season and for the minimum time from application to harvest, see **Table 1**.

II. Restrictions and Limitations

Do not plant other crops to be harvested for 30 days after application unless **Poast** is registered for use on that crop.

Poast should only be applied as a directed spray to the grove floor for bearing pistachio trees. Avoid all direct or indirect contact with any desired grass crop unless otherwise recommended on the **Poast** label.

Do not apply to grasses or crops under stress such as stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control may result. Irrigation may be needed.

Do not apply to crops that show injury (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced or prolonged.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Poast** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. BASF does not recommend using tank mixes other than those listed on BASF labeling. Local agricultural authorities may be a source of information when using other than BASF recommended tank mixes.




Table 1. Crop-Specific Restrictions and Limitations for Poast® herbicide

Crop	Minimum Time From Application to Harvest (PHI)	Maximum Rate Per Acre per Application	Maximum Rate Per Acre per Season	Livestock Grazing or Feeding
Borage	23 days	2.5 pints	5 pints	n/a
Buckwheat	21 days	2.5 pints	5 pints	n/a
Dill	14 days	2.5 pints	5 pints	n/a
Okra	14 days	1.5 pints	5.5 pints	n/a
Root vegetables (except sugar beet) subgroup: Edible burdock Celeriac Chervil, turnip-rooted Chicory Ginseng Oriental radish Parsley, turnip-rooted Parsnip Radish Rutabaga Salsify Black Salsify Spanish salsify Skirret Turnip	14 days	2.5 pints	2.5 pints	n/a
Beet, garden	60 days	2.5 pints	5 pints	n/a
Carrot	30 days	2.5 pints	5 pints	Yes
Horseradish	60 days	2.5 pints	5 pints	n/a

n/a = not applicable

Aerial application is permitted for all labeled crops.

Poast is a registered trademark of BASF

© 2005 BASF Corporation
All rights reserved

007969-00180.20050223.NVA 2005-04-025-0066.pdf

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

