

SUPPLEMENTAL LABELING

RONILAN^R FUNGICIDE FL
(EPA Reg. No. 7969-62)

IN STONEFRUIT

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of application.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

Ronilan FL is effective for the control of brown rot blossom and twig blight, and fruit brown rot of stonefruit including apricots, cherries, nectarines, peaches, plums, and prunes. Ronilan FL is a contact fungicide, therefore, thorough coverage of plant parts to be protected is essential for effective disease control.

TIME AND RATE OF APPLICATION

Depending on conditions which favor disease development, one to three applications should be made during bloom for control of brown rot blossom and twig blight. One or two pre-harvest applications should be made for control of fruit brown (See Table).

ACCEPTED
APR 24 1987
7969-62

STONEFRUIT

APPLICATION RATES AND TIMING FOR CONTROL OF BROWN ROT (MONILINIA SPP.)

CROP	<u>Pints of Ronilan FL Per</u> 100 Gallons	<u>FL Per</u> Acre	TIMING
<p>Apricots Cherries Nectarines Peaches Plums Prunes</p>	<p>0.75 to 1.0</p> <p>Rate is Based on 200 GPA</p> <p>Do not exceed 2 pints product per acre per application.</p>	<p>1.5 to 2.0</p>	<p><u>Brown Rot Blossom & Twig Blight</u></p> <p>Apply at <u>Early Bloom</u> Apricots - Red Bud Peaches & Nectarines - Pink Bud Cherries - Early Popcorn Plums and Prunes - Green Tip</p> <p>Apply again at <u>Full Bloom</u> if conditions persist which favor disease development.</p> <p>Apply again at <u>Petal Fall</u> if conditions persist which favor disease development.</p> <p>Do not apply more than 3 bloom treatments.</p> <p><u>Fruit Brown Rot</u></p> <p>Apply when conditions favor disease development within 3 weeks of harvest. One additional application should be made 7 to 14 days later if these conditions persist.</p> <p>Do not apply more than 2 pre- harvest treatments.</p> <p>Do not apply within 3 days of harvest.</p>

2/4

METHOD OF APPLICATION

Rates of Ronilan FL per 100 gallons are based on an application of 200 gallons per acre of dilute spray. Refer to the Rates per Acre if other than 200 gallons per acre is applied.

Apply Ronilan FL as a spray with ground equipment, using sufficient water to obtain THOROUGH COVERAGE (50-400 gallons per acre).

Under high disease pressure or in conditions very favorable for disease development, use higher treatment rates, multiple applications, and shorter application intervals. For large mature trees, use higher treatment rates.

Ronilan FL is physically compatible with most pesticides used in a tank mix on stonefruit. To assure compatibility of Ronilan FL with other products, add correct proportions in a small container with water. Mix thoroughly and let the mixture stand for five minutes. The mixture is compatible if the combination remains mixed, or can be readily remixed.

Mixing Instructions: Partially fill spray tank with clean water and begin agitation. Measure the required amount of Ronilan FL and add to the tank. Fill the tank to the total volume required. Maintain agitation to keep the material in suspension and apply with properly calibrated spray equipment.

RESTRICTIONS AND LIMITATIONS FOR STONEFRUITS

Do not apply Ronilan FL within 3 days of harvest.

Do not apply more than 2 pints of Ronilan FL per acre per application.

Do not apply more than 10 pints of Ronilan FL per acre per season (maximum of 6 pints at bloom and 4 pints pre-harvest).

Do not apply Ronilan FL during rain. Apply when conditions will permit spray to dry on the plants.

CONDITIONS OF SALE AND WARRANTY

The Directions for Use of this product reflects the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. In no case shall BASF or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. BASF and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

EPA ESTABLISHMENT NO. 7969-WG-1

Ronilan is a registered trademark of BASF AG Made in West Germany.