

BASF

RT Date: 1-12-94

Conclude™ B

herbicide

Postemergence Herbicide

Active Ingredients:

Sodium salt of bentazon: sodium (3-isopropyl-1H-2,1,3-benzothiadiazin-4(3H)-one-2,2-dioxide*29.2%

Sodium salt of acifluorfen: sodium 5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoate13.4%

Inert Ingredients57.4%

Total100.0%

*Equivalent to 2.67 pounds per gallon bentazon: 3-isopropyl-1 H-2,1,3-benzothiadiazin-4-(3H)-one 2,2-dioxide; and 1.33 pounds per gallon of sodium acifluorfen, sodium 5-[2-chloro-4-(trifluoromethyl) phenoxy]-2-nitrobenzoate.

EPA Reg. No. 7969-76

KEEP OUT OF REACH OF CHILDREN.

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to read it to you in detail.)

Statement of Practical Treatment

If in eyes: Flush with large amounts of water for at least 15 minutes. Get medical attention. If on skin: Wash with plenty of soap and water. Consult a physician if irritation persists. If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Note to physician: Emesis is recommended.

See inside booklet for complete Directions For Use and Conditions of Sale and Warranty.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions For Use section for information about this standard.

Net contents 7.5 gallons

BASF Corporation
P.O. Box 13528, Research Triangle Park, NC, 27709

BEST AVAILABLE COPY

Specimen Label

**Precautionary Statements
Hazards to Humans (And
Domestic Animals):**

Causes irreversible eye damage. Harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes. Avoid breathing vapor or spray mist and contact with skin or clothing. This product may cause an allergic skin response.

**Personal Protective Equipment
(PPE)**

Applicators and other handlers must wear:

- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations
Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

In Case of Emergency

In case of large-scale spillage regarding this product, call:

CHEMTREC 800-424-9300
BASF Corporation 800-832-HELP

In case of medical emergency regarding this product, call:

1. Your local doctor for immediate treatment
2. Your local poison control center (hospital)
3. BASF Corporation 800-832-HELP

**Directions For Use —
Tank Mix of Conclude B
and Conclude G**

(Hereafter referred to as Conclude)

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry to treated areas during the REI of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or vitron ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

General Information

Conclude is intended for the early postemergence control of a wide spectrum of broadleaf weeds and annual grasses in soybeans (See Table 1).

General Use Area

Use only in the following areas: Arkansas, Indiana (Only in Crawford, Daviess, Dubois, Gibson, Knox, Martin, Perry, Pike, Posey, Spencer, Vanderburgh, and Warrick counties), Kentucky, Louisiana, Michigan, Mississippi, Missouri (bootheel counties of Butler, Dunklin, Mississippi, New Madrid, Pemiscott, Ripley, Scott, and Stoddard), and Tennessee.

BEST AVAILABLE COPY

Maximum Weed Height: Controlled by Tank Mixing Conclude B (1.5 pints per acre) and Conclude G (1.5 pints per acre) with Crop Oil Concentrate (1 pint per acre)

3 of 9

Broadleaves	Maximum Weed Height	Grasses	Maximum Weed Height
Bristly Starbur	3"	Barnyardgrass	4"
Carpetweed	2"	Broadleaf Signalgrass	4"
Cocklebur	6"	Crabgrass, Large	4"
Common Lambsquarters	2"	Crabgrass, Smooth	4"
Common Ragweed	3"	Foxtail, Giant	4"
Crotalaria	6"	Foxtail, Green	4"
Croton, Tropic	<2"	Foxtail, Yellow	4"
Woolly	<2"	Goosegrass	4"
Giant Ragweed	6"	Johnsongrass (seedling)	4"
Jimsonweed	6"	Junglegrass	4"
Ladysthumb	6"	Panicum, Browntop	4"
Morningglories	2"	Panicum, Fall	4"
Nightshade, Black	2"	Panicum, Texas	4"
Pigweed, Redroot	2"	Red Spangletop	4"
Smooth	3"	Witchgrass	4"
Prickly Sida/Teaweed*	2"	Woolly Cupgrass	4"
Redweed	3"		
Sesbania	6"		
Smartweed, Pennsylvania	4"		
Spurred Anoda*	2"		
Texasweed	3"		
Velvetleaf	2"		
Vernice Mallow	2"		
Wild Mustard	4"		
		Perennials: (top growth suppression)	Maximum Weed Height
		Canada Thistle	6"
		Johnsongrass (Rhizome)	4"
		Yellow Nutsedge	4"

For new germination or perennial regrowth, follow up no sooner than 15 days later with Basagran® herbicide, Poast Plus® herbicide, Storm® herbicide, or Blazer® herbicide. Refer to the respective labels for Directions For Use.
* Control may be inconsistent. A later application of Basagran may be necessary. (See Basagran label.)

Storage and Disposal

Keep from freezing. Store above 32°F.

Do not contaminate water, food or feed by storage or disposal.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law.

If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Triple rinse container (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Mode of Action:

Conclude is effective through postemergence contact and systemic activity. Weeds must be thoroughly covered with spray. Large crop-and-weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage.

Crop Tolerance:

All soybean varieties are tolerant to Conclude at all stages of growth. Leaf speckling may occur but plants generally outgrow this condition within 10 days.

Cultivation:

Do not cultivate before application or within five days after application of Conclude. Cultivation may put weeds under stress and reduce control obtained. A timely cultivation 2-3 weeks after applying Conclude may assist weed control in soybeans grown in rows more than 10 inches apart.

Application Rate and Timing

Apply Conclude B and Conclude G at the rate of 1.5 pints each per acre at early postemergence when weeds are small and actively growing (generally when soybeans are in the 2nd to 3rd trifoliate leaf stage of growth). Add oil concentrate at a rate of 1 pint per acre. Under excessively dry, wet, or cold conditions, which may reduce herbicidal activity, add oil concentrate at the rate of 2 pints per acre.

Mixing

Fill tank of a thoroughly clean sprayer half to two-thirds full with clean water. Start agitation and add the recommended amounts of product in the following order: Conclude B, oil concentrate, and Conclude G. Then add the remaining quantity of water. Maintain constant agitation during application. After mixing, spray within 48 hours.

BEST AVAILABLE COPY

Ground Application

Use a minimum of 10-20 gallons of water per broadcast acre at a minimum of 40 psi pressure (measured at the boom, not at the pump or in the line) to ensure adequate spray coverage. Use standard high pressure pesticide hollow cone or flat fan nozzles spaced 18-20 inches apart. Do not use flood, whirl chamber, or controlled droplet applicator nozzles.

Aerial Application

Use a minimum of 5 gallons of water per acre and a maximum of 40 psi pressure. To obtain uniform coverage and to avoid drift hazards, the following application equipment and practices should be used:

Nozzle Type: Use only diaphragm-type nozzles producing cone or fan spray patterns.

Nozzle Height: 6-10 feet above crop.

Nozzle Orientation: Nozzles must be oriented so as to discharge straight back with the air stream (opposite the direction of travel of the aircraft) and not more than 20° downwind.

Nozzles must be located no farther out than $\frac{3}{4}$ the distance from the center of the aircraft to the end of the wing or rotor.

Do not apply Conclude by aircraft when wind is blowing at a velocity above 10 mph. Coarse sprays (larger droplets) are less likely to drift.

Do not apply Conclude by air if ornamental or sensitive nontarget crops such as cotton, sugar beets, sunflowers, or okra are within 200 feet downwind.

Applicator must follow the most restrictive use cautions to avoid drift hazard and must follow labeling as well as applicable state and local regulations and ordinances.

Procedure For Cleaning Spray Equipment

Clean sprayer thoroughly before and after application of Conclude, particularly if a herbicide with the potential to injure crops was previously used. Consult the label of the previously used herbicide for cleaning instructions. If no instructions are available, the steps listed below are suggested for thorough cleaning of spray equipment prior to or following application of Conclude.

Step 1: Hose down thoroughly the inside as well as the outside of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of this rinse water.

Step 2: Refill tank with water while adding 1 gallon household ammonia or 1 pint household dishwashing detergent or 1 pound of dishwasher detergent per 100 gallons of water. Or add a commercial sprayer cleaner according to the manufacturer's directions. Operate the pump to circulate the detergent solution through the sprayer system for 5-10 minutes and discharge a small amount of solution through the boom and nozzles. Let the solution stand for 24 hours.

Step 3: Flush the detergent solution out of the spray tank through the boom.

Step 4: Remove the nozzles and screens and flush the system with two tankfuls of water.

Restrictions and Limitations

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive label applies when using tank mixes.

Only one application of Conclude per acre per season should be made.

Do not apply to weeds under stress, such as stress due to lack of moisture, previous herbicide injury, mechanical injury or cold temperatures, as unsatisfactory weed control

could result.

Do not apply if rainfall or irrigation is expected soon after application.

Physical incompatibility, reduced weed control, or crop injury may result from mixing Conclude with other pesticides (fungicides, herbicides, insecticides or miticides) additives or fertilizers not recommended on the label. BASF does not recommend the use of Conclude in tank mixes other than those listed on BASF labels, supplemental labels, or technical bulletins. Local agricultural authorities may be a source of information when using other BASF recommended combinations.

Do not apply Conclude to soybeans that show injury, leaf phytotoxicity, and/or other prior herbicide applications because this injury may be enhanced or prolonged.

Do not apply Conclude through any type of irrigation system.

Do not apply to soybeans within 75 days of harvest.

Do not use treated plants for feed or forage.

In case of crop failure, only soybeans or peanuts may be immediately be replanted.

Crop rotation restriction: Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in field treated with Conclude for 18 months following treatment.

Do not apply more than 2 pints of Basagran or 1 pint of Blazer following an application of Conclude.

Do not apply more than 1.5 pints of Storm following an application of Conclude.

Do not apply sequential applications of Blazer, Basagran, or Poast Plus within 15 days following the application of Conclude.

After application of Conclude, do not apply more than 3.5 pints of Poast per acre, per season.

Do not use selective application equipment such as recirculating sprayers, wiper applicators, or shielded applicators when applying Conclude.

4 2 9

BLANK PAGE

6 7 9
Conditions of Sale and Warranty
The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the

Directions For Use, subject to the inherent risks, referred to above.

BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Basagran is a registered trademark of BASF AG.

Conclude B and Conclude G are trademarks of BASF Corporation.

Storm, Poast Plus, and Blazer are registered trademarks of BASF Corporation.

© 1994 BASF Corporation

XX ----- JJ9154 A1661293


BEST AVAILABLE COPY

BASF Corporation
PO. Box 13528
Research Triangle Park, NC 27709

BASF

Supplemental Labeling

Storm[®]
herbicide



RT: 12-29-93

Tank mix with Poast[®] herbicide for postemergence use in soybeans

EPA Reg. No 7969-76

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

General Information

A tank mix of Storm[®] herbicide plus Poast[®] herbicide is intended for the early postemergence control of a wide spectrum of broadleaf weeds and annual grasses in soybeans (See Table 1).

General Use Area

Use only in the following areas: Arkansas, Kentucky, Indiana (Only in Crawford, Daviess, Dubois, Gibson, Knox, Martin, Perry, Pike, Posey, Spencer, Vanderburgh, and Warrick counties), Louisiana, Michigan, Mississippi, Missouri (boothel counties of Butler, Dunklin, Mississippi, New Madrid, Pemiscott, Ripley, Scott, and Stoddard), and Tennessee.

Storage and Disposal

Keep from freezing. Store above 32°F.

Do not contaminate water, food or feed by storage or disposal.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law.

If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide

or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Triple rinse container (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Mode of Action:

A tank mix of Storm plus Poast is effective through postemergence contact and systemic activity. Weeds must be thoroughly covered with spray. Large crop-and-weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage.

Crop Tolerance:

All soybean varieties are tolerant to A tank mix of Storm plus Poast at all stages of growth. Leaf speckling may occur but plants generally outgrow this condition within 10 days.

Cultivation:

Do not cultivate before application or within five days after application of the tank mix of Storm plus Poast. Cultivation may put weeds under stress and reduce control obtained. A timely cultivation 2-3 weeks after applying the tank mix of Storm plus Poast may assist weed control in soybeans grown in rows more than 10 inches apart.

Application Rate and Timing

Apply the tank mix of Storm plus Poast at the rate of 1.5 pints each per acre at early postemergence when weeds are small and actively growing (generally when soybeans

are in the 2nd to 3rd trifoliate leaf stage of growth). Add oil concentrate at a rate of 1 pint per acre. Under excessively dry, wet, or cold conditions, which may reduce herbicidal activity, add oil concentrate at the rate of 2 pints per acre.

Mixing

Fill tank of a thoroughly clean sprayer half to two-thirds full with clean water. Start agitation and add the recommended amounts of product in the following order: Storm, oil concentrate, and Poast. Then add the remaining quantity of water. Maintain constant agitation during application. After mixing, spray within 48 hours.

Ground Application

Use a minimum of 10-20 gallons of water per broadcast acre at a minimum of 40 psi pressure (measured at the boom, not at the pump or in the line) to ensure adequate spray coverage. Use standard high pressure pesticide hollow cone or flat fan nozzles spaced 18-20 inches apart. Do not use flood, whirl chamber, or controlled droplet applicator nozzles.

Aerial Application

Use a minimum of 5 gallons of water per acre and a maximum of 40 psi pressure. To obtain uniform coverage and to avoid drift hazards, the following application equipment and practices should be used:

Nozzle Type: Use only diaphragm-type nozzles producing cone or fan spray patterns.

Nozzle Height: 6-10 feet above crop.

Nozzle Orientation: Nozzles must be oriented so as to discharge

straight back with the air stream (opposite the direction of travel of the aircraft) and not more than 20° downwind.

Nozzles must be located no farther out than 3/4 the distance from the center of the aircraft to the end of the wing or rotor.

Do not apply the tank mix of Storm plus Poast by aircraft when wind is blowing at a velocity above 10 mph. Coarse sprays (larger droplets) are less likely to drift.

Do not apply the tank mix of Storm plus Poast by air if ornamental or sensitive nontarget crops such as cotton, sugar beets, sunflowers, or okra are within 200 feet downwind. Applicator must follow the most restrictive use cautions to avoid drift hazard and must follow labeling as well as applicable state and local regulations and ordinances.

Procedure For Cleaning Spray Equipment

Clean sprayer thoroughly before and after application of the tank mix of Storm plus Poast, particularly if a herbicide with the potential to injure crops was previously used. Consult the label of the previously used herbicide for cleaning instructions. If no instructions are available, the steps listed below are suggested for thorough cleaning of spray equipment prior to or following application of the tank mix of Storm plus Poast.

Step 1: Hose down thoroughly the inside as well as the outside of equipment while filling the spray tank half full of water.

Flush by operating sprayer until the system is purged of this rinse water.

Step 2: Refill tank with water while adding 1 gallon household

ammonia or 1 pint household dishwashing detergent or 1 pound of dishwasher detergent per 100 gallons of water. Or add a commercial sprayer cleaner according to the manufacturer's directions. Operate the pump to circulate the detergent solution through the sprayer system for 5-10 minutes and discharge a small amount of solution through the boom and nozzles. Let the solution stand for 24 hours.

Step 3: Flush the detergent solution out of the spray tank through the boom.

Step 4: Remove the nozzles and screens and flush the system with two tankfuls of water.

Table 1: Maximum Weed Heights Controlled by Tank Mixing Storm (1.5 pints per acre) and Poast (1.5 pints per acre) with Crop Oil Concentrate (1 pint per acre)

Broadleaves:	Max. Weed Height	Grasses:	Max. Weed Height	Perennials: (top growth suppression)	Max. Weed Height
Bristly Starbur	3"	Barnyardgrass	4"	Canada Thistle	6"
Carpetweed	2"	Broadleaf Signalgrass	4"	Johnsongrass (Rhizome)	4"
Cocklebur	6"	Crabgrass, Large	4"	Yellow Nutsedge	4"
Common Lambsquarters*	2"	Crabgrass, Smooth	4"		
Common Ragweed	3"	Foxtail, Giant	4"		
Crotalaria	6"	Foxtail, Green	4"		
Oxton, Tropic	<2"	Foxtail, Yellow	4"		
Woolly	<2"	Goosegrass	4"		
Giant Ragweed	6"	Johnsongrass (seedling)	4"		
Jimsonweed	6"	Junglegrass	4"		
Ladysthumb	6"	Panicum, Browntop	4"		
Morningglories	2"	Panicum, Fall	4"		
Nightshade, Black	2"	Panicum, Texas	4"		
Pigweed, Redroot	2"	Red Spangletop	4"		
Smooth	3"	Witchgrass	4"		
Prickly Sida/Tea Weed*	2"	Woolly Cupgrass	4"		
Redweed	3"				
Sesbania	6"				
Smartweed, Pennsylvania	4"				
Spurred Anoda*	2"				
Texasweed	3"				
Velvetleaf*	2"				
Verice Mallow	2"				
Wild Mustard	4"				

For new germination or perennial regrowth, follow up no sooner than 15 days later with Basagran, Poast Plus, Storm, or Blazer. Refer to the respective labels for Directions For Use.

* Control may be inconsistent. A later application of Basagran may be necessary. (See Basagran label.)

Restrictions and Limitations

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive label applies when using tank mixes.

Only one application of the tank mix of Storm plus Poast per acre per season should be made.

Do not apply to weeds under stress, such as stress due to lack of moisture, previous herbicide injury, mechanical injury or cold temperatures, as unsatisfactory weed control could result.

Do not apply if rainfall or irrigation is expected soon after application.

Physical incompatibility, reduced weed control, or crop injury may result from mixing the tank mix of Storm plus Poast with other pesticides (fungicides, herbicides, insecticides or miticides) additives or fertilizers not recommended on the label. BASF does not recommend the use of Storm plus Poast in tank mixes other than those listed on BASF labels, supplemental labels, or technical bulletins. Local agricultural authorities may be a source of information when using other BASF recommended combinations.

Do not apply a tank mix of Storm plus Poast to soybeans that show injury, leaf phytotoxicity, and/or other prior herbicide applications because this injury may be enhanced or prolonged.

Do not apply a tank mix of Storm plus Poast through any type of irrigation system.

Do not apply to soybeans within 75 days of harvest.

Do not use treated plants for feed or forage.

In case of crop failure, only soybeans or peanuts may be immediately be replanted.

Crop rotation restriction: Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in field treated with a tank mix of Storm plus Poast for 18 months following treatment.

Do not apply more than 2 pints of Basagran or 1 pint of Blazer following an application of a tank mix of Storm plus Poast.

Do not apply more than 1.5 pints of Storm following an application of Conclude.

Do not apply sequential applications of Blazer, Basagran, or Poast Plus within 15 days following the application of a tank mix of Storm plus Poast.

After application of a tank mix of Storm plus Poast, do not apply more than 3.5 pints of Poast per acre, per season.

Do not use selective application equipment such as recirculating sprayers, wiper applicators, or shielded applicators when applying a tank mix of Storm plus Poast.

Conditions of Sale and Warranty

The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Storm and Blazer are registered trademarks of BASF Corporation.

Basagran and Poast are registered trademarks of BASF AG.

© 1993 BASF Corporation

XX ---- JJ8996 A1181093

BASF Corporation
P.O. Box 13528
Research Triangle Park, NC 27709

BASF