

Supplemental Labeling

Galaxy®

herbicide

ACCEPTED
APR 20 1994
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7969-77

Tank mix with Poast® herbicide for postemergence use in soybeans

EPA Reg. No 7969-77

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

General Information

A tank mix of Galaxy plus Poast will control broadleaf weeds and annual grasses not controlled by Galaxy or Poast alone. This tank mix is effective through contact and systemic action; therefore, weeds must be thoroughly covered with spray. Large crop- and weed-leaf canopies shelter smaller weeds and prevent adequate spray coverage. Galaxy may cause soybean leaf-speckling and leaf-bronzing under certain conditions. Soybeans are tolerant and generally outgrow these conditions in 7-10 days.

Time and Rate of Application

Apply Galaxy at 2 pints per acre plus Poast at 1 pint per acre before weeds reach the maximum size listed in Table 1. Soybeans generally should be in the first to third trifoliate stage. Early application to weeds results in the most beneficial effect on weed control and makes it easier to obtain thorough coverage. Delay in application which permits weeds to exceed the maximum size stated could result in inadequate control. Cultivation before or during applica-

tion is not recommended. Cultivation at this time may cause inadequate control because weeds may be under stress. A timely cultivation 7-10 days after application may assist in weed control.

Water Volume and Spray Pressure

Ground Equipment: Use a minimum of 10 gallons of water per broadcast acre and a maximum of 60 psi (measured at the boom, not at the pump or in the line). When crop and weed foliage is dense, use up to 20 gallons of water and a minimum of 40 psi pressure. Use standard high pressure pesticide hollow cone or flat fan nozzles spaced 20 inches apart. Do not use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles.

Aerial Application

Use a minimum of 5 gallons of water per acre and a maximum of 40 psi pressure. To obtain uniform coverage and to avoid drift hazards, the following application equipment and practices should be used:

Nozzle Type: Use only diaphragm-type nozzles producing cone or fan spray patterns.

Nozzle Height: Maximum of 10 feet above crop.

Nozzle Orientation: Nozzles must be oriented so as to discharge straight back with the air stream (opposite the direction of travel of the aircraft) or at some angle between straight back and straight down.

Nozzles must be located no farther out than 3/4 the distance from the center of the aircraft to the end of the wing or rotor. Do not apply by aircraft within 200

feet upwind of ornamental or sensitive nontarget crops such as corn, cotton, small grains, sugar beets, or sunflowers.

Applicator must follow the most restrictive use cautions to avoid drift hazard and must follow labeling as well as applicable state and local regulations and ordinances.

Spray Additive:

A nonphytotoxic oil concentrate (commonly referred to as crop oil concentrate) is required to achieve consistent weed control, especially grass control with Poast.

The oil concentrate must contain either a petroleum or vegetable oil base and must meet the following criteria:

- 1) be nonphytotoxic,
- 2) contain only EPA-exempt ingredients,
- 3) provide good mixing quality in the jar test, and
- 4) prove beneficial in local experience

With the addition of oil concentrate to Galaxy on soybeans, some leaf burn may occur, but generally all new growth is normal and crop vigor is not reduced. The potential for leaf burn is increased when relative humidity and temperature are high. A few oil concentrates have exhibited excessive leaf burn. Refer to your supplier for information concerning successful local experience prior to purchasing any oil concentrate.

Rate of Oil Concentrate Ground Application: 1.25% v/v (2 pints per acre maximum)

Table 1: Application Rate Table for Weed Growth Stages

Weeds Controlled	Maximum Height (inches)	Rate per Acre
Anoda, Spurred	3	
Beggarticks	6	
Buckwheat, Wild	3	
Canada Thistle ^{a,b}	8 to bud stage	
Cocklebur	6	
Dayflower	4	
Devilclaw ^a	3	
Galinoga ^a	2	
Jimsonweed	6	
Ladythumb	6	
Lambquarters, Common ^a	2	
Mallow, Venice	4	
Morningglories ^a	2	
Mustard, Wild	4	
Nightshade, Black	2	
Nutsedge, Yellow ^a	6-8	
Pigweed, Redroot	2	
Smooth	2	
Poinsettia, Wild	4	
Purslane, Common	1	
Ragweed, Common	3	
Giant	6	
Redweed	6	
Shepherdspurse ^a	4	
Sida, Prickly (Teaweed)	3	
Smartweed, Pennsylvania	6	
Starbur, Bristly	2	
Sunflower, Wild	5	
Velvetleaf	5	
Waterhemp, Tall	2	
Grasses Controlled		
Annual Ryegrass	4	
Barnyardgrass	4	
Broadleaf Signalgrass	4	
Foxtail, Giant	6	
Green	6	
Yellow	6	
Goosegrass	4	
Johnsongrass, Seedling	4	
Panicum, Browntop	4	
Fall	4	
Texas	4	
Millet, Wild Proso	4	
Crabgrass, Large	4	
Smooth	4	
Cupgrass, Woolly	4	
Sprangletop, Red	4	
Jungle rice	4	
Volunteer Corn	12	
Witchgrass	4	

Galaxy: 2 pints
+
Poast 1 pint
+
1.25% w/v oil concentrate

Apply 1 Application
0.625% v/v (1 pint per acre maximum)

Restrictions and Limitations
Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive label applies when using a tank mix.

Do not use treated plants for feed or forage.

Do not apply this tank mix through any type of irrigation equipment.

Do not apply this tank mix within 75 days of harvest.

Do not apply more than a total of 5 pints (80 fluid ounces) of Poast per acre per season.

Do not apply this tank mix during prolonged periods of drought or during unseasonably cold weather, as unsatisfactory weed control may result.

Do not apply to soybeans that have been subject to stress conditions such as hail damage, flooding, drought, injury from other herbicides, or widely fluctuating temperatures, as crop injury may result.


Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in a field treated with Galaxy for 18 months after treatment.

Do not apply more than 3 pints of Galaxy per acre, per season.

Rainfall immediately following application may reduce control.

BEST AVAILABLE COPY

- Do not treat earlier than leaf stage shown and do not count cotyledon leaves.
- Control may be inconsistent with the rate of Galaxy[®] herbicide. A later application of Basagran[®] herbicide may be necessary (see label for Basagran).
- Control may be partial or inconsistent with this rate of Galaxy. A later application of Blazer[®] herbicide may be necessary (see label for Blazer).
- Do not treat rosette before seed stalk appears.



3 2 7
Conditions of Sale and Warranty
The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Galaxy is a trademark and Poast is a registered trademark of BASF AG.

© 1993 BASF Corporation

XX ---- JJ9116 A1561193


BEST AVAILABLE COPY

BASF Corporation
P.O. Box 13528
Research Triangle Park, NC 27709

BASF

Supplemental Labeling

Galaxy[®]
herbicide



RT: 2-8-94
COPY 3

Tank mix with Poast Plus[®] herbicide with boron and nitrogen for postemergence use in soybeans

EPA Reg. No 7969-77

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

General Information

A tank mix of Galaxy plus Poast Plus will control broadleaf weeds and annual grasses not controlled by Galaxy or Poast Plus alone.

This tank mix is effective through contact and systemic action, therefore, weeds must be thoroughly covered with spray.

Adding a micronutrient fertilizer containing boron, which is an essential plant micronutrient, may significantly reduce leaf burn used in a tank mix with a nitrogen solution.

Time and Rate of Application

Apply Galaxy at 2 pints per acre plus Poast Plus at 1 1/2 pints per acre before weeds reach the maximum size listed in Table 1.

Soybeans generally should be in the first to third trifoliate stage. Early application provides the most beneficial effect on weed control and makes thorough coverage easier. Delay in application which permits weeds to exceed the maximum size stated could result in inadequate control.

Cultivation before or during application is not recommended as weeds may be under stress. Cultivating 7-10 days after application may assist in weed control.

Water Volume and Spray Pressure

Ground Equipment: Use a minimum of 10 gallons of water per broadcast acre and a minimum of 60 psi (measured at the boom, not at the pump or in the line). When crop and weed foliage is dense, use up to 20 gallons of water and a minimum of 40 psi pressure. Use standard high pressure pesticide hollow cone or flat fan nozzles spaced 20 inches apart. Do not use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles.

Air Application: Use a minimum of 5 gallons of water per acre and a maximum of 40 psi pressure. To obtain uniform coverage and to avoid drift hazards, use the following application equipment and practices:

Nozzle Type: Use only diaphragm-type nozzles producing cone or fan spray patterns.

Nozzle Height: Maximum of 10 feet above crop.

Nozzle Orientation: Nozzles must be oriented to discharge straight back with the air stream (opposite the direction of travel of the aircraft) or at some angle between straight back and straight down.

Nozzles must be located no farther out than 3/4 the distance from the center of the aircraft to the end of the wing or rotor. Do not apply Galaxy and Poast Plus tank mix by aircraft within 200 feet upwind of ornamental or sensitive nontarget crops. Applicator must follow the most restrictive use cautions to avoid drift hazard and must follow labeling as well as applicable state and local regulations and ordinances.

Spray Additives

A sprayable boron additive at the rate of 0.25 pounds of boron plus either 1 quart per acre of 28% UAN or 10-34-0; or 1 pound per acre of AMS should be added to the spray solution. The key factor in reducing crop injury is not to exceed this boron/nitrogen ratio. Increasing the nitrogen contribution in the additive mix will increase the chance of soybean leaf burn, thus decreasing the benefit of boron in the mixture.

Mixing

Fill the tank of a thoroughly clean sprayer half to two-thirds full with clean water. Start agitation and add Galaxy followed by Poast Plus. Allow the combination to mix thoroughly. Add boron and nitrogen and remaining volume of water. Maintain constant agitation during application.

Restrictions and Limitations

Always read and follow all label directions when using any pesticide alone or in tank mix combinations.

The most restrictive label applies when using a tank mix.

Do not use treated plants for feed or forage.

Do not apply this tank mix through any type of irrigation equipment.

Do not apply this tank mix within 75 days of harvest.

Do not apply more than a total of 7 1/2 pints (120 fluid ounces) of Poast Plus per acre per season.

Do not apply this tank mix during prolonged periods of drought or during unseasonably cold weather, as unsatisfactory weed control may result.

Do not apply to soybeans that have been subject to stress conditions such as hail damage, flooding, drought, injury from other herbi-

cides, or widely fluctuating temperatures, as crop injury may result. Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in a field treated with Galaxy for 18 months after treatment. Do not apply more than 3 pints of Galaxy per acre, per season.

A maximum of 2.5 pints per acre of Basagran and 13 fluid ounces per acre of Blazer may be applied following an initial 2.0 pint per acre application of Galaxy. Do not apply if rainfall is expected within one hour after treatment.

5 9 7
 Conditions of Sale and Warranty
 The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

Table 1: Application Rate Table for Weed Growth Stages

Weeds Controlled	Maximum Height (inches)	Rate per Acre
Beggarticks	6	Galaxy 2 pints + Post Plus 1 1/2 pints + boron* 0.25 pounds + 1 quart/acre UAN** or 1 pound AMS
Black Nightshade	< 2	
Bristly Starbur	2	
Canada Thistle*	8 to bud stage	
Cocklebur*	6	
Common Lambquarters*	2	
Common Furlane	1	
Common Ragweed	3	
Dayflower	4	
Devilclaw*	3	
Galinoga*	2	
Giant Ragweed	6	
Jimsonweed	6	
Ladythumb	6	
Morningglories*	2	
Pennsylvania Smartweed	6	
Prickly Sida (Teaweed)	3	
Redroot Pigweed	2	
Redweed	6	
Shepherdspurse*	4	
Smooth Pigweed	2	
Spurred Anoda	3	
Tall Waterhemp	2	
Velvetleaf	5	
Vernice Mallow	4	
Wild Buckwheat	3	
Wild Mustard	4	
Wild Poinsettia	4	
Wild Sunflower	5	
Yellow Nutsedge*	6-8	
Grasses Controlled		
Annual Ryegrass	4	Galaxy 2 pints + Post Plus 1 1/2 pints + boron* 0.25 pounds + 1 quart/acre UAN** or 1 pound AMS
Barnyardgrass	4	
Broadleaf Signalgrass	4	
Browntop Panicum	4	
Fall Panicum	4	
Giant Foxtail	6	
Green Foxtail	6	
Goosegrass	4	
Junglerice	4	
Large Crabgrass	4	
Red Sprangletop	4	
Seedling Johnsongrass	4	
Smooth Crabgrass	4	
Texas Panicum	4	
Volunteer Corn	12	
Wild Proso Millet	4	
Witchgrass	4	
Woolly Cupgrass	4	
Yellow Foxtail	6	

- * Do not treat earlier than leaf stage shown and do not count cotyledon leaves.
- * Control may be inconsistent with the rate of Galaxy® herbicide. A later application of Basagran® herbicide may be necessary (see label for Basagran).
- * Control may be partial or inconsistent with this rate of Galaxy. A later application of Blazer® herbicide may be necessary (see label for Blazer).
- * Do not treat rosette before seed stalk appears.
- * 0.25 pounds of boron is equivalent to 1.25 pounds of Solubor.
- ** See section titled Spray Additives for alternate nitrogen sources.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Blazer and Post Plus are registered trademarks of BASF Corporation.

Solubor is a trademark of U.S. Borax.

Basagran is a registered trademark and Galaxy is a trademark of BASF AG.

© 1994 BASF Corporation

XX ---- JJ9115 A1551193

BEST AVAILABLE COPY


BASF Corporation
 P.O. Box 13528
 Research Triangle Park, NC 27709

BASF

Supplemental Labeling

Galaxy[®]

herbicide



RT: 2-8-94
COPY 3

Tank mix with Poast Plus[®] herbicide and silicone adjuvant for postemergence use in soybeans

EPA Reg. No 7969-77

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

General Information

A tank mix of Galaxy plus Poast Plus will control broadleaf weeds and annual grasses not controlled by Galaxy or Poast Plus alone. This tank mix is effective through contact and systemic action; therefore, weeds must be thoroughly covered with spray. The use of a silicone adjuvant reduces the tendency for severe crop injury while providing good foliar spray coverage and increasing the rainfastness of the tank mix.

Time and Rate of Application

Apply this tank mix before weeds reach the maximum size listed in Table 1. Soybeans generally should be in the first to third trifoliate stage. Early application to weeds results in the most beneficial effect on weed control and makes it easier to obtain thorough coverage. Delay in application which permits weeds to exceed the maximum size stated could result in inadequate control. Cultivation before or during application is not recommended as weeds may be under stress. A timely cultivation 7-10 days after application may assist in weed control.

Water Volume Spray Pressure

Apply recommended rates of this tank mix as follows:

Ground Applications: Use a minimum of 10 gallons of water per broadcast acre at a minimum of 60 psi pressure (measured at the boom, not at the pump or in the line) to ensure adequate spray coverage. Use standard high pressure pesticide hollow cone or flat fan nozzles spaced 20 inches apart. Do not use flood or whirl chamber nozzles.

Air Applications: Use a minimum of 5 gallons of water per acre and a maximum of 40 psi pressure. To obtain uniform coverage and to avoid drift hazards, the following application equipment and practices should be used:

Nozzle Type: Use only diaphragm-type nozzles producing cone or fan spray patterns.

Nozzle Height: Maximum of 10 feet above crop.

Nozzle Orientation: Nozzles must be oriented so as to discharge straight back with the air stream (opposite the direction of travel of the aircraft) or at some angle between straight back and straight down.

Nozzles must be located no farther out than $\frac{3}{4}$ the distance from the center of the aircraft to the end of the wing or rotor. Do not apply Galaxy and Poast Plus tank mix by aircraft within 200 feet upwind of ornamental or sensitive nontarget crops. Applicator must follow the most restrictive use cautions to avoid drift hazard and must follow labeling as well as applicable state and local regulations and ordinances.

Mixing

Fill tank of a thoroughly clean sprayer half to two-thirds full with clean water. Start agitation. Add Galaxy + Poast Plus. Add silicone additive and remaining volume of water. Allow to mix thoroughly. Maintain constant agitation during application. Apply soon after mixing.

Silicone Adjuvants

The silicon adjuvant must meet the following criteria: 1) be nonphytotoxic, 2) contain only EPA-exempt ingredients, 3) provide good mixing quality, 4) be successful in local experience. Nonionic surfactants, crop oil, nitrogen (UAN or AMS) solutions or other additive combinations should not be added to tank mixes containing silicone adjuvants because excessive crop injury may result. Use 0.25% v/v (1 quart per 100 gallons of spray solution) of silicone adjuvant. The rate may need to be adjusted according to the adjuvant used.

Restrictions and Limitations

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive label applies when using a tank mix.

Do not use treated plants for feed or forage.

Do not apply this tank mix through any type of irrigation equipment.

Do not apply this tank mix within 75 days of harvest.

Do not apply more than a total of 7 $\frac{1}{2}$ pints (120 fluid ounces) of Poast Plus per acre per season.

Do not apply this tank mix during prolonged periods of drought or during unseasonably cold weather,

BEST AVAILABLE COPY

Agricultural Products

BASF

TABLE 1: Application rate table for weeds (wet weather)

Weeds Controlled	Maximum Height (inches)	Rate per Acre	
Beggarticks	6	Galaxy 2 pints + Poast Plus 1 1/2 pints + 0.25% v/v silicone adjuvant	
Black Nightshade	<2		
Bristly Starbur	2		
Canada Thistle ^a	8 to bud stage		
Cocklebur ^a	6		
Common Lambquarters ^a	2		
Common Purslane	1		
Common Ragweed	3		
Dayflower	4		
Devilsclaw ^a	3		
Gallinoger ^a	2		
Giant Ragweed	6		
Jimsonweed	6		
Ladythumb	6		
Morningglories ^a	2		
Pennsylvania Smartweed	6		
Prickly Sida (Tasweed)	3		
Redroot Pigweed	2		
Redweed	6		
Shepherdspurse ^a	4		
Smooth Pigweed	2		
Spurred Anoda	3		
Tail Waterhemp	2		
Velvetleaf	5		
Verice Mallow	4		
Wild Buckwheat	3		
Wild Mustard	4		
Wild Poinsettia	4		
Wild Sunflower	5		
Yellow Nutsedge ^a	6-8		
Grasses Controlled			
Annual Ryegrass	4		
Barnyardgrass	4		
Broadleaf Signalgrass	4		
Browntop Panicum	4		
Fall Panicum	4		
Giant Foxtail	6		
Green Foxtail	6		
Goosegrass	4		
Junglerice	4		
Large Crabgrass	4		
Red Sprangletop	4		
Seedling Johnsongrass	4		
Smooth Crabgrass	4		
Texas Panicum	4		
Volunteer Corn	12		
Wild Proso Millet	4		
Witchgrass	4		
Woolly Cupgrass	4		
Yellow Foxtail	6		

- Do not treat earlier than maximum height shown.
- Control may be inconsistent with the rate of Galaxy[®] herbicide. A later application of Basagran[®] herbicide may be necessary (see label for Basagran).
- Control may be partial or inconsistent with this rate of Galaxy. A later application of Blazer[®] herbicide may be necessary (see label for Blazer).
- Do not treat rosette before seed stalk appears.
- Rate may be adjusted according to the adjuvant used.

as unsatisfactory weed control may result.

Do not apply to soybeans that have been subject to stress conditions such as hail damage, flooding, drought, injury from other herbicides, or widely fluctuating temperatures, as crop injury may result.

Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in a field treated with Galaxy

for 18 months after treatment.

Do not apply more than 3 pints of Galaxy per acre, per season. A maximum of 2.5 pints per acre of Basagran and 13 fluid ounces per acre of Blazer may be applied following an initial 2.0 pint per acre application of Galaxy.

Do not apply if rainfall is expected within one hour after treatment.

The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Galaxy is a trademark of BASF AG.

Poast Plus is a registered trademark of BASF Corporation.

© 1994 BASF Corporation

XX --- JJ9117 A1571193

BEST AVAILABLE COPY

BASF Corporation
P.O. Box 13528
Research Triangle Park, NC 27709

BASF