

16
7969-79

Supplemental Label

Blazer + Whip 1 EC - Tank Mix Application in Soybeans
Blazer^R herbicide EPA Reg. No. 7969-79
Whip^R 1 EC herbicide EPA Reg. No. 8340-23

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA - registered label are to be followed. This labeling must be in the possession of the user at the time of herbicide application.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General Information

Blazer herbicide may be tank mixed or applied sequentially with Whip 1 EC for postemergence control of broadleaf and grass weeds. Weeds must be actively growing and at the recommended growth stages. The tank mix combination should not be used for the control of rhizome Johnsongrass.

Time of Application

Applications should be made to actively growing weeds and prior to grass tillering. Whenever the broadleaf and grass weeds are not both in the proper stage of growth according to the respective label, a sequential application should be utilized.

Sequential Application

Sequential applications should be made if all weeds to be controlled are not in the correct growth stage for treatment at the same time: when Whip is applied first, a waiting period of 3 days is necessary before applying Blazer. When Blazer is applied first, a waiting period of 7 days is necessary before applying Whip.

Rate

Use Blazer at the rate of 1.5 to 2 pints per acre, Whip 1 EC at the rate of 1.6 pints per acre for postemergence control of selected annual broadleaf/grass weeds in soybean.

Spray Additive

No surfactant or crop oil concentrate should be used when tank mixing Blazer and Whip.

REC'D
with comments
NOV 25 1981
7969-79

Page 2

Water Volume and Spray Pressure

For additional information, refer to the section entitled "Directions for Use" on the Blazer specimen label.

Ground Equipment

For the tank mix of Blazer + Whip use 20 gallons of spray solution per acre (broadcast basis) and a minimum of 40 psi pressure. Use standard high pressure hollow cone or flat fan nozzles spaced 20 inches apart. Do not use flood or whirl chamber nozzles.

Air Equipment

Use a minimum of 10 gallons of spray solution per acre.

Mixing

Fill the spray tank one-half full with water and add the recommended amount of product in the following order - Blazer, Whip-while the agitator is running, then add the remaining quantity of water.

Coverage

Thorough coverage of actively growing weeds is essential. Large crop-and-weed leaf canopies shelter smaller weeds and can prevent adequate spray coverage. Soybeans are tolerant to the tank mix; however, under certain conditions soybeans may burn, crinkle and bronze.

Restrictions and Limitations (Partial List)

Read and follow restrictions and limitations on the Blazer and Whip labels. The most restrictive labeling applies in tank mixtures.

Do not use treated plants for feed or forage.

In the case of crop failure, only soybeans or peanuts may be immediately replanted.

Crop rotation restriction: Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in fields treated with Blazer for a period of 18 months following treatment.

Do not apply the tank mix less than 90 days before harvesting.

Do not apply this tank mix through any irrigation system.

Blazer is a registered trademark of BASF Corp.

Whip 1 EC is a registered trademark of Hoechst A.G.