

7969-79

09/10/2003

1/12

**BASF**

**ACCEPTED**  
 SEP 10 2003  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 this is certified for the pesticide  
 registered under  
 EPA Reg. No. 7969-79

# ULTRABLAZER®

herbicide

**For use on peanuts, rice, and soybeans**

**Active Ingredient:**

Sodium salt of acifluorfen: Sodium 5-[2-chloro-4-(trifluoromethyl)  
 phenoxy]-2-nitrobenzoate\* .....20.1%

**Inert Ingredients:** .....79.9%

**Total** .....100.0%

\* Equivalent to 2 pounds of active ingredient per gallon.

**EPA Registration Number: 7969-79**

**EPA Est Number:**

**KEEP OUT OF REACH OF CHILDREN.**

**DANGER/PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (if you do not understand the label, find someone to explain it to you in detail.)

See inside labeling for complete **First Aid, Precautionary Statements, Directions For Use, and Conditions of Sale and Warranty.**

**Net contents: 2.5 gallons (9.462 liters)**

### FIRST AID

- If in eyes**
  - Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - Call a poison control center or doctor for treatment advice.
- If on skin or clothing**
  - Take off contaminated clothing.
  - Rinse skin immediately with plenty of water for 15 - 20 minutes.
  - Call a poison control center or doctor for treatment advice.
- If swallowed**
  - Call poison control center or doctor immediately for treatment advice.
  - Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.
  - Do not give anything by mouth to an unconscious person.
- If Inhaled**
  - Move person to fresh air.
  - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
  - Call a poison control center or doctor for further treatment advice.

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-832-HELP (4357)

### NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

**DANGER.** Corrosive. Causes irreversible eye damage. Harmful if swallowed, absorbed through skin, or inhaled. Do not get in eyes or on clothing. Avoid contact with skin and breathing vapor or spray mist.

#### Personal Protective Equipment (PPE)

##### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark, except as specified on this label for application to rice. Do not contaminate water by disposal of equipment washwaters.

Do not apply when weather conditions favor drift from target area.

#### Groundwater Advisory

Acifluorfen is present in this product. Residues of acifluorfen have been found in groundwater as a result of agricultural use. Use of this product in areas where soils are permeable, such as sand and soils with loamy sand textures, and where water tables are shallow could result in contamination of groundwater. The utilization of irrigated water in these areas will increase the likelihood of contamination.

### Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** are to be followed. This labeling must be in the user's possession during application.

#### Agricultural Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **48 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

#### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Do not store below 32° F.

**Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate violates federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### Container Disposal:

- **Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

- **Bulk/Mini-bulk Containers:** Reusable containers should be returned to the point of purchase for cleaning and refilling because the container must be thoroughly cleaned before refilling.

#### In Case of Emergency

In case of large-scale spillage regarding this product, call:

CHEMTREC 800-424-9300  
BASF Corporation 800-832-HELP

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment.
- Your local poison control center (hospital).
- BASF Corporation (800-832-HELP)

#### Steps to be taken in case material is released or spilled:

Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal. Remove contaminated clothing, and wash affected skin areas with soap and water. Wash clothing before re-use. Keep the spill out of all sewers and open bodies of water.

### I. General Information

**Ultra Blazer® herbicide** is intended for selective postemergence control of certain broadleaf weeds and grasses in peanuts, soybeans, and rice.

#### Crop Tolerance

All listed crops are tolerant to **Ultra Blazer** at all stages of growth listed. Leaf speckling may occur, but plants generally outgrow this condition within 10 days. New growth is normal and crop vigor is not reduced.

#### Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions and then triple rinsing the equipment before and after applying this product.

### II. Application Instructions

Apply recommended rates of **Ultra Blazer** as follows unless instructed differently in section **VI. Crop-Specific Information**. Applications can be made to actively growing weeds as aerial, banding, or broadcast applications at the rates and growth stages listed in **Table 1. Application Rates for Ultra Blazer® herbicide — Peanuts and Soybeans** and in **VI. Crop-Specific Information** for rice. The most effective control will result from making postemergence applications of **Ultra Blazer** early, when weeds are small. Early application to weeds results in improved weed control, allows use of the lower rate (depending on weed species), and makes thorough spray coverage easier to obtain. Delaying application permits weeds to exceed the maximum size stated and will prevent adequate control.

Avoid drift to all other crops and nontarget areas. Do not apply when conditions favor drift from target area or when windspeed is greater than 10 mph.



### Irrigation

In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth. Weeds growing under drought conditions usually are not adequately controlled.

### Spray Coverage

Weeds must be thoroughly covered with spray. Always use an adequate volume of spray solution to ensure thorough coverage. Dense leaf canopies shelter smaller weeds and can prevent adequate spray coverage.

### Cultivation

Do not cultivate within 5 days before or 7 days after applying **Ultra Blazer® herbicide**.

### Aerial Application Methods and Equipment

**Water Volume:** Use a minimum of 10 gallons of water per acre. A minimum of 5 gallons of water per acre has been effective where adequate coverage can be achieved.

**Spray Pressure:** Use up to 40 psi.

**Application Equipment:** Use only diaphragm-type nozzles that produce cone or fan spray patterns.

### Special Directions for Aerial Application

To obtain uniform coverage and to avoid drift hazards, follow these guidelines:

- Use coarse sprays (larger droplets) as they are less likely to drift.
- Do not apply **Ultra Blazer** by air if sensitive species are within 100 yards downwind or 50 yards upwind.

The applicator must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances. A drift control agent may

reduce drift, however, it may also decrease weed control.

### Ground Application (Banding)

Follow **Ground Application (Broadcast)** instructions for band applications. When row banding equipment is used, it should be adjusted to provide maximum coverage of weeds in the row. Thorough coverage of the weeds can be obtained with two nozzles directed from either side of the crop row toward the weeds in the center rows. The recommended minimum band width is 15 inches with a minimum of 15 gallons of water per acre on the band. Application with a single nozzle over the row is not recommended.

### Ground Application Methods and Equipment (Broadcast)

**Water Volume:** Use 10-20 gallons of spray solution per broadcast acre for optimal performance. Increase water volume up to 50 gallons if crop or weed foliage is dense.

**Spray Pressure:** Use a minimum of 40 psi (measured at the boom, not at the pump or in the line).

**Note:** When using the lower water volume (i.e., 10 gallons per acre) or when crop and weed foliage is dense, use a minimum of 60 psi for best results.

**Application Equipment:** Use standard high-pressure pesticide flat fan or hollow cone nozzles spaced up to 20" apart. Do not use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles as erratic coverage can cause inconsistent weed control. Do not use selective application equipment such as recirculating sprayers or wiper applicators.

**Table 1. Application Rates for Ultra Blazer® herbicide — Peanuts and Soybeans**  
**Refer to section VI. Crop-Specific Information for rate and timing details for rice.**

**Note:** Weed height will vary depending on environmental conditions and is only given as a guide. Emphasis should be placed on leaf stages. Refer to section III. **Additives** for more information.

Weeds Controlled (including triazine and ALS-resistant biotypes)	0.5 pint per acre		1.0 pint per acre		1.5 pints per acre	
	Leaf Stage <sup>a</sup> (up to)	Maximum Height	Leaf Stage <sup>a</sup> (up to)	Maximum Height	Leaf Stage <sup>a</sup> (up to)	Maximum Height
Balloonvine	—	—	—	—	2	2"
Beggarweed, Florida	—	—	—	—	2	< 2" <sup>b</sup>
Buckwheat, Wild	—	—	—	—	2	2" <sup>b</sup>
Buffalobur	—	—	—	—	2	2" <sup>b</sup>
Burgherkin	—	—	—	—	2	2" <sup>b</sup>
Carpetweed	—	—	Multi 3" dia.	< 2"	Multi. 6" dia.	2"
Citron (Wild Watermelon)	—	—	—	—	2	2" <sup>b</sup>
Cocklebur <sup>c</sup>	—	—	—	—	2	2"
Copperleaf, Hophornbeam , Virginia	—	—	2	2"	4	4"
Crotolaria, Showy	—	—	6	6" <sup>b</sup>	6	6" <sup>b</sup>
Croton, Tropic , Woolly	—	—	1-2	< 2"	2	2"
Crownbeard, Golden	—	—	1-2	< 2"	2	2"
Eclipta	—	—	—	—	2	< 2"
Galinsoga, Hairy , Smallflower	—	—	—	—	6	< 2"
Groundcherry, Cutleaf , Lanceleaf	—	—	—	—	4	< 2"
Indigo, Hairy	—	—	—	—	2	1"
Jimsonweed	—	—	—	—	2	1"
Ladysthumb	—	—	—	—	3	< 2"
Lambsquarters, Common <sup>c</sup>	—	—	4	4"	6	6"
Morningglory <sup>c</sup> , Cypressvine , Entireleaf	—	—	4	4"	6	6"
, Ivyleaf	—	—	2	2"	2	2"
, Purple Moonflower	—	—	2	2"	4	4"
, Scarlet	—	—	2	2"	4	4"
, Smallflower	—	—	2	2"	4	4"
, Small White (pitted)	—	—	2	2"	4	4"
, Tall (common)	—	—	2	2"	4	4"
, Willowleaf (Palmleaf)	—	—	2	2"	4	4"
Mustard, Wild	2	2"	4	< 4"	4	4"
Nightshade, Eastern Black , Black	—	—	2-3	< 2"	6	2"
Pigweed, Palmer	4	< 2"	2-3	< 2"	6	2"
, Prostrate	—	—	6	< 4"	6	4"
, Redroot	4	< 2"	—	—	4	4"
, Smooth	4	< 2"	6	< 4"	6	4"
, Spiny	—	—	2	< 2"	6	4"
Poinsettia, Wild	—	—	—	—	2	2" <sup>b</sup>
Poorjoe	—	—	—	—	2	2"
Purslane, Common	—	—	—	—	Multi. 6" dia.	1"
Pusley, Florida	—	—	2	2"	4	4"
Ragweed, Common , Giant	—	—	2	2"	4	3"
Senna, Coffee	—	—	2	< 2"	2	3"
Sesbania, Hemp	—	—	—	—	2	2" <sup>b</sup>
Smartweed, Pennsylvania	—	—	4	4" <sup>b</sup>	6	6" <sup>b</sup>
Smellmelon	—	—	4	4"	6	6"
Spurge, Prostrate , Spotted	—	—	—	—	2	2" <sup>b</sup>
Starbur, Bristly	—	—	—	—	Multi. 0.5" dia. Multi. 0.5" dia.	—
Waterhemp, Common , Tall	4	2"	6	< 4"	2	2" <sup>b</sup>
	4	2"	6	< 4"	6	4"
	4	2"	6	< 4"	6	4"
<b>Annual Grasses</b>						
Foxtail, Giant <sup>c</sup>	—	—	—	—	2	1"
, Green <sup>c</sup>	—	—	—	—	2	1"
, Yellow <sup>c</sup>	—	—	—	—	2	1"
Johnsongrass, Seedling <sup>c</sup>	—	—	—	—	2	1"
Panicum, Fall <sup>c</sup>	—	—	—	—	2	1"
Shattercane <sup>c</sup>	—	—	—	—	2	1"
Volunteer Small Grains <sup>c</sup>	—	—	—	—	2	1"

<sup>a</sup> Do not count leaves as pairs; count each leaf separately. Do not count cotyledon leaves. Spraying weeds in the cotyledon growth stage is not recommended.  
<sup>b</sup> Refer to **Special Use Directions**.  
<sup>c</sup> Suppression or partial control.

**Special Use Directions for Additional Weed Problems in Peanuts and Soybeans**

For the following weeds, use 1.5 pints of **Ultra Blazer® herbicide** per acre and 2 pints of spray surfactant per 100 gallons of spray mix unless otherwise stated. Activity depends on good soil moisture during and after the spray applications.

**Beggarweed, Florida**

Controlling Florida beggarweed is difficult because of the weed's long germination season. Apply **Ultra Blazer® herbicide** when beggarweed seedlings have no more than 2 young expanding true leaves. Weeds at this time will not be more than 1.5" high. It is important to obtain maximum control of the earliest weed flush. Time the cultivation to give maximum control of regrowth or secondary weed flushes. **Ultra Blazer** will suppress or partially control weeds growing under conditions of high soil moisture and high relative humidity.

**Buckwheat, Wild**

**Buffalobur**

Partial control of wild buckwheat and buffalobur can usually be obtained when the seedlings have fewer than 2 true leaves. Use **Ultra Blazer** in 30 gallons of water per acre.

**Cucurbits: Burgherkin**

**Citron (Wild Watermelon)**

**Smellmelon**

Members of the cucumber family germinate over an extended period of time. Therefore, control is difficult to obtain with a single spray. For **Ultra Blazer** to be effective, the initial application should be made to weeds no later than the 2-leaf growth stage.

**Morningglories**

More consistent control of morningglories can be achieved by using sequential applications of 1 pint of **Ultra Blazer**.

**Poinsettia, Wild**

The recommended application of **Ultra Blazer** will usually kill or severely stunt wild poinsettia. Apply before the the third true leaf has formed. This treatment will usually cause a height differential between soybeans and surviving wild poinsettia which will allow directed applications and even greater control.

**Sesbania, Hemp**

**Crotalaria, Showy**

Sesbania and crotalaria are very sensitive to **Ultra Blazer**. Apply 1 pint of **Ultra Blazer** per acre. Effective control can be obtained at just about all plant heights, however, it is important that **Ultra Blazer** be applied prior to bloom. Applications after bloom are usually not effective and therefore not

recommended. To control these weeds, the application should be timed to occur after maximum weed emergence has taken place. Care must be exercised to make certain that crop canopies do not shade this weed from spray deposits. Waiting for the sesbania to break through the crop canopy may be advisable to control late season infestations.

**Starbur, Bristly**

**Senna, Coffee**

The recommended application of **Ultra Blazer** will kill or suppress seedlings that are not past the 2-leaf stage. Applications after the 2-leaf stage are usually ineffective.

**Perennial Weeds:**

**Bindweed, Field and Hedge**

**Milkweed, Climbing and Common**

**Redvine**

**Trumpet creeper**

Growth of perennial weeds from underground rootstocks is very difficult to control. Apply **Ultra Blazer** as recommended above with 2-4 pints of spray surfactant per 100 gallons of spray mix to burn back the above-ground plant parts and retard regrowth. **Ultra Blazer** will not kill the underground rootstocks of these weeds.

**Annual Grasses:**

**Foxtail, Giant, Green, and Yellow**

**Johnsongrass, Seedling**

**Panicum, Fall**

**Shattercane**

**Ultra Blazer** must not be the basic component of a grass management program. Rather, **Ultra Blazer** can be used for additional control of escaped grasses following a preplant incorporated or pre-emergence herbicide. Grasses not exceeding the 2-leaf stage will be stunted or killed.

**Volunteer Small Grains:**


**Barley**

**Oats**

**Rye**

**Wheat**

**Ultra Blazer** applied to emerging volunteer small grains in the 1-2 leaf stage will kill or stunt many plants.



7/12

### III. Additives

To achieve consistent weed control, one of the following additives is needed: ammonium sulfate, crop oil concentrate, nonionic surfactant, or urea ammonium nitrate. AMS (or UAN) should be used when velvetleaf is a target weed. Additives may cause some leaf burn, but new growth is normal and crop vigor is not reduced. The potential for leaf burn is increased when relative humidity and temperature are high. Consult your local BASF representative for recommendations for your area. See **Table 3 Additive Rates Per Acre** for additive rates and **Table 2 Additive Options for Ultra Blazer® herbicide Tank Mixes**.

#### Ammonium Sulfate (AMS)

AMS is a dry, granular nitrogen-source fertilizer. Use only fine feed-grade or spray-grade AMS because inferior grades of AMS do not dissolve adequately and can plug spray nozzles. BASF does not recommend applying AMS if applied in less than 10 gallons per acre because of potential problems with precipitation in reduced volumes. Use AMS only if it has been demonstrated to be successful in local experience.

#### Nonionic Surfactant

The standard label recommendation is 1-2 pints of an 80% active nonionic spray surfactant per 100 gallons of water. For certain weeds, the higher spray surfactant rate is recommended.

#### Oil Concentrate

The oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:

- be nonphytotoxic,
  - contain only EPA-exempt ingredients,
  - provide good mixing quality in the compatibility test, and
  - be successful in local experience.
- The exact composition of suitable products will vary;

however, vegetable and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. For additional information, see **Compatibility Test for Mix Components**.

Some oil concentrates cause excessive leaf burn. Refer to your supplier for information concerning successful local experience before purchasing any oil concentrate.

#### Urea Ammonium Nitrate (UAN)

Commonly referred to as 28%, 30% or 32% nitrogen solution, UAN may be added in place of other spray additives to improve weed control.

Because most nitrogen solutions are mildly corrosive to galvanized, mild steel, and brass spray equipment, rinse the entire spray system with water soon after use. Do not use brass or aluminum nozzles when spraying UAN.

#### Temperature and Relative Humidity Effects

The following standard will help determine the optimum adjuvant rate to use. If the temperature and relative humidity exceed 150 (e.g., temperature of 85° F plus 70% relative humidity = 155), use the lower adjuvant rates.

**Table 3. Additive Rate Per Acre**

Additive	Ground Application	Air Application
Nonionic Surfactant	1-2 pints per 100 gallons	1-2 pints per 100 gallons
AMS Oil Concentrate UAN Solution	2.5 pounds 1-2 pints 4-8 pints	2.5 pounds 1-2 pints 4 pints

**Table 2. Additive Options for Ultra Blazer Tank Mixes**

Additive Options	Nonionic Surfactant (1-2 pints per 100 gallons)	AMS (2.5 pounds) or UAN (4-8 pints per acre)	Crop Oil Concentrate (1-2 pints per acre)	Nonionic Surfactant (1-2 pints per 100 gallons) + AMS (1-2 pounds per acre) or UAN (2-4 pints per acre)	Crop Oil Concentrate (1 pint per acre) + AMS (1-2 pounds per acre) or UAN (2-4 pints per acre)
Option A	✓				
Option B		✓			
Option C			✓		
Option D				✓	
Option E					✓

## IV. General Tank Mixing Information

### Tank Mix Partners/Components

The following products may be tank mixed with **Ultra Blazer® herbicide** according to the tank mixing instructions in this label and respective product labels.

- **Assure® II** (quizalofop)
- **Basagran®** (bentazon)
- **Cadre®** (imazamethapyr)
- **Classic®** (chlorimuron ethyl)
- **Concert® SP** (thifensulfuron methyl + chlorimuron ethyl)
- **Dual® 8E** (metolachlor)
- **Facet 75 DF®** (quinclorac)
- **FirstRate®** (chloransulam-methyl)
- **Frontier® 6.0** (dimethenamid)
- **Fusilade® DX** (fluazifop-p-butyl)
- **Fusion®** (fluazifop-p-butyl + fenoxaprop-p-ethyl)
- **Lasso® 4E** (alachlor)
- **Matador®** (quizalofop)
- **Pinnacle®** (thifensulfuron methyl)
- **Poast®** (sethoxydim)
- **Poast® HC** (sethoxydim)
- **Poast Plus®** (sethoxydim)
- **Propanil**
- **Pursuit®** (imazethapyr)
- **Raptor®** (imazamox)
- **Reliance® STS**  
(thifensulfuron methyl + chlorimuron ethyl)
- **Resource®** (flumiclorac)
- **Roundup® Ultra** (glyphosate)
- **Scepter®** (imazaquin)
- **Select®** (clethodim)
- **Skirmish®** (chlorimuron ethyl)
- **Synchrony® STS™**  
(thifensulfuron methyl + chlorimuron ethyl)
- **2,4-DB**
- **2,4-DB LVE** (preplant burndown only)

See section **VI. Crop-Specific Information** for more details. Read and follow the applicable **Restrictions and Limitations** and **Directions For Use** on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Ultra Blazer** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. BASF does not recommend using tank mixes other than those listed on BASF labeling. Local agricultural authorities may be a source of information when using other than BASF recommended tank mixes.

### Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test.

For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Only use water from the intended source at the source temperature.

Add components in the sequence indicated in the **Mixing Order** using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

### Mixing Order

- 1) **Water.** Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
- 2) **Agitation.** Maintain constant agitation throughout mixing and application.
- 3) **Products in PVA bags.** Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
- 4) **Water-dispersible products** (such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions). If an inductor is used, rinse it thoroughly after the component has been added.
- 5) **Water-soluble products** (such as **Ultra Blazer**). If an inductor is used, rinse it thoroughly after the component has been added.
- 6) **Emulsifiable concentrates** (such as oil concentrate when applicable). If an inductor is used, rinse it thoroughly after the component has been added.
- 7) **Water-soluble additives** (such as AMS or UAN when applicable). If an inductor is used, rinse it thoroughly after the component has been added.
- 8) **Remaining quantity of water.** Maintain constant agitation during application.



## V. Restrictions and Limitations

- **Maximum seasonal use rate:** Do not apply more than a total of **2 pints** (0.5 pound of active ingredient) of **Ultra Blazer® herbicide** per acre, per season in soybeans and peanuts. Do not apply more than a total of **1 pint** (0.25 pound active ingredient) of **Ultra Blazer** per acre, per season in rice.
- **Maximum application use rate:** Do not apply more than **1.5 pints** (0.375 pound of active ingredient) of **Ultra Blazer** per acre, per application in peanuts and soybeans. Do not apply more than **1 pint** (0.25 pound of active ingredient) of **Ultra Blazer** per acre, per application in rice.
- **Preharvest Interval (PHI):** See **Table 4**.
- **Restricted Entry Interval (REI): 48 hours**
- Allow a minimum of **15 days** between **sequential applications** of **Ultra Blazer**.
- Do not use treated plants for feed or forage.
- In case of crop failure, only peanuts, soybeans, or rice may be immediately replanted.
- **Crop Rotation Restriction:** Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in fields treated with **Ultra Blazer** for **18 months** following treatment.
- **Stress:** Do not apply to weeds or crops under stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control may result.
- Do not apply **Ultra Blazer** to crops that show **injury** (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced or prolonged.
- **Rainfast period:** Rainfall or overhead irrigation within 4 hours after application may reduce the effectiveness of **Ultra Blazer**.
- Do not apply through any type of **irrigation** equipment.

**Table 4. Crop-Specific Restrictions and Limitations**

Crop	Minimum Time from Application to Harvest (PHI)	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application
Peanuts	75 days	1.5 pints	2 pints	No	Yes
Rice	50 days	1 pint	1 pint	No	Yes
Soybeans	50 days	1.5 pints	2 pints	No	Yes

## VI. Crop-Specific Information

### Peanuts

Apply the rates of **Ultra Blazer® herbicide** recommended in **Table 1** to peanuts pre-emergence, at cracking stage (initiation of soil cracking, but before peanut emergence from the soil), or postemergence to peanuts to control susceptible weeds.

### Peanut Tank Mixes

**Ultra Blazer** may be applied in a tank mix with one the following herbicides:

Tank Mix Partner	Additive Option
• <b>Basagran®</b> .....	A or C
• <b>Cadre®</b> .....	A
• <b>Dual® 8E</b> .....	A
• <b>Frontier® 6.0</b> .....	A
• <b>Lasso® 4E</b> .....	A
• <b>Poast®</b> .....	C
• <b>Poast® HC</b> .....	C
• <b>Poast Plus®</b> .....	C
• <b>2,4-DB*</b> .....	A or C

\* Do not apply this tank mix after pod-filling stage begins.

Refer to **Table 2** for the additive option appropriate for each tank mix.

### Rice

**Ultra Blazer** may be applied when rice is at the late tillering stage up to the early boot stage, which normally occurs in June or July. Rice must be past the 3-leaf stage. Apply **Ultra Blazer** to hemp sesbania plants before sesbania is in the flowering stage. Best results are obtained when the sesbania growth extends above the rice.

Apply 0.5 pint of **Ultra Blazer** per acre to hemp sesbania plants. A second application of 0.5 pint of **Ultra Blazer** per acre can be made to control later germinating sesbania. To achieve consistent weed control, add 1-2 pints of an 80% active nonionic spray surfactant per 100 gallons of water. Using a spray adjuvant is important for effective control of hemp sesbania.

### Specific Restrictions and Limitations

Do not apply **Ultra Blazer** after the rice reaches the boot stage.

The maximum application rate for rice is 1 pint per acre, per season and should only be used to control hemp sesbania.

Do not apply more than 2 applications to rice per season nor exceed 1 pint per acre per season.

Do not use water from treated rice fields for irrigation purposes for other than those labeled for use with **Ultra Blazer® herbicide**.

Do not harvest crayfish from treated rice areas for food.

### Rice Tank Mixes

**Ultra Blazer** may be applied in a tank mix with one the following herbicides:

Tank Mix Partner	Additive Option
• <b>Basagran®</b> .....	A
• <b>Facet® 75 DF</b> .....	A
• <b>Propanil</b> .....	A

Refer to **Table 2** for the additive option appropriate for each tank mix.

### Soybeans

To ensure optimum spray coverage of weeds, apply **Ultra Blazer® herbicide** to small actively growing weeds. Refer to section II. **Application Instructions** and **Table 1** for more information. A sequential application of 1 pint of **Ultra Blazer** following 1 pint of **Ultra Blazer** can be used to control subsequent weed flushes or escaped weeds before they reach the maximum weed size listed in **Table 1**.

### Soybean Tank Mixes

**Ultra Blazer** may be applied in a tank mix with one the following herbicides:

Tank Mix Partner	Additive Option
• <b>Assure® II</b> ¹ .....	A
• <b>Basagran®</b> .....	A or C
• <b>Classic®</b> .....	A
• <b>Concert® SP</b> (up to 0.25 ounce) .....	D
• <b>FirstRate®</b> .....	D
• <b>Frontier® 6.0</b> .....	A
• <b>Fusilade® DX</b> .....	A
• <b>Fusion®¹</b> .....	A
• <b>Matador®¹</b> .....	A
• <b>Pinnacle®</b> (up to 0.25 ounce) .....	A or D
• <b>Poast®¹</b> .....	C
• <b>Poast® HC</b> ¹ .....	C
• <b>Poast Plus®¹</b> .....	C
• <b>Pursuit®</b> .....	D
• <b>Raptor®</b> .....	D
• <b>Reliance® STS SP</b> ² (up to 0.25 ounce) .....	D
• <b>Resource®</b> .....	C
• <b>Roundup® Ultra</b> .....	8.5-17 pounds of AMS per 100 gallons
• <b>Scepter®</b> .....	A
• <b>Select® 2 EC</b> .....	C
• <b>Skirmish®</b> .....	D
• <b>Synchrony® STS</b> ² (up to 0.5 ounce) .....	E
• <b>2,4-DB</b> .....	A

¹ For best results if applying as part of a weed control program with **Ultra Blazer**, follow these guidelines:

- If the partner is applied prior to the **Ultra Blazer** application, wait 24 hours before applying **Ultra Blazer**.
- If the partner is applied following the **Ultra Blazer** application, wait 7 days before applying.

² When applying this tank mix to soybean varieties other than those designated as **STS**, do not add oil concentrate. Application to soybean varieties not designated as **STS** will result in severe crop injury or yield loss.

Refer to **Table 2** for the additive option appropriate for each tank mix.

### Burndown Treatment Before Planting Soybeans

**Ultra Blazer** alone can be applied any time before planting soybeans to control susceptible weed species present (see **Table 1**). This application is not intended to replace a full-season weed control program, but is intended to control susceptible weed species present before soybeans are planted. Use a spray additive to enhance burndown activity before planting soybeans.

### Burndown Tank Mixes

**Ultra Blazer** may be applied in a tank mix with one the following herbicides:

Tank Mix Partner	Additive Option
• <b>Poast®</b> .....	C or E
• <b>Poast® HC</b> .....	C or E
• <b>Poast Plus®</b> .....	C or E
• <b>2,4-D LVE</b> .....	C

Refer to **Table 2** for the additive option appropriate for each tank mix.

Weeds listed in this label: Broadleaves	
Common Name	Scientific Name
Artichoke, Jerusalem	Helianthus tuberosus
Balloonvine	Cardiospermum halicacaburn
Beggarweed, Florida	Desmodium tortuosum
Beggarticks	Bidens frondosa
Bindweed, Field	Convolvulus arvensis
, Hedge	Convolvulus sepium
Buckwheat, Wild	Polygonum convolvulus
Buffalobur	Solanum rostratum
Burgherkin	Cucumis anguria
Carpetweed	Mollugo verticillata
Citron (Wild Watermelon)	Citrullus vulgaris
Cocklebur, Common	Xanthium pensylvanicum
, Heartleaf	Xanthium strumarium
Copperleaf, Hophornbeam	Acalypha ostryaefolia
, Virginia	Acalypha virginica
Crotalaria, Showy	Crotalaria spectabilis
Croton, Tropic	Croton glandulosus
	Croton capitatus
Crownbeard, Golden	Verbesina encelioides
Cucumber, Wild Spiny	Cucumis dipsaceus
Eclipta	Eclipta alba
Galinsoga, Hairy	Galinsoga ciliata
, Smallflower	Galinsoga parviflora
Groundcherry, Cutleaf	Physalis angulata
, Lanceleaf	Physalis lanceifolia
Indigo, Hairy	Indigo fera hirsuta
Jimsonweed	Datura stramonium
Ladysthumb	Polygonum persicaria
Lambsquarters, Common	Chenopodium album
Milkweed, Climbing	Sarcostemma cyanchoides
, Common	Asclepias syriaca
Morningglory, Cypressvine	Ipomoea quamoclit
, Entireleaf	Ipomoea hederacea
	var. integruscula
, Ivyleaf	Ipomoea hederacea
	var. hederacea
, Purple Moonflower	Ipomoea muricata
, Scarlet	Ipomoea coccinea
, Smallflower	Jacquemontia tamnifolia
, Small White (Pitted)	Opomoea lacunosa
, Tall (Common)	Ipomoea purpurea
, Willowleaf (Palmleaf)	Ipomoea wrightii
Mustard, Wild	Brassica kaber
Nightshade, Eastern Black	Solanum ptycanthum
, Black	Solanum nigrum
Pigweed, Palmer	Amaranthus palmeri
, Prostrate	Amaranthus blitoides
, Redroot	Amaranthus retroflexus
, Smooth	Amaranthus hybridus
, Spiny	Amaranthus spinosus
Poinsettia, Wild	Euphorbia heterophylla
Poorjoe	Diodia teres
Purslane, Common	Protulaca oleracea
Pusley, Florida	Richardia scabra
Ragweed, Common	Ambrosia artemisifolia
, Giant	Ambrosia trifida
Redvine	Brunnichia cirrhosa
Senna, Coffee	Cassia occidentalis
Sesbania, Hemp	Sesbania exaltata
Smartweed, Pennsylvania	Polygonum pensylvanicum
Smellmelon	Cucumis melo
Spurge, Prostrate	Euphorbia supina
, Spotted	Euphorbia maculata
Starbur, Bristly	Acanthospermum hispidum
Teaweed (See Sida, Prickly)	Sida spinosa
Trumpetcreeper	Campsis radicans
Velvetleaf	Abutilon theophrastic
Waterhemp, Common	Amaranthus rudis
, Tall	Amaranthus tuberculatus

Weeds listed in this label: Grasses	
Common Name	Scientific Name
Foxtail, Giant	Setaria faberi
, Green	Setaria viridis
, Yellow	Setaria lutescens
Johnsongrass, Seedling	Sorghum halepense
, Rhizome	Sorghum halepense
Panicum, Fall	Panicum dichotomiflorum
, Texas	Panicum texanum
Shattercane	Sorghum bicolor
Volunteer Barley	Hordeum vulgare
Corn	Zea mays
Oats	Avena sativa
Rye	Secale cereale
Wheat	Triticum aestivum

12/12

**Crops**

This product can be used on the following crops:

**Peanut  
Rice  
Soybeans**

Look inside for complete **Restrictions and Limitations** and **Application Instructions**.

**Conditions of Sale and Warranty**

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer. BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. **BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.** BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

*Basagran, Blazer, Cadre, Facet, Frontier, Poast, Poast Plus, Pursuit, Raptor and Scepter are registered trademarks of BASF.*

*Assure, Classic, Concert, Pinnacle, Reliance, and Synchrony are registered trademarks and STS is a trademark of E.I. DuPont de Nemours and Company.*

*Dual is a registered trademark of Novartis.*

*FirstRate is a registered trademark of DowElanco.*

*Fusilade is a registered trademark of Zeneca, Limited.*

*Fusion is a registered trademark of Zeneca, Inc.*

*Lasso, Roundup, and Roundup Ready are registered trademarks of Monsanto Company.*

*Matador and Skirmish are registered trademarks of FMC Corporation.*

*Resource and Select are registered trademarks of Valent USA Corporation.*

© 2003 BASF Corporation  
All rights reserved  
007969-03106.20030626  
NVA 2003-04-098-106  
Supersedes NVA 99-4-98-0142  
EPA Approved

BASF Corporation  
Agricultural Products  
26 Davis Drive  
Research Triangle Park, NC 27709

