

Ms. Karen R. Blundell
BASF Corporation
Agricultural Products
P.O. Box 13528
Research Triangle Park, NC 27709-3528

Dear Ms. Blundell:

SUBJECT: Label Amendment Adding Ground Water Advisory and
WPS Label Language
BAS 530 04H Herbicide
EPA Reg. No.: 7969-82
Your Application Dated May 21, 1997, as Amended by FAX on July 18, 1997

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, provided that you:

- 1. Revise the second paragraph under "HAZARDS TO HUMANS AND DOMESTIC ANIMALS" to read as follows:

"Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals."

- 2. Revise the Statement Of Practical Treatment to read as follows:

"If In Eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

If Swallowed: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, eggwhites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

If On Skin: Wash with plenty of soap and water. Get medical attention."

RD:STANTON:PM Team 23:Rm. 237:CM-2:305-5218:Disk # 6:5524542. LET

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	S. Stanton							
DATE ▶	Jul 18, 1997							

3. Add the following Note To Physician in close proximity to but clearly distinguished from the Statement of Practical Treatment:

"Probable mucosal damage may contraindicate the use of gastric lavage."

4. Delete the words "puncture and" from the glass container disposal instructions on page 2.
5. Add the following statement under "Restrictions and Limitations" on page 5 (as on the previously accepted label):

"Make applications before soybeans bloom."

A stamped copy is enclosed for your records. Submit one copy of the final printed labeling incorporating these changes before you release the product for shipment.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

3/8

RT Date: 5-27-97

ACCEPTED
with COMMENTS
In EPA Letter Dated

JUL 18 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
7969-82

BASF

BAS 530 04H

herbicide

Postemergence Herbicide

For broad spectrum weed control in soybeans.

A soluble liquid formulation containing:

Active Ingredients: *	
Sodium salt of bentazon:	29.0%
Sodium salt of fomesafen	12.9%
Inert ingredients	58.1%
Total	100.0%

*Equivalent to 2.66 pounds per gallon bentazon (3-(1-methylethyl)-1H-2,1,3-benzothiadiazin-4(3H)-one 2,2 dioxide and 1.21 pounds per gallon of fomesafen (5-(2-chloro-4-trifluoromethyl) phenoxy)-N-methylsulfonyl)-2-nitrobenzamide.

EPA Reg. No. 7969-82

KEEP OUT OF REACH OF CHILDREN.

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to read it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the **Directions For Use** section for information about this standard.

Net contents 2 1/2 gallons

BASF Corporation
P.O. Box 13528, Research Triangle Park, NC, 27709-3528

Specimen Label

4/8

**Precautionary Statements
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS
DANGER**

This product contains fomesafen which has been determined to cause tumors in laboratory animals (mice). Risks can be reduced by closely following use directions and precautions and by wearing the protective clothing specified elsewhere on this label.

This product causes irreversible eye damage. Do not get in eyes, on skin or on clothing. Avoid breathing vapors or spray mist.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in WPS.

If swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician or Poison Control Center.

Environmental Hazards

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. Do not apply when weather conditions favor drift from target area. This product contains fomesafen sodium which is known to leach through soil into ground water under certain conditions as a result of label use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Directions for use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry to treated areas during the restricted entry interval REI of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, and water, includes:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Storage and disposal

Do not allow product to freeze. Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide spray mixture, is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA regional office for guidance.

Container Disposal

Metal Containers: Triple-rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Plastic Containers: Triple-rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Glass Containers: Triple-rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other state and local procedures.

Mini Bulk Container Precautions-

Before refilling, inspect thoroughly for damage, such as cracks, puncture, bulges, dents, abrasions and damaged or worn thread on closure devices. **REFILL ONLY WITH BAS 530 04 H.** The contents of this container cannot be completely removed by cleaning. Refilling with materials other than BAS 530 04 H will result in contamination and may weaken container. After filling and before transporting, check for leaks. Do not refill or transport damaged or leaking container. **CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

In Case of Emergency

In case of large-scale spillage, regarding this product, call: CHEMTREC - 800-424-9300 BASF Corporation - 8(10-832-HELP

In case of medical emergency regarding this product, call:

1. Your local doctor for immediate treatment.
2. Your local poison control center (hospital).
3. BASF Corporation - 800-832-HELP.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Statement of Practical Treatment

If in eyes: Flush eyes immediately with large amounts of water for at least 15 minutes. Get medical attention.

If on skin: Immediately remove contaminated clothing and wash thoroughly with plenty of soap and water. Get medical attention if irritation persists.

Steps to be taken in case material is released or spilled: Dike and contain spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal. Remove contaminated clothing and wash affected skin areas with water. Wash clothing before re-use. Keep spill out of all sewers and open bodies of water.

General Information

BAS 530 04 H herbicide is intended for selective postemergence control of certain broadleaf weeds in soybeans. **BAS 530 04 H** is effective mainly through contact action; therefore, weeds must be thoroughly covered with spray. Soybeans are tolerant to **BAS 530 04 H** when applied at recommended rates. **BAS 530 04 H** may cause some soybean leaf-speckling and leaf-bronzing to occur under certain conditions; however, soybeans soon outgrow these effects with no subsequent adverse effects noted (See **Restrictions and Limitations**).

BAS 530 04 H is intended for use in the following states or parts of states:

Alabama, Arkansas, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma (east of U.S. Highway 75 and east of Indian Nation Parkway), South Carolina, Tennessee, and Texas (counties east of U.S. Highway 75 and Interstate 45, additional counties of Austin, Brazoria, Chambers, Colorado, Fort Bend, Harris, Matagorda, Waller, and Wharton).

Indiana, Illinois, Maine, Massachusetts, Missouri, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont and West Virginia.

Iowa, Kansas (all counties east of U.S. Highway 75), Michigan (all counties south of U.S. Highway 46), Minnesota (all counties south of Highway 212), Nebraska (all counties east of U.S. Highway 81), and Wisconsin (all counties south of Interstate 94).
Delaware, Kentucky, Maryland, and Virginia.

Timing of applications

Make postemergence applications of **BAS 530 04 H** early, when weeds are small and actively growing and before weeds reach the maximum size listed in the **Application Rate Table**. Delay in application which

permits weeds to exceed the maximum size stated will result in inadequate control.

Cultivation before or during the application is not recommended. Cultivation may put weeds under stress thus making control more difficult to obtain. Timely cultivation 5-7 days after application will usually assist in weed control.

Water Volume and Spray Pressure

Apply recommended rates of **BAS 530 04 H** as follows:

Ground equipment: Use sufficient spray volume and pressure to ensure thorough weed coverage. Use 10-20 gallons of water per broadcast acre and a minimum of 40 psi pressure (measured at the boom, not at the pump or in the line). When crop and weed foliage is dense, use up to 50 gallons of water and up to 80 psi pressure. Use standard high pressure pesticide hollow cone or flat fan nozzles spaced 20 inches apart. Do not use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles.

Air equipment: Use sufficient spray volume and pressure to ensure thorough weed coverage. Use 5-10 gallons of water per acre and a maximum of 40 psi pressure. Use only diaphragm-type nozzles producing cone or fan spray patterns.

Aerial Application - special directions and restrictions

To obtain uniform coverage and to avoid drift hazards, the following application equipment and practices should be used:

Nozzle height: 6 to 10 feet above crop.

Nozzle orientation: Nozzles must be oriented so as to discharge straight back with the air stream (opposite the direction of travel of the aircraft) and not more than 20 degrees downward.

Nozzles must not be located further out than $\frac{3}{4}$ the distance from the center of the aircraft to the end of the wing or rotor.

Do not apply **BAS 530 04 H** by aircraft when the wind is blowing at a velocity above 10 mph. Coarse sprays (larger droplets) are less likely to drift.

Do not apply **BAS 530 04 H** by air if ornamentals or sensitive non-target crops, such as cotton, sugar beets, sunflowers or okra are within 200 feet downwind.

5/8
Applicator must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

Addition of Oil Concentrate

A nonphytotoxic crop oil concentrate (commonly referred to as oil concentrate) should be added to the spray tank with **BAS 530 04 H** to achieve consistent weed control.

The oil concentrate must contain either a petroleum or vegetable oil base and must meet the following criteria: 1) be nonphytotoxic, 2) contain only EPA-exempt ingredients, 3) provide good mixing quality in the jar test, and 4) prove beneficial in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers which provide good mixing quality. For vegetable oil concentrates, it has been observed that highly refined vegetable oils are more satisfactory than unrefined vegetable oils. For additional information, see **Jar test for estimating suitability of oil concentrates** at the end of this section.

With the addition of oil concentrate to **BAS 530 04 H** on soybeans, some leaf burn may occur, but generally all new growth is normal and crop vigor is not reduced. The potential for leaf burn is increased when relative humidity and temperature are high. A few oil concentrates have exhibited excessive leaf burn. Refer to your supplier for information concerning successful local experience prior to purchasing any oil concentrate.

Rate of oil concentrate:

Ground application - 1 pint/acre maximum

Air application - 1 pint/acre maximum

Jar test for estimating suitability of oil concentrates

- Water supply:** Use only water from intended source and at the source temperature.
- Amount of water in jar:** Ground application - For 20 gals./A spray volume use $3\frac{1}{3}$ cups (800 ml) of water. Air application - For 10 gals./A spray volume use $1\frac{2}{3}$ cups (400 ml) of water. For other spray volumes, adjust proportionately to above.

BEST AVAILABLE COPY

9/8

APPLICATION RATE TABLE

Applications of **BAS 530 04H** should be made when weeds are small and actively growing and before weeds reach the maximum size listed in the **Application Rate Table**. Such applications generally correspond to the soybean growth stages of unifoliate to two expanded trifoliate leaves. Soybeans may experience slight yellowing, bronzing, speckling, or burning of leaves under certain conditions. Soybean plants generally outgrow this condition within 10 days.

Weeds Controlled*	Weed Growth Stages		BAS 530 04H Rate Per Acre	Oil Concentrate Rate
	Leaf Stage Up to	Maximum Height		
Bristly Starbur	2-6	3"	1.5 pints/A	1.0 pint/A
Black Nightshade	2-4	3"		
Carpetweed	6" diameter	2"		
Cocklebur*	2-4	6"		
Common Lambsquarters ^b	2-4	2"		
Common Purslane	Multi-leaf	6" diameter		
Common Ragweed	4-6	3"		
Giant Ragweed	Up to 4	6"		
Hemp Sesbania	4	6"		
Jimsonweed	Up to 4	4"		
Ladysthumb	Up to 6	6"		
Morningglory, Cypressvine	4	4"		
Entireleaf	2	2"		
Ivyleaf	2	2"		
Purple Moonflower	2	2"		
Small White (pitted)	2	2"		
Tall (Common)	2	2"		
Pennsylvania Smartweed	Up to 6	6"		
Prickly Sida or Teaweed ^b	Up to 4	2"		
Redroot Pigweed	Up to 6	3"		
Redweed	2-4	3"		
Showy Crotalaria	4	2"		
Smooth Pigweed	Up to 6	3"		
Spurred Anoda	4	2"		
Tropic Croton	2	1"		
Velvetleaf	Up to 4	2"		
Venice Mallow	Up to 6	2"		
Wild Mustard	Up to 6	4"		
Wild Sunflower	2	3"		
Wild Poinsetia	4	4"		

a Do not treat earlier than leaf stage shown and do not count cotyledon leaves.

b Control may be inconsistent with this rate of **BAS 530 04H**. A later application of **Basagran** may be necessary (see label for **Basagran**).

* See Appendix for scientific names.

- 3. Amount of BAS 530 04 H and oil concentrate to add: Add BAS 530 04 H and oil concentrate at the rate of 1 teaspoon (5 ml) for each pint of recommended label rate.
- 4. Add components in following sequence, gently mixing between component additions:
 - a. BAS 530 04 H
 - b. Oil concentrate
- 5. Cap jar, invert 10 cycles, let stand for 15 minutes, evaluate.
- 6. Evaluation: An ideal tank mix combination will be uniform; thus, the suitability of the oil concentrate is questionable if any of the following are observed:
 - Free oil (film or globules) at the surface.
 - Flocculation - fine particles which may be suspended in the liquid or found as a precipitated layer at the bottom of the jar.
 - Clabbering - thickening texture (coagulated) resembling yogurt or a curd-like texture as with cottage cheese.

Mixing/spraying
 Fill tank of a thoroughly clean sprayer half to two-thirds full with clean water. Start agitation and add BAS 530 04 H; allow to mix thoroughly. Add spray additive and remaining volume of water. Maintain constant agitation during application.

Restrictions and Limitations:
 Do not apply more than a total of 1 1/2 pints of BAS 530 04 H per acre per season.
 Do not apply BAS 530 04 H to soybeans that have been subject to stress conditions such as hail damage, flooding, drought, injury from other herbicides, or widely fluctuating temperatures, as crop injury may result.
 Do not apply BAS 530 04 H to soybeans that show injury (leaf phytotoxicity and/or plant stunting) produced by other prior herbicide applications, because this injury may be enhanced and/or prolonged.
 Do not apply BAS 530 04 H during prolonged periods of drought or during unseasonably cold weather, as unsatisfactory weed control may result.
 Rainfall or overhead irrigation soon after application may nullify the effectiveness of BAS 530 04 H.
 Physical incompatibility, reduced weed control, or crop injury may result from mixing BAS 530 04 H with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. BASF does not recommend the use of BAS 530 04 H tank mixes other than those listed on BASF labels, supplemental labels, or technical information bulletins. Local agricultural authorities may be a source of information when using other than BASF approved tank mixes.
 Clean sprayer thoroughly prior to application of BAS 530 04 H, particularly if a herbicide was used which has the potential to injure the crop to be sprayed.

Do not graze treated areas or harvest for forage or hay.
 In the case of crop failure, only soybeans may be immediately replanted.
 Avoid overlapping spray swaths.
 Otherwise, injury may occur to rotational crops.
 Avoid drift to all other crops and nontarget areas.

Do not apply BAS 530 04 H through any type of irrigation system.
 Do not apply Reflex® 2LC herbicide during the same year after application of BAS 530 04 H.
 Do not apply more than 3.0 pints of Basagran® herbicide following an application of 1.5 pints of BAS 530 04 H per acre per season.

Do not apply BAS 530 04 H to any field more than once every two years in the following states or parts of states:
 Indiana, Illinois, Maine, Massachusetts, Missouri, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, West Virginia, Iowa, Kansas (all counties east of U.S. Highway 75), Michigan (all counties south of U.S. 40), Minnesota (all counties south of highway 212), Nebraska (all counties east of U.S. Highway 81), Wisconsin (all counties south of Interstate 94), Delaware, Kentucky, Maryland and Virginia.

Rotational Crop Restrictions
 The following rotational crops may be planted after applying BAS 530 04H at recommended rates in soybeans:

Crop to be Planted	Minimum Rotation Interval (Months After Last BAS 530 04H Application)
Small grains, such as wheat, barley, rye	4
Corn, cotton, peanuts, rice	10
To avoid crop injury do not plant sunflowers, sugar beets, sorghum or any other crop not listed above within	18

Do not graze rotated small grain crops or harvest for livestock forage or straw.

BEST AVAILABLE COPY

APPENDIX

The following are scientific names for the weeds listed in this section. For specific recommendations on the control of these weeds, refer to the **Application Rate Table**.

Common Name	Scientific Name
Bristly Starbur	<i>Acanthospermum hispidum</i>
Black nightshade	<i>Solanum nigrum</i>
Carpetweed	<i>Mollugo verticillata</i>
Cocklebur	<i>Xanthium strumarium</i>
Common Lambsquarter	<i>Chenopodium album</i>
Common Purslane	<i>Portulaca oleracea</i>
Common Ragweed	<i>Ambrosia artemisiifolia</i>
Giant Ragweed	<i>Ambrosia trifida</i>
Hemp Sesbania	<i>Sesbania exalta</i>
Jimsonweed	<i>Datura stramonium</i>
Ladysthumb	<i>Polygonum persicaria</i>
Morningglory, Common	<i>Ipomoea purpurea</i>
, Cypressvine	<i>Ipomoea quamoclit</i>
, Ivyleaf	<i>Ipomoea hederacea</i>
, Entireleaf	<i>Ipomoea hederacea</i>
, Pitted	<i>Ipomoea lacunosa</i>
, Purple Moonflower	<i>Ipomoea muricata</i>
Pennsylvania Smartweed	<i>Polygonum pennsylvanicum</i>
Pigweed, Redroot	<i>Amaranthus retroflexus</i>
, Smooth	<i>Amaranthus hybridus</i>
Prickly Sida or Teaweed	<i>Sida spinosa</i>
Redweed	<i>Melochia corchorifolia</i>
Showy Crotalaria	<i>Crotalaria breviflora</i>
Spurred Anoda	<i>Anoda cristata</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Venice Mallow	<i>Hibiscus trionum</i>
Wild Mustard	<i>Sinapsis arvensis</i>
Wild Sunflower	<i>Helianthus annuus</i>
Wild Poinsettia	<i>Euphorbia heterophylla</i>

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Reflex is a registered trademark of ICI Group Companies.

BASF Corporation
 P.O. Box 13528
 Research Triangle Park, NC 27709

BASF