79	عا	9	_	8	2
			1		~

# 12-15-2-003

	reverse before comple	ting form.	-	Form Approve	d. OMB No. 2070	0-0060. Approval expires 2-28-95	
United States Environmental Protection Washington, DC 204			ion Agency		Registration Amendment Other	n OPP Identifier Number	
		Application	n for Pestic	de - Section	ı <b>1</b>		
1. Company/Product Number 7969-85			2. EPA Product Manager Ms. M. Waller		T T	3. Proposed Classification  None Restricted	
4. Company/Product (Name) Curalan EG			PM# 21	PM#			
5. Name and Address of Applicant (Include ZIP Code) BASF Corporation, Agricultural Products PO Box 13528 Research Triangle Park, NC, 27709-3528  Check if this is a new address			6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to:  EPA Reg. No.  Product Name				
			Section -				
Amendment - Explain Resubmission in resp  Notification - Explain  Explanation: Use addition  Minor correction to final print	below.		I and Section II.)	Final printed lab Agency letter da "Me Too" Appli Other - Explain b	cation.	NOTIFICATION DEC 15 2003	
Material This Product Wil	l Be Packaged In:		Section - I				
Child-Resistant Packaging  Yes  No  * Certification must	Yes Yes No		Weter Soluble if Yes No If "Yes" Package wgt	No Place Oss* No. per  Place Place Pap		etal astic ass	
Container 4 X 1 p  6. Manner in Which Label is Affixed to Product Lithography		Lithogra	pound or 4 X 2.75 pounds  Other				
	·	Paper of Stencile	Section - I	v			
1. Contact Point (Complete	items directly below fi	or identification			cessary, to process	s this eoficetion l	
Name T.R. Neisen		T	Title T Manager, Regulatory Affairs (		Tele	phone No.º (Indivde Area Code) 9)547-2914	
	ments I have made on y knowlingliy false or law.		ll attachments th				
2. Signature Welsen		,	. Title Manager, Regulatory Affairs				
4. Typed Name T.R. Nelsen		5.	5. Date 12/3/2003			•• ••	



 Active Ingredient:
 Vinclozolin [3-(3, 5-dichlorophenyl)-5-ethenyl-5-methyl-2, 4-oxazolidinedione]
 .50.0%

 Inert Ingredients:
 .50.0%

 Total
 .100.0%

 EPA Reg. No. 7969-85
 EPA Establishment No.

Keep out of reach of children.

# CAUTION

See inside for complete Precautionary Statements, Directions For Use and Conditions of Sale and Warranty.

### STATEMENT OF PRACTICAL TREATMENT

If on skin: Wash with plenty of soap and water. Get medical attention.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to

mouth. Get medical attention.

If in eyes: Flush with plenty of water. Call a physician if irritation persists.

See additional Precautionary Statements inside booklet.

### **Net contents:**

Product of Germany; Formulated in the United States with U.S. and imported ingredients









### **Precautionary Statements**

Hazards to Humans and Domestic Animals Caution. Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Do not get on skin, in eyes, or on clothing. Avoid breathing vapor or spray mist. Prolonged exposure or frequently repeated skin contact may cause allergic reactions in some individuals

### Personal Protective Equipment (PPE) Applicators and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- · Chemical-resistant gloves made of any waterproof
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading
- For exposure in enclosed areas, a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC 23-C), or a canister approved for pesticides (MSHA/NIOSH approval prefix TC-14G)
- For exposure outdoors, dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)

Wash thoroughly with soap and water after handling. Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **Engineering Controls Statement**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### **User Safety Recommendations**

#### **Users should:**

- · Wash hands before eating, drinking, chewing
- gum, using tobacco, or using the toilet.
   Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **Environmental Hazards**

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the high water mark. Do not contaminate water when disposing of equipment washwaters.

### **Directions For Use**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other

persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe. consult the agency responsible for pesticide

This product is not permitted for residential use. This product may only be used as follows:

Commercial: lawn and landscape areas at business and office complex sites, and turf at professional sports complexes or arenas.

Industrial: lawn and landscape areas at manufacturing sites

Golf Course: tee boxes, greens and turf mowed at 1" or less

Sod Farms: mechanically harvested turf only

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants.
- Chemical resistant gloves made of any waterproof material
- Chemical-resistant footear plus socks
- Protective eyewear
- Chemical-resistant headwear for overhead exposure

### **Nonagricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouses.

### Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Do not store below 32° F or above 100° F. Store in a dry place away from heat or open flame.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### Container Disposal:

• Plastic Containers: Triple rinse container (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local,



authorities, by burning. If burned, stay out of smoke.

• Water-soluble packaging: This package contains water-soluble bags inside a foil liner (outer bag). The bags will dissolve in water and the contents will disperse. If all the water-soluble bags are not used, carefully reseal the full bag. Do not remove the water-soluble bags from the overwrap bags except for immediate use. If exposed to moisture, the water-soluble bags may break

**Emergency Telephone Numbers** 

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact:

800-424-9300 CHEMTREC (Transportation and spills)
BASF Corporation 800-832-HELP

# Steps to be taken in case material is released or spilled:

Keep the spilled product dry to reduce volume. Scoop up the recaptured dry product and return it to the original container. Apply product according to normal label directions. If the spilled product gets wet, add a drying agent to contain the spill and dispose of as directed.

### I. General Information

**Curalan® EG fungicide** is effective on all turfgrasses. Refer to the appendix for a list of diseases controlled.

### Mode of Action

**Curalan EG** is a contact fungicide. It should be applied preventatively to protect susceptible plant tissues from infection. **Curalan EG** inhibits spore germination. If spores have already germinated, **Curalan EG** will prevent any further growth of the germ tube.

**Crop Tolerance** 

All listed plants are tolerant to **Curalan EG** at listed stages of growth. Consult your local representative or agricultural advisor for assistance.

Notice to user: Due to variability within species, and in application techniques and equipment, and the number of possible tank mix combinations, neither the manufacturer nor the seller has determined whether or not Curalan EG can safely be used on all varieties and species of turf under all conditions. It is recommended, therefore, that the professional user determine if Curalan EG can be used safely before broad use by applying the recommended use rate of Curalan EG on a nonlabeled species or variety under the conditions expected to be encountered on a small test area. Any adverse effects should be visible within 14 days.

Irrigation

Allow foliage to dry after application and before irrigation.

Coverage

Thorough coverage of all foliage is important, therefore enough spray volume and pressure must be used to achieve desired disease control.

Cleaning Spray Equipment

Clean application equipment thoroughly using a strong detergent or commercial sprayer cleaner according to

the manufacturer's directions before and after applying this product

### II. Application Instructions

Apply Curalan EG as instructed by section VII. Crop-Specific Information

### III. Additives

To achieve consistent disease control, an additive, such as a wetting agent, may be needed. Refer to section **VII. Crop-Specific Information** for more details. Consult your local BASF representative for recommendations for your area.

**Compatibility Test for Mix Components** 

Add components in the following sequence using 2 teaspoons for each pound or 1 leaspoon for each pint of recommended label rate per acre.

- Water: For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
- Products in PVA bags: Cap the jar and invert 10 cycles.
- Water-dispersible products: (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions) Cap the jar and invert 10 cycles.
- 4) Water-soluble products: Cap the jar and invert 10 cycles.
- Emulsifiable concentrates: (oil concentrate or methylated seed oil when applicable) Cap the jar and invert 10 cycles.
- 6) Water-soluble additives: Cap the jar and invert 10 cycles.
- Let the solution stand for 15 minutes.
- 8) Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. Do not use any spray solution that could clog spray nozzles.



### IV. Mixing Order

- 1) **Water:** Begin by agitating a thoroughly clean sprayer tank half full of clean water.
- 2) Products in PVA bags: Place the water-soluble PVA bag into the mixing tank. The water-soluble PVA bag will dissolve in water to allow the contents to disperse. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing. To prepare spray solution for aerial application, use a mixing tank or mixing vat first to get product into suspension before transferring suspension to air application equipment.
- Water-dispersible products: (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
- 4) Water-soluble products:
- Emulsifiable concentrates (oil concentrate or methylated seed oil when applicable)
- 6) Water-soluble additives (AMS or UAN when applicable)
- 7) Remaining quantity water

Maintain constant agitation during application. For more information, refer to section **V. Tank Mixing Application**.

### V. General Tank Mixing Information

See section VII. Crop-Specific Information for more details. Read and follow the applicable Restrictions and Limitations and Directions For Use on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

Tank Mix Partners/Components

The following products may be tank mixed with **Curatan® EG fungicide** according to the specific tank mixing instructions in this label and respective product labels.

#### Spotrete\* fungicide

BASF does not recommend using tank mixes other than those listed on BASF labeling. Physical incompatibility, reduced disease control, or crop injury may result from mixing **Curalan EG** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. Local agricultural authorities may be a source of information when using other than BASF recommended tank mixes.

# VI. General Restrictions and Limitations - All Crops

- Maximum seasonal use rate: Do not apply more than a total of 8 pounds of Curalan EG per acre (3 ounces per 1,000 square feet), per season.
- Preharvest Interval (PHI): Do not harvest turf for transplant into residential areas until 24 days have elapsed from the last application.
- Restricted Entry Interval (REI): 12 hours.
- Refer to section VII. Crop-Specific Information for sequential application intervals for Curalan EG.
- Do not apply to areas likely to be grazed by livestock and do not feed clippings to livestock or poultry.
- Do not apply through any type of irrigation equipment.
- This product cannot be used to **formulate** or reformulate any other peshcide product.

Table 1. Crop-Specific Restrictions and Limitations

Crop	Minimum Time from Application to Harvest (PHI)	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application
Turfgrasses	n/a	8 pounds	No	No

## VII. Crop-Specific Information

## Turfgrasses

Rate: Apply Curalan® EG fungicide at the rates indicated in Table 2 in 1-4 gallons of water per 1,000 square feet (40-160 gallons per acre) after mowing. The preventive rate of Curalan EG should be applied before the disease is expected to appear. Apply subsequent treatments at the preventative rate once control has been achieved

Table 2. Application Rates and Timing for Turfgrasses

Disease	Application Intervals*	Preventative Treatment		
		Rate per Acre	Rate per 1,000 Square Feet	
Dollar Spot	21-28 days	2.7 pounds	1 ounce	
Leaf Spots and Melting Out	14-28 days	2.7 pounds	1 ounce	
Red Thread and Pink Patch	14-28 days	2.7 pounds	1 ounce	
Brown Patch	14-28 days	2.7 pounds	1 ounce	
Gray Snow Mold**	10-21 days	2.7 pounds	1 ounce	
Fusarium Patch & Pink Snow Mold"	10-21 days	2.7 pounds	1 ounce	

Re-treat at shorter intervals or use higher rates if conditions favorable to disease exist. Tank mix with Fungo\* fungicide or Spotrete\* fungicide for improved control.

### Crops

This product can be used on the following crops:

### **Turfgrasses**

Look inside for complete Restrictions and Limitations and Application Instructions.

Diseases listed in this label:			
Common Name	Scientific Name		
Brown Patch Dollar Spot	Rhizoctonia solani Sclerotinia homoeocarpa; Lanzia sp.;		
Fusarium Patch Pink Snow Mold Gray Snow Mold Helminthosporium Leaf Spots	Moellerodiscus sp. Fusarium nivale Microdochium Typhula incarnata Helminthosporium Drechslera spp.		
Melting Out Pink Patch Red Thread	Bipolaris sp. Limonomyces roseipellis Laetisaria fuciformis Ciborinia		

### Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer. BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions** For Use, subject to the inherent risks, referred to above, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASE and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASE

**WARNING:** This product contains a chemical known to the State of California to cause birth defects or other reproductive harm.

**WARNING:** This product contains a chemical known to the State of California to cause cancer.

Curalan and the Specialty Products logo are registered trademarks of BASF Corporation.

Fungo is a registered trademark of Scotts-Sierra Crop Protection Company.

Spotrete is a trademark of W.A. Cleary Corporation

© 2003 BASE Corporation All rights reserved 007969-00085.20031202.NVA 2003-04-012-0135.pdf Supersedes NVA 99-4-12-0047

> BASE Corporation 26 Davis Drive Research Triangle Park, NO 27 109

**Agricultural Products** 



December 3, 2003

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U.S. Environmental Protection Agency
Ariel Rios Building
1200 Pennsylvania Avenue, N.W.
Washington, DC 20460

Agricultural Froducts

RE: EPA Registration Number 7969-85 Notification of Minor Label Change as sper PR Notice 98-10.

Dear Sir:

BASF has found a typographical error in our Curalan® Fungicide label and is notifying the Agency of its correction. Specifically on page 2 of the label under the heading **Notice to user** the following statement appears:

"Due to variability within species, and in application techniques and equipment, and the number of possible tank mix combinations, neither the manufacturer nor the seller has determined whether or not **Curalan EG** can safely be used on all varieties and species of nonbearing food crops, trees, shrubs, ornamentals, bedding plants, ground covers, nursery, turf and other non-food crops under all conditions. It is recommended, therefore, that the professional user determine if **Curalan EG** can be used safely before broad use by applying the recommended use rate of **Curalan EG** on a nonlabeled species or variety under the conditions expected to be encountered on a small test area. Any adverse effects should be visible within 14 days."

This statement has been changed to read

Due to variability within species, and in application techniques and equipment, and the number of possible tank mix combinations, neither the manufacturer nor the seller has determined whether or not Curalan EG can safely be used on all varieties and species of turf under all conditions. It is recommended, therefore, that the professional user determine if Curalan EG can be used safely before broad use by applying the recommended use rate of Curalan EG on a nonlabeled species or variety under the conditions expected to be encountered on a small test area. Any adverse effects should be visible within 14 days.

We have made this change because the label does not permit the use on nonbearing food crops, trees, shrubs, ornamentals, bedding plants, ground covers, nursery, and other non-food crops. This reference should have been deleted when BASF voluntarily deleted those uses from the label.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in

violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

I am enclosing five copies of the modified label for your records. Should you have any questions regarding this notification, please call me at (919) 547-2974.

Sincerely,

BASF Corporation Agricultural Products

T.R. Nelsen, Ph.D.

Manager, Regulatory Affairs