

Poast Plus®

herbicide

Directions For Use for Poast Plus alone to control annual grasses in soybeans in the Midwest (IA, IL, IN, KS, MN, MO, NE, OH, and SD)*

EPA Reg. No. 7969-88

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of herbicide application

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

General Information

Poast Plus® herbicide is an improved formulation of Poast® herbicide that contains a unique adjuvant which improves user convenience. Poast Plus without additive is intended for the postemergence control of annual grasses in soybeans. Poast Plus is effective through postemergence contact and systemic activity. For effective control, grasses must be thoroughly covered with spray solution.

Cultivation:

Do not cultivate within five days before application of Poast Plus or 7 days following application. Cultivation may put grass under

stress and reduce overall control. A timely cultivation after applying Poast Plus may assist grass control in soybeans.

Application Rate and Timing

Apply Poast Plus at 24 ounces per acre early postemergence when weeds are small and actively growing (generally when soybeans are in the 1st to 2nd trifoliate leaf stage of growth).

Poast Plus used alone at the 24 ounces per acre rate will control annual grasses and volunteer corn as listed in the following table. No additional additive is needed with the 24 ounces per acre rate of Poast Plus to achieve complete control of these grasses.

Recommendations

For grasses other than those listed on the previous table, consult the current label for Poast Plus.

Ground Application

Use 5-20 gallons of water per

broadcast acre at a minimum of 40 psi pressure (measured at the boom, not at the pump or in the line) to ensure adequate spray coverage. Use standard high pressure pesticide hollow cone or flat fan nozzles spaced 20 inches apart. Do not use flood or whirl chamber nozzles.

Applicator must follow the most restrictive use cautions to avoid drift hazard and must follow labeling as well as applicable state and local regulations and ordinances.

Restrictions and Limitations

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix. Do not apply to weeds under stress, such as stress due to lack of moisture, previous herbicide injury, mechanical injury or cold temperatures, as unsatisfactory control could result. Do not apply if rainfall or irrigation is

Grass Controlled	Height (inches)	Application Rate
Barnyardgrass Fall Panicum Foxtails: Giant, Green, Yellow Woolly Cupgrass Wild Proso Millet	Up to 8	24 fluid ounces
Shattercane/Wildcane If needed, re-treat at the same rate and stage of growth.	6-18	
Volunteer Corn Maintain sufficient boom height above volunteer corn	Up to 20	

ACCEPTED

APR 20 1994

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7969-88

BEST AVAILABLE COPY

Agricultural Products

BASF

exp. ted within one hour following application.

Physical incompatibility, reduced weed control, or crop injury may result from mixing Poast Plus with other pesticides (fungicides, herbicides, insecticides or miticides), additives or fertilizers not recommended on the label. BASF does not recommend the use of Poast Plus in tank mixes other than those listed on BASF labels, supplemental labels, or technical bulletins. Local agricultural authorities may be a source of information when using other than BASF recommended combinations. Do not apply Poast Plus as a pre-plant or preemergent treatment prior to corn, millet, sorghum, or small grain crops.

Do not apply Poast Plus through any type of irrigation system.

Do not apply to soybeans within 75 days of harvest.

() not graze treated soybean fields () do not feed treated soybean forage (green succulent) or ensilage to livestock.

() bean hay may be fed.

Do not apply Poast Plus to soybeans under stress due to lack of moisture, previous, herbicide injury, mechanical injury, or cold temperatures, as crop injury may result.

Do not apply more than a total of 7.5 pints each of Poast Plus per acre in one season.

* The specific program area includes the states of IA, IL, IN, OH, and SD as well as portions of:

Nebraska including the counties
() Jams, Antelope, Blaine, Boone, Boyd, Brown, Buffalo, Burt, Butler, Cass, Cedar, Clay, Colfax, Cuming, Custer, Dakota, Dawson, Dixon, Dodge, Douglas, Fillmore, Franklin, Frontier, Furnas, Gage, Garfield, Gosper, Greeley, Hall, Hamilton, Harlan, Holt, Howard, Jefferson, Johnson, Kearney, Keya Paha, Knox, Lancaster, Loup, Madison, Merrick, Nance, Nemaha, Nuckolls, Otoe, Pawnee, Phelps, Pierce, Platte, Polk, Red Willow, Richardson, Rock, Saline, Sarpy, Saunders, Seward, Sherman, Stanton, Thayer, Thurston, Valley, Washinaton, Wayne, Webster, Wheeler, and York.

Kansas including the counties of
Allen, Anderson, Atchison, Bourbon, Brown, Butler, Chase, Chautaugua, Cherokee, Clay, Cloud, Coffey, Cowley, Crawford, Dickinson, Doniphan, Douglas, Elk, Franklin, Geary, Greenwood, Harvey, Jackson, Jefferson, Jewell, Johnson, Labette, Leavenworth, Linn, Lyon, McPherson, Marion, Marshall, Miami, Montgomery, Morris, Nemaha, Neosho, Ness, Osage, Ottawa, Phillips, Pottawatomie, Republic, Riley, Saline, Sedgwick, Shawnee, Smith, Sumner, Wabaunsee, Washington, Wilson, Woodson and Wyandotte.

Missouri excluding the counties of the Missouri Bootheel: Butler, Dunkin, Mississippi, New Madrid, Pemiscot, Ripley, Scott, and Stoddard.

Minnesota including the counties of
Big Stone, Blue Earth, Brown, Carver, Chippewa, Cottonwood, Dodge, Faribault, Fillmore, Freeborn, Goodhue, Houston, Jackson, Kandigohi, Lac Qui Parle, LeSueur, Lincoln, Lyon, Martin, McLeod, Meeker, Mower, Murry, Nicollet, Nobles, Olmsted, Pipestone, Redwood, Renville, Rice, Rock, Scott, Sibley, Steele, Stevens, Swift, Wabasha, Waseca, Watonwan, Winona, and Yellow Medicine.

2 7 2
Conditions of Sale and Warranty
The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Poast Plus is a registered trademark of BASF Corporation.

© 1994 BASF Corporation

XX ----- JJ9118 A1581193

BEST AVAILABLE COPY

BASF Corporation
P.O. Box 13528
Research Triangle Park, NC 27709

BASF