

7969-88

06-28-2007

1/20

 **BASF**
The Chemical Company

ACCEPTED
JUN 28 2007
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 7969-88

Poast Plus[®]

herbicide

Active Ingredient:

sethoxydim: 2-[1-(ethoxyimino)butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one* 13.0%

Inert Ingredients: 87.0%

Total: 100.0%

*Equivalent to 1.0 pounds of sethoxydim per gallon

EPA Reg. No. 7969-88

EPA Est. No. _____

KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

See next page for additional **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

Net Contents: _____

BASF Corporation, Agricultural Products
26 Davis Drive, Research Triangle Park, NC 27709

2/20

FIRST AID

If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye injury. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. For more options, refer to **Category E** on an EPA chemical-resistance category-selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to aquatic organisms. For terrestrial uses, **DO NOT** apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters.

Endangered Species Concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of federal law. This pesticide is toxic to vascular plants and should be used strictly in accordance with drift precautions on this label in order to minimize off-site exposures.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Unless otherwise stated in supplemental labeling, all applicable directions, restrictions and precautions are to be followed. This labeling must be in the user's possession during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, or viton \geq 14 mils
- Shoes plus socks

Storage and Disposal

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage: DO NOT store below 32° F or above 100° F. Store in a dry place away from heat or open flame. Avoid contamination of feed or foodstuffs.

Pesticide Disposal: Pesticide wastes are toxic. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Bulk/Mini-bulk Containers:

Refillable/reusable containers should be returned to the point of purchase for cleaning and refilling.

In Case of Emergency

In case of large-scale spillage regarding this product, call:

CHEMTREC 1-800-424-9300
BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment.
- Your local poison control center (hospital).
- BASF Corporation 1-800-832-HELP (4357).

Steps to be taken in case material is released or spilled:

Wear the personal protective equipment specified on this label. Recover the material for reuse according to label whenever possible. Cover the liquid with an absorbent material (such as pet litter). Sweep up and place in an appropriate container for disposal. Remove and wash clothing and personal protective equipment prior to reuse. Keep the spill out of all sewers and open bodies of water.

I. General Information

Poast Plus® herbicide is a selective, broad spectrum, postemergence herbicide for control of annual and perennial grass weeds. **Poast Plus** does not control sedges or broadleaf weeds. Essentially, all grass crops, such as sorghum, corn, small grains, and rice, as well as ornamental grasses, such as turf, are susceptible to **Poast Plus**.

Mode of Action

Poast Plus rapidly enters the target weed through its foliage and translocates throughout the plant. The effects range from slowing or stopping growth (generally within 2 days), to foliage reddening and leaf tip burn. Subsequently, foliage burnback may occur. These symptoms will generally be observed within 3 weeks depending on environmental conditions.

Crop Tolerance

All labeled crops are tolerant to **Poast Plus** at all stages of growth.

Herbicide Resistance

Repeated use of **Poast Plus** (or similar postemergence grass herbicides with the same mode of action) may lead to the selection of naturally occurring biotypes with resistance to these products. If poor performance cannot be attributed to adverse weather conditions or improper application methods, a resistant biotype may be present. Consult your local representative or agricultural advisor for assistance.

Irrigation

In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth.

Cultivation

DO NOT cultivate within 5 days before or 7 days after applying **Poast Plus**. Cultivating 7 days or later after treatment may help provide season-long control.

3/20

Cleaning Spray Equipment

Clean spray equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product.

II. Application Instructions

Applications can be made to actively growing weeds as aerial, broadcast, band, or spot spray applications at the rates and growth stages listed in **Tables 1, 2 and 3**, unless instructed differently in section **VI. Crop-Specific Information**. The most effective control will result from making postemergence applications of **Poast Plus® herbicide** early, when weeds are small. Delaying application permits weeds to exceed the maximum size stated and may prevent adequate control.

Apply **Poast Plus** to the foliage of grasses uniformly and completely because large leaf canopies shelter smaller weeds and can prevent adequate spray coverage. **DO NOT** spray to the point of runoff.

Spray Drift Management

General Information Pertaining to Aerial and Ground Applications

Make aerial or ground application when the wind velocity favors on-target product deposition. Apply only when the wind speed is less than or equal to 10 mph. For all non-aerial applications, wind speed must be measured to the application site on the upwind side immediately prior to application. **DO NOT** make aerial or ground applications into areas of temperature inversions. Inversions are characterized by stable air and increasing distance above the ground. Mist or fog may indicate the presence of an inversion in humid areas. When permissible by local regulations, the applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

Aerial Application Methods and Equipment

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

DO NOT apply under circumstances where possible drift to unprotected persons, to food, forage, or other plantings that might be damaged, or crops thereof rendered unfit for sale, use or consumption can occur.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements **DO NOT** apply to forestry applications, public health uses or to applications using dry formulations.

- 4/20
1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
 2. Nozzles must always point backward parallel with the air stream and never be pointed downward more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the [Aerial Drift Reduction Advisory Information](#).

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. Use the largest droplet size consistent with acceptable efficacy. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly or under unfavorable environmental conditions (see **Wind, Temperature and Humidity**, and **Temperature Inversions**).

Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. For ground boom and aerial applications, use medium or coarser spray nozzles according to ASAE 572 definition for standard nozzles or a volume mean diameter (VMD) of 300 microns or greater for spinning atomizer nozzles.

Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets. Use a minimum of 5 gallons of water per acre. Increase water volume to a least 10 gallons of water per acre if grass foliage or crop canopy is dense.
- **Pressure** - **DO NOT** exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid-stream nozzles oriented straight back produce the largest droplets and the lowest drift.

5/20

Boom Length

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential.

NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

Low humidity and high temperatures increase the evaporation of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperatures. When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, or nontarget crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Ground Application Methods and Equipment (Broadcast)

For ground boom applications, apply with nozzle height no more than 4 feet above ground or crop canopy.

DO NOT apply when conditions favor drift from target area or when windspeed is greater than 10 mph.

Water Volume: Use 5-20 gallons of spray solution. In the West and in the High and Rolling Plains Region, (see regional descriptions in **Table 1**), **DO NOT** use less than 10 gallons of spray solution per acre.

Spray Pressure: Use 40-60 psi (measured at the boom, not at the pump or in the line). When crop and weed foliage are dense, use a maximum of 20 gallons of water and 60 psi.

Application Equipment: Use standard high-pressure pesticide flat fan or hollow cone nozzles spaced up to 20 inches apart. **DO NOT** use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles as erratic coverage can cause inconsistent weed control. When tall weeds such as volunteer corn are to be controlled, the boom should be high enough to cover the entire plant. Refer to the nozzle manufacturer's directions for recommended height. When a crop such as cotton is 24 inches or taller and the grasses are below the crop canopy, use drop nozzles to ensure good coverage of the grass species.

DO NOT use selective application equipment such as recirculating sprayers or wiper applicators.

Ground Application (Banding)

Poast Plus may be applied by banding to control annual grasses. Banding is not recommended for perennial grasses.

Follow **Ground Application Methods and Equipment (Broadcast)** instructions for band applications. When applying **Poast Plus** by banding, determine the amount of herbicide and water volume needed using the following formula:

Bandwidth in inches	x	Broadcast rate per acre	=	Banding herbicide rate per acre
Row width in inches				
Bandwidth in inches	x	Broadcast volume per acre	=	Banding water volume per acre
Row width in inches				

Spot or Small Area Application

DO NOT make spot treatments in addition to broadcast or band treatments. When using knapsack sprayers or high-volume spray equipment with hand guns or other suitable nozzle arrangements, prepare a 1 to 1.5% solution of **Poast Plus**® herbicide in water unless otherwise specified under specific crops. Use a concentration of 0.5% for **Dash**® HC and **Sundance**® HC spray adjuvants, or 1% for oil concentrate. Prepare the desired volume of spray solution by mixing the amount of **Poast Plus** and the amount of **Dash HC**, **Sundance HC** or oil concentrate in water according to **Tables 5** and **6**.

Rescue Treatment for Controlling Selected Annual Grasses

If **Poast Plus** cannot be applied at the recommended time, larger annual grasses may be controlled with a later application by increasing the rate of **Poast Plus** (see **Table 3**). **DO NOT** exceed the maximum rate per acre, per season, for specific crops (see **Table 7**).

III. Additives

To achieve consistent weed control, always use one of the following additives: **Dash HC**, **Sundance HC**, methylated/modified seed oil, or crop oil concentrate. In addition, urea ammonium nitrate or ammonium sulfate is recommended for use on alfalfa, beans, cotton, flax, peanuts, peas, potatoes, soybeans, **Poast Protected**™ field corn, **Poast Protected** sweet corn, sugar beets, and sunflowerers to enhance activity on certain grass species. See **Table 4. Additive Rates Per Acre** for more information. However, when used in many vegetable crops under the following conditions, **Poast Plus** plus adjuvants should be used with caution due to potential crop leaf injury: when the temperature exceeds 90° F and the relative humidity is 60% or greater, or anytime the temperature exceeds 100° F, regardless of the humidity.

Because most nitrogen solutions are mildly corrosive to galvanized, mild steel, and brass spray equipment, rinse the entire spray system with water soon after use. UAN and AMS are not registered in California.

Consult a BASF representative or local agricultural authority for more information on the use of additives.

Dash HC, Sundance HC, Crop Oil Concentrate, or Methylated Seed Oils

A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:

- be non-phytotoxic,
- contain only EPA-exempt ingredients,
- provide good mixing quality in the jar test, and
- be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers to provide good mixing quality.

Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. For more information, see

6/20
Compatibility Test for Mix Components. For most crops, **Dash HC** or **Sundance HC** may be substituted for crop oil concentrate or methylated seed oil; however, for some crops and tank mixes, **Dash HC**, **Sundance HC** and MSO are not recommended. (See section **VI. Crop-Specific Information** for more information.)

Nitrogen Source

- **Urea Ammonium Nitrate (UAN):** Commonly referred to as 28%, 30%, or 32% nitrogen solution, UAN may be used in addition to **Dash HC**, **Sundance HC**, or crop oil concentrate to improve weed control. **DO NOT** use brass or aluminum nozzles when spraying UAN.
- **Ammonium Sulfate (AMS):** AMS per acre may be substituted for UAN. When liquid AMS is used, 3.0 quarts of 8-8-0 analysis may be substituted for 2.5 pounds of dry AMS. Use high-quality AMS (spray grade) to avoid plugging of nozzles. Other sources of nitrogen are not as effective as those mentioned. If the AMS is added directly to the spray tank, add slowly while agitating. Adding the mix too quickly may clog outlet lines. Be sure the AMS is completely dissolved before adding any other products. BASF does not recommend applying AMS if applied in less than 10 gallons per acre because of potential problems with precipitation in reduced volumes.

UAN and AMS are not registered for use in California.

IV. General Tank Mixing Information

Tank Mix Partners/Components

The following products, listed with associated common name, may be tank mixed with **Poast Plus** according to the specific tank mixing instructions in this label and respective product labels.

- atrazine
- **Basagran**®/bentazon
- **Betamix**®/desmedipham + phenmedipham
- **Betanex**®/desmedipham
- **Blazer**®/acifluorfen
- **Bronate**®/bromoxynil + MCPA
- **Buctril**®/bromoxynil
- **Clarity**®/dicamba
- **Classic**®/chlorimuron
- **Cobra**®/lactofen
- **Dual**®/metolachlor
- **Dual II**®/metolachlor
- **FirstRate**®/cloransulam-methyl
- **Flexstar**®/fomesafen
- **Frontier**®/dimethenamid
- **Galaxy**®/bentazon + acifluorfen
- glyphosate (e.g. **Roundup**)
- **Guardman**®/atrazine + dimethenamid
- **Harness**®/acetochlor
- **Laddok**® S-12/bentazon + atrazine
- **Liberty**®/glufosinate
- **Marksman**®/atrazine + dicamba
- MCPA
- metribuzin
- **Outlook**® dimethenamid-P
- **Pursuit**®/imazethapyr
- **Pursuit**® DG/imazethapyr
- **Pursuit**® W/imazethapyr
- **Pursuit**® W DG/imazethapyr
- **Raptor**®/imazamox
- **Reflex**®/fomesafen
- **Reliance**® STS®/chlorimuron + thifensulfuron
- **Resource**®/flumiclorac
- **Sencor**® DF/metribuzin
- **Staple**®/pyrithiobac
- **Stellar**®/flumiclorac + lactofen
- **Stinger**®/clopyralid
- **Storm**®/bentazon + acifluorfen
- **Surpass**®/acetochlor
- **Synchrony**® STS/ chlorimuron + thifensulfuron
- **Touchdown**®/sulfosate
- **UpBeet**®/triflusaluron
- 2,4-D amine
- 2,4-DB
- 2,4-D (LVE)

7/20

See section **VI. Crop-Specific Information** for more details. Read and follow the applicable Restrictions and Limitations and **Directions For Use** on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

Separate applications should be made if all target weeds are not at the labeled growth stage for treatment at the same time.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Poast Plus® herbicide** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. BASF does not recommend using tank mixes other than those listed on BASF labeling. Local agricultural authorities may be a source of information when using other than BASF recommended tank mixes.

Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test.

For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Only use water from the intended source at the source temperature.

Add components in the sequence indicated in the **Mixing Order** using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, **DO NOT** mix the ingredients in the same tank.

Mixing Order

- 1) **Water.** Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
- 2) **Agitation.** Maintain constant agitation throughout mixing and application.
- 3) **Products in PVA bags.** Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
- 4) **Water-dispersible products** (such as dry flowables, wettable powders, suspension concentrates, or suspensions).
If an inductor is used, rinse it thoroughly after the component has been added.
- 5) **Water-soluble products.** If an inductor is used, rinse it thoroughly after the component has been added.

- 6) **Emulsifiable concentrates** (such as **Poast Plus** or oil concentrate when applicable). If an inductor is used, rinse it thoroughly after the component has been added.
- 7) **Water-soluble additives** (such as AMS or UAN when applicable). If an inductor is used, rinse it thoroughly after the component has been added.
- 8) **Remaining quantity of water.**

Maintain constant agitation during application.

8/20

Table 1. Standard Application Rates and Timing - Annual Grasses

All application rate and timing recommendations are based on growing region. Therefore, refer to the maps below and descriptions below to ensure application accuracy. Follow the **Application Rate and Timing** tables for your region only. Refer to **Table 7** for the maximum allowable use rates for specific crop and use sites.

Annual Grass	Midwest, South, and Northeast		West & High and Rolling Plains	
	Maximum Height	Rate Per Acre (pints)	Maximum Height	Rate Per Acre (pints)
Barnyardgrass	8"	1.5	8"	2.25
Crabgrass, Large ¹	6"	1.5	4"	2.25
, Smooth ¹	6"	1.5	4"	2.25
Cupgrass, Southwestern	—	—	8"	2.25
, Woolly	8"	1.5	—	—
Fescue, Tall (seedling)	6"	2.25	—	—
Foxtail, Giant	8"	1.5	8"	2.25
, Green	8"	1.5	8"	2.25
, Yellow	8"	1.5	8"	2.25
Goosegrass	6"	1.5	4"	2.25
Itchgrass	4"	3.0	—	—
Johnsongrass (seedling)	8"	1.5	8"	2.25
Junglerice	8"	1.5	8"	2.25
Lovegrass	6"	2.25	—	—
Millet, Wild Proso	10"	0.75	10"	1.5
Oats, Tame	6"	2.25	—	—
, Wild ¹	4"	1.5	4"	2.25
Orchardgrass (seedling)	6"	2.25	—	—
Panicum, Browntop	8"	1.5	8"	2.25
, Fall	8"	1.5	8"	2.25
, Texas	8"	1.5	8"	2.25
Red Rice ¹	4"	3.0	—	—
Ryegrass, Annual	8"	1.5	8"	2.25
Sandbur, Field	3"	1.875	—	—
Shattercane/Wildcane ¹	18"	1.5	18"	2.25
Signalgrass, Broadleaf	8"	1.5	8"	2.25
Sprangletop, Red ³	8"	1.5	8"	2.25
Stinkgrass	6"	2.25	—	—
Volunteer ^{2,4} Barley ¹	4"	2.25	4"	3.0
Corn ¹	20"	1.5	12"	2.25
Oats ¹	4"	2.25	4"	3.0
Rye ¹	4"	2.25	4"	3.0
Wheat ¹	4"	2.25	4"	3.0
Witchgrass ¹	8"	1.5	8"	2.25

¹ Add nitrogen to the crop oil concentrate to improve grass control on indicated species.

² Apply **Poast Plus**[®] herbicide before tillering.

³ **Poast Plus** is not recommended for use on red sprangletop in California, Arizona, or western New Mexico.

⁴ In the West Region, volunteer cereals that emerge from late spring through early summer (May through July) may be partially or incompletely controlled because of unfavorable conditions at application time.

Regional Descriptions

West & High and Rolling Plains: An area of the western United States, including western Texas, Oklahoma and Kansas, west of a line running north from Del Rio to Gainesville, Texas, and extending along Interstate 35 to the Oklahoma-Kansas border, then west along border to Highway 83 and then north to the Kansas-Nebraska border, west to Colorado, all of Colorado to the Continental Divide, then west of the Continental Divide North to the US-Canada border.

Midwest, South, and Northeast: All other regions not listed above.

9/20

Table 2. Standard Application Rates and Timing - Perennial Grasses¹

All application rate and timing recommendations are based on growing region. Therefore, refer to the maps below and descriptions in **Table 1** to ensure application accuracy. Follow the **Application Rate and Timing** tables for your region only. Refer to **Table 7** for the maximum allowable use rates for specific crop and use sites.

Perennial Grass	Midwest, South, and Northeast		West & High and Rolling Plains	
	Maximum Height	Rate Per Acre (pints)	Maximum Height	Rate Per Acre (pints)
Bermudagrass	6" stolon	2.25	6" stolon	3.0 ² -3.75
Guineagrass	8"	3.75	—	—
Johnsongrass (Rhizome)	25"	2.25	10"	2.25 ² -3.75
Johnsongrass (No-Till)	20"	2.25	—	—
Muhly, Wirestem	6"	1.875	—	—
Quackgrass ¹	8"	2.25	8"	3.75
Ryegrass, Perennial	8"	2.25	8"	2.25
Torpedograss	8"	3.75	—	—
Sequential Application	Maximum Height	Rate Per Acre (pints)	Maximum Height	Rate Per Acre (pints)
Bermudagrass	4" stolon	1.5	4" stolon	2.25 ²
Guineagrass	8"	3.75	—	—
Johnsongrass (Rhizome)	12"	1.5	8"	1.5 ² -2.25
Johnsongrass (No-Till)	12"	1.5	—	—
Muhly, Wirestem	6"	1.875	—	—
Quackgrass ¹	8"	1.5	8"	2.25
Ryegrass, Perennial	8"	2.25	8"	2.25
Torpedograss	8"	3.75	—	—

¹ Add nitrogen to the crop oil concentrate to improve grass control on indicated species. Cultivate 7-14 days after an initial or sequential application to aid control.

² Use 3.75 pints per acre for the following forage crops: alfalfa, clover, birdsfoot trefoil, sainfoin.

Table 3. Special Application Rates and Timing for Midwest, South and Northeast

Annual Grass	Midwest, South, and Northeast			
	Special Early Maximum Height	Early Rate Per Acre (pints)	Rescue Maximum Height	Rescue Rate Per Acre (pints)
Barnyardgrass	4"	1.125 ¹	12"	2.25
Crabgrass, Large ³	—	—	8"	2.25
Smooth ³	—	—	8"	2.25
Foxtail, Giant ²	4"	1.125	16"	2.25
Green ²	4"	1.125	16"	2.25
Yellow ²	—	—	16"	2.25
Goosegrass	3"	1.125	8"	2.25
Johnsongrass (seedling)	—	—	16"	2.25
Millet, Wild Proso	10"	0.75	24"	1.5
Panicum, Fall	4"	1.125	12"	2.25
Texas	4"	1.125	12"	1.5
Signalgrass, Broadleaf	4"	1.125	12"	2.25
Volunteer Corn ³	12"	1.125	—	—

¹ In the following states use 1.5 pint: AL, AR, FL, GA, LA, MS, NC, SC, TN, TX, and VA.

² For flax, use 0.75 pint per acre when foxtails are less than 1.5" high. When using the special early rate, the foxtail species should not have started to tiller.

³ Add nitrogen to the crop oil concentrate to improve grass control on indicated species.

10/20

Additive	Ground Application	Aerial Application
AMS	2.5 pounds	2.5 pounds
Dash® HC/Sundance® HC	1.0 pint	1.0 pint
Crop Oil Concentrate	2.0 pints	2.0 pints
Methylated Seed Oils/MSO	1.5 pints	1.5 pints
UAN Solution	4.0-8.0 pints	4.0-8.0 pints

Spray Solution Volume	Amount of Product to be Added					
	Poast Plus® herbicide (1.0%)	or	Poast Plus (1.5%)	Oil Concentrate (1.0%)	or	Dash HC/Sundance HC (0.5%)
1 gallon	1.9 fl oz		2.9 fl oz	1.3 fl oz		0.6 fl oz
3 gallons	5.8 fl oz		8.75 fl oz	3.8 fl oz		1.9 fl oz
5 gallons	9.6 fl oz		14.5 fl oz	6.4 fl oz		3.2 fl oz
25 gallons	3.0 pints		4.5 pints	2.0 pints		1.0 pint
50 gallons	6.0 pints		9.0 pints	4.0 pints		2.0 pints
100 gallons	12.0 pints		18.0 pints	8.0 pints		4.0 pints

2 tablespoons = 1 fluid ounce

Grass (see Tables 1, 2, and 3 for the complete list of grasses controlled)	Concentration in Spray Solution ¹			
	Poast Plus	Crop Oil Concentrate/ Methylated Seed Oil	or	Dash HC/ Sundance HC
Annual grasses up to 6" height	1.0%	1.0%		0.5%
Annual grasses up to 12" height	1.5%	1.0%		0.5%
Perennial grasses²	1.5%	1.0%		1.0%

¹ Refer to Table 5. Spot Treatment Dilution for preparing the desired solution volume.
² Repeat application as needed.

V. General Restrictions and Limitations - All Crops

- **Maximum seasonal use rate:** See Table 7 for crop-specific maximum seasonal use rates.
- **Preharvest Interval:** See Table 7 for crop-specific preharvest intervals.
- **Restricted-Entry Interval (REI): 12 hours**
- Avoid all direct or indirect contact with any desired grass crop unless otherwise recommended on the **Poast Plus** label.
- **Stress: DO NOT** apply to grasses or crops under stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control may result. In irrigated areas, it may be necessary to irrigate before application to insure active weed growth.
- **DO NOT** apply to crops that show **injury** (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications because this injury may be enhanced or prolonged.
- **DO NOT** apply as a **preplant** or **preemergence treatment** before planting grass crops, such as corn, millet, or sorghum, unless otherwise specified on supplemental labeling.
- **DO NOT** use UAN or AMS in California.
- **DO NOT** use **selective application equipment** such as recirculating sprayers, wiper applicators, or shielded applicators.
- **Rainfast Period: Poast Plus** is rainfast **1 hour** after application.
- **DO NOT** apply through any type of **irrigation** equipment.
- **DO NOT** plant other crops to be harvested for 30 days after application unless **Poast Plus** is registered for use on that crop.

11/20

Table 7. Crop-Specific Restrictions and Limitations for Poast Plus® herbicide

Crop	Minimum Time From Application to Harvest (PHI)	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application
Alfalfa, birdsfoot trefoil, and sainfoin ¹	14 days before cutting for (dry) hay	3.75 pints	9.75 pints	Yes	Yes
Alfalfa, birdsfoot trefoil, and sainfoin ¹ (undried)	7 days before grazing, feeding, or cutting for (undried) forage	3.75 pints	9.75 pints	Yes	Yes
Apricot	25 days	3.75 pints	7.5 pints	n/a	No
Artichoke, Globe	7 days	3.75 pints	7.5 pints	No	Yes
Asparagus	1 day	3.75 pints	7.5 pints	No	Yes
Avocado (nonbearing) ¹	1 year	3.75 pints	11.25 pints	n/a	No
Beans ¹ , Dry	30 days	3.75 pints	6.0 pints	Yes	Yes
, Succulent	15 days	3.75 pints	6.0 pints	Yes	Yes
Beet (Garden)	60 days	3.75 pints	7.5 pints	No	Yes
Blueberry ¹	30 days	3.75 pints	7.5 pints	No	Yes
Brassica ¹ including: Broccoli (including Chinese & Raab), Brussels Sprouts, Cabbage (Bok Choy, Chinese Mustard, Napa), Cauliflower, Collards, Kale, Kohlrabi, Mustard Greens, Rape Greens	30 days ¹	2.25 pints	4.5 pints	No	Yes
Bulb Vegetables including: Garlic, Leek, Onion (Dry Bulb & Green), Shallot	30 days	2.25 pints	6.75 pints	No	Yes
Caneberries ¹ including: (All varieties and/or hybrids of these) Blackberry, Raspberry (Red, Black), Loganberry, Youngberry	45 days	3.75 pints	7.5 pints	No	Yes
Canola/Crambe/Rapeseed ¹	60 days	3.75 pints	7.5 pints	No	Yes
Carrot	30 days	3.75 pints	7.5 pints	No	Yes
Cherries (sweet and sour)	25 days	3.75 pints	7.5 pints	n/a	No
Citrus ¹	15 days	3.75 pints	15.0 pints	Yes	No
Clover	7 days before grazing, feeding, or cutting for (undried) forage	3.75 pints	9.75 pints	Yes	Yes
Clover hay	20 days before grazing, feeding, or cutting for (dry) hay	3.75 pints	9.75 pints	Yes	Yes
Corn (Poast Protected™ field corn) ¹	60 days (grain or fodder) 45 days (forage and silage)	2.25 pints	4.5 pints	Yes	Yes
Corn (Poast Protected™ sweet corn) ¹	45 days (grain or fodder) 30 days (fresh sweet corn or forage and silage)	2.25 pints	4.5 pints	Yes	Yes
Cotton ¹	40 days	3.75 pints	11.25 pints	No	Yes
Cranberry ¹	60 days	3.75 pints	6.75 pints	No	Yes
Cucurbits ¹ including: Cantaloupes (all), Cucumber, Gherkin, Honeydew Melon, Muskmelon (all), Pumpkin, Squash (all), Watermelon	14 days ¹	2.25 pints	4.5 pints	No	Yes
Date (nonbearing) ¹	1 year	3.75 pints	11.25 pints	n/a	No
Deciduous Trees, Nonfood Crop Areas, Fallow Land ¹	n/a	3.75 pints	n/a	No	No
Fescue, Tall ¹	n/a	3.75 pints	n/a	No	Yes
Fig (nonbearing) ¹	1 year	3.75 pints	11.25 pints	n/a	No
Flax ¹	75 days	2.25 pints	6.0 pints	Yes	Yes

12/20

Table 7. Crop-Specific Restrictions and Limitations for Poast Plus® Herbicide (Continued)

Crop	Minimum Time From Application to Harvest (PHI)	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application
Fruiting Vegetables ¹ including: Eggplant, Groundcherry, Pepino, Peppers (all) ¹ , Tomatillo, Tomato ¹	20 days ¹	2.25 pints	6.75 pints	No	Yes
Grape ¹	50 days	3.75 pints	7.5 pints	No	No
Head & Petiole Type Vegetables ¹ Cardoon, Celery ¹ , Celery (Chinese), Celtuce, Fennel (Florence), Lettuce (Head), Radicchio, Rhubarb ¹ , Swiss Chard	30 days ¹	2.25 pints	4.5 pints	No	Yes
Horseradish ¹	60 days	3.75 pints	7.5 pints	No	No
Leafy Vegetables Amaranth, Arugula, Chervil, Chrysanthemum (Edible, Garland), Cilantro, Corn Salad, Cress (Garden, Upland), Dandelion, Dock, Endive (Escarole), Lettuce (Leaf), Orach, Parsley, Purslane (Garden, Winter), Spinach (including New Zealand & Vine)	15 days	2.25 pints	4.5 pints	No	Yes
Lentil ¹	50 days	3.75 pints	6.0 pints	No	Yes
Lingonberry, Salal, Juneberry	45 days	3.75 pints	7.5 pints	No	Yes
Mint ¹	20 days	3.75 pints	7.5 pints	No	Yes
Nectarine	25 days	3.75 pints	7.5 pints	n/a	No
Olives (nonbearing) ¹	1 year	3.75 pints	11.25 pints	n/a	No
Orchard floor middles ¹	n/a	0.75 pint	0.75 pint	n/a	No
Peach	25 days	3.75 pints	7.5 pints	n/a	No
Peanut ¹	40 days	2.25 pints	3.75 pints	No	Yes
Peas, Dry , Succulent	30 days 15 days	3.75 pints 3.75 pints	6.0 pints 6.0 pints	Yes Yes	Yes Yes
Pistachio ¹	15 days	3.75 pints	15.0 pints	n/a	No
Plum (nonbearing) ¹	1 year	3.75 pints	11.25 pints	n/a	No
Pome Fruits ¹ including: Apples, Crabapples, Pears, and Quince	14 days	3.75 pints	11.25 pints	No	No
Pomegranate (nonbearing) ¹	1 year	3.75 pints	11.25 pints	n/a	Yes
Potatoes ¹ , Field , Sweet (East US) , Sweet (West US)	30 days 30 days 60 days	3.75 pints 3.75 pints 2.25 pints	7.5 pints 7.5 pints 7.5 pints	No No No	Yes Yes Yes
Prune (nonbearing) ¹	1 year	3.75 pints	11.25 pints	n/a	No
Tuberous & Corm Vegetables Arracacha, Arrowroot, Artichoke (Chinese, Jerusalem), Canna (Edible), Cassava (Bitter, Sweet), Chayote Root, Chufa, Dasheen (Taro), Ginger, Leren, Potato ¹ , Tanier, Turmeric, Yam Bean, Yam (True)	30 days	3.75 pints	7.5 pints	No	Yes
Safflower	30 days	3.75 pints	7.5 pints	No	Yes
Set Aside Conservation Land ¹	n/a	3.75 pints	11.25 pints	n/a	Yes
Soybean ¹	75 days	3.75 pints	7.5 pints	Only seed and hay	Yes
Strawberry ¹	7 days	3.75 pints	3.75 pints	No	Yes
Sugar Beet ¹	60 days	3.75 pints	7.5 pints	Yes	Yes
Sunflower ¹	70 days	3.75 pints	3.75 pints	No	Yes
Tobacco ¹	42 days	2.25 pints	6.0 pint	No	Yes
Tree Nuts ¹	15 days	3.75 pints	15.0 pints	No	No

¹ See VI. Crop-Specific Information for more details concerning use restrictions and PHI exceptions.
n/a = not applicable.

VI. Crop-Specific Information

Crops Grown For Seed

Poast Plus® herbicide is recommended for use on all crops on this label when they are grown for seed production. Use the **Poast Plus** rates given for each food crop listed in other sections on this label. Slight modifications in application methods may be required for certain seed crops due to crop canopy or different cultural methods from the corresponding food crop.

Field Crops

Always add 1.0 pint of **Dash**® HC or **Sundance**® HC spray adjuvant, or 2 pints of oil concentrate per acre. Add 4.0-8.0 pints of UAN or 2.5 pounds of AMS to control crabgrass, volunteer corn and all volunteer cereals. UAN and AMS are not registered in California.

Beans, Dry

Poast Plus may be applied in a tank mix with one of the following herbicides:

- **Basagran**®
- **Frontier**®

Canola/Crambe/Rapeseed

Not registered in California.
Processed meal may be fed.

Corn, Field

Only Poast Protected™ field corn hybrids are tolerant to **Poast Plus** applications. **Severe crop injury will occur to corn hybrids not designated as Poast Protected corn.**

Not for use in California.

Over-the-top applications of **Poast Plus** in **Poast Protected** field corn may be made until the onset of pollen shed provided the appropriate preharvest intervals are met. **DO NOT** apply **Poast Plus** after pollination occurs.

Poast Plus may be applied in a tank mix with one of the following herbicides:

- atrazine
- **Basagran**
- **Dual**® & **Dual II**®
- **Frontier**
- **Guardsman**®
- **Harness**®
- **Laddok**® S-12
- **Surpass**®
- 2,4-D (LVE)

Corn, Sweet

Only Poast Protected sweet corn hybrids are tolerant to **Poast Plus** applications. **Severe crop injury will occur to sweet corn hybrids not labeled as Poast Protected sweet corn.**

Applications of **Poast Plus** in **Poast Protected** sweet corn may be made until the onset of pollen shed. **DO NOT** apply **Poast Plus** after pollination occurs. A second application of **Poast Plus** in **Poast Protected** sweet corn

13/20
may be made 10 days or later following the first application.

Poast Plus may be applied in a tank mix with one of the following herbicides:

- atrazine
- **Basagran**
- **Frontier**
- **Guardsman**
- **Outlook**®
- **Laddok S-12**

Cotton

Processed meal may be fed to animals.

Poast Plus may be applied in a tank mix with one of the following herbicides: (including herbicides registered for use in cotton tolerant to glyphosate and bromoxynil):

- **Buctril**®
- glyphosate (e.g. **Roundup**®)
- **Staple**®

For best grass control, apply **Poast Plus** 3 days prior to **Staple**.

Flax

Not registered in California

Processed meal may be fed to animals.

Poast Plus may be applied in a tank mix with one of the following herbicides:

- **Buctril**
- MCPA
- **Bronate**®

Buctril, MCPA or **Bronate** herbicides applied with **Poast Plus** may cause leaf burn, retarded growth, and delayed maturity of the crop.

Tank Mixing Restrictions (partial list)

DO NOT delay spraying broadleaf weeds even though grassy weeds are not in the correct stage for treatment.

DO NOT add AMS or UAN solution to a tank mix of **Poast Plus** + **Buctril** or MCPA or **Bronate**.

Lentil

Not registered in California.

Mint

Poast Plus may be applied in a tank mix with one of the following herbicides:

- **Basagran**
- **Buctril**

Peanut

Processed meal may be fed to animals.

Poast Plus may be applied in a tank mix with one of the following herbicides.

- **Basagran**
- **Blazer**®
- **Storm**®
- 2,4-DB

Soybean

In California, the maximum rate per acre per application is 3.0 pints.

Only processed meal from seed or hay may be fed to animals.

14/20

Poast Plus® herbicide may be applied in a tank mix with one of the following herbicides (including uses in **RoundUp Ready®, Liberty Link®** and **STS®** varieties):

- **Basagran®**
- **Blazer®**
- **Classic®**
- **Cobra®**
- **FirstRate®**
- **Flexstar®**
- **Frontier®**
- **Galaxy®**
- glyphosate (e.g. **Roundup®**)
- **Liberty®**
- **Pursuit®**
- **Pursuit® DG**
- **Pursuit® W**
- **Pursuit® W DG**
- **Raptor™**
- **Reflex®**
- **Reliance® STS**
- **Resource®**
- **Stellar®**
- **Storm®**
- **Synchrony® STS**
- **Touchdown®**
- 2,4-D (LVE)*

*For use as preplant burndown only.

Tank Mix Specific Restrictions

Tank mixes of **Poast Plus** with **Basagran + Blazer, Galaxy** or **Storm** herbicides are not for use in California.

DO NOT use MSO with any tank mix combination except with **Basagran, Pursuit** or **Raptor** herbicides.

Sugar Beets

Processed pulp and molasses may be fed to animals.

Poast Plus may be applied in a tank mix with one of the following herbicides:

- **Betamix®**
- **Stinger®**
- **Betanex®**
- **UpBeet®**

Poast Plus may be tank mixed with other postemergence herbicides that are registered for use on sugar beet varieties tolerant to those herbicides.

Tank Mixing Restrictions

Not for use in California.

The use of UAN solution or AMS with a **Poast Plus + Betamix/Betanex** herbicides tank mix is not recommended.

DO NOT use **Poast Plus + Betamix/Betanex** if grasses to be controlled include rhizome Johnsongrass, quackgrass, Bermudagrass, wirestem muhly, volunteer corn, shattercane, red rice, or itchgrass.

Sunflower

Commercially released varieties of sunflower are tolerant to **Poast Plus** at all stages of growth; however, leaf speckling has been occasionally observed on sunflowers with no corresponding reduction in vigor or growth. **Poast Plus** is not recommended for use on sunflower inbred lines grown for seed because crop safety of these lines has not been adequately established.

Processed meal and soapstock may be fed to animals.

Tobacco

Not registered in California.

1st application: Make the first application to plantbed seedlings up to 4 weeks prior to transplanting to the field. Maximum application rate: 1.5 pints/acre.

2nd application: The second application may follow up to 3 weeks after transplanting. Maximum application rate: 2.25 pints/acre

3rd application: The third application may be made up to 7 weeks after transplanting. Maximum application rate: 2.25 pints/acre.

Poast Plus may be applied at the seedbed stage of growth.

Forage Crops

Alfalfa, Birdsfoot Trefoil, Clover, Sainfoin

Poast Plus may be applied to seedling or established alfalfa and clover grown for hay, silage, green chop, direct grazing, or for seed.

Mowing: The best control of annual grasses can be achieved by applying **Poast Plus** before grass weeds are mowed. Once a grass is mowed it becomes tougher to control, as much of the leaf surface may be removed, putting the grass under stress. In areas without a killing frost, some annuals can overwinter after having been mowed a number of times. These grasses can form large crowns and contain many viable buds. A large crown, even if it is an annual grass, may require repeated applications of **Poast Plus** for partial or complete control.

Tank Mixing in Alfalfa, Birdsfoot Trefoil and Sainfoin Only

Poast Plus may be applied in a tank mix with:

- 2,4-DB

Tank Mix Specific Restrictions

DO NOT add UAN solution or AMS to a tank mix of **Poast Plus + 2,4-DB**.

DO NOT use **Poast Plus + 2,4-DB** in the High and Rolling Plains of Texas, western Oklahoma, western Kansas, and eastern New Mexico.

Irrigated Alfalfa, Clover, Birdsfoot Trefoil, and Sainfoin:

Irrigation practices can be very critical to the successful use of **Poast Plus** and may be necessary to start grass weeds growing again. Generally, applications 2-4 days after an irrigation are most effective because:

- grasses resume active growth,
- grasses have less chance to grow too large,
- by waiting later, the clover or alfalfa begins to canopy and interferes with spray coverage.

15/20

Irrigation shortly after application (2 days) can be effective, but more consistent grass control is obtained when the irrigation is made before the application.

Annual Grass Control

Apply **Poast Plus**® herbicide at the grass sizes and rates indicated in **Tables 1** and **3**. If grass has been cut, apply **Poast Plus** after the regrowth reaches the minimum height (so there will be enough leaf area for absorption) and before it exceeds the maximum height indicated.

Apply before the clover or alfalfa canopies cover the grasses and interfere with the spray coverage. Also, applications after a clover or alfalfa cutting may need to be timed to follow an irrigation or rainfall which will allow the grasses to regrow to a treatable size.

Some annual grasses are spring- and summer-germinating plants, while others are fall-germinating plants, and the time they are actively growing and most susceptible to **Poast Plus** may vary from area to area. Also, some annuals germinate over a long time, and because control of small grasses is desired, applications after each weed flush may be needed. As a general guideline, spray spring- and summer-germinating grasses as early in the season as possible. The optimum application timing may occur very early in the spring after initial greenup. Spray fall-germinating weeds in the fall soon after they begin growing but before any killing frosts. Late fall applications may be less effective due to environmental changes, such as frosts or the onset of flowering.

Perennial Grass Control

Poast Plus effectively controls or suppresses perennial grasses, such as Bermudagrass, Johnsongrass, quackgrass, wirestem muhly, and perennial ryegrass (see **Table 2**). However, their growth characteristics are such that they are more difficult to control than annual grasses, especially in a perennial crop such as established alfalfa or clover. A program of repeated applications is usually necessary for best results.

The most economical way of controlling perennial grasses is to do so in the year of stand establishment before rhizomes or stolons become large and difficult to kill. The field should be disked before seeding to thoroughly fragment rhizomes or stolons.

In summer and fall seedings, cool season grasses (quackgrass, wirestem muhly, and perennial ryegrass) can become very competitive under cool fall conditions. Fall applications of **Poast Plus** will reduce late season grass growth and limit the ability of grasses to accumulate nutrient reserves in roots and rhizomes.

In established stands, it is important to begin applying in the spring when conditions favor active growth and before storage tissues have increased their nutrient reserves. Additional applications should be made on any grass regrowth in later cuttings.

Interseeded Oats

Oats interseeded with clover, alfalfa, birdsfoot trefoil, and sainfoin may be killed by applying **Poast Plus**. Their removal allows the seedling crops to grow with less competition. This application should be made before the interseeded oats reach the boot stage or later to be most effective.

Fruit and Nut Crops

Blueberry

Not registered in California.

Caneberries

Aircraft use not registered in California.

Citrus

Pulp and waste may be fed to livestock.

Cranberry

Not registered in California.

Grape

Pomace and raisin waste may be fed to animals.

Pistachio

Poast Plus should only be applied as a directed spray to the grove floor for bearing pistachio trees.

Pome Fruits

Pressed or processed apple waste may be fed to animals.

Strawberry

Not for aircraft application in California.

Tree Nuts

Poast Plus may be used for grass control and suppression in bearing or nonbearing tree nuts. Tree nuts are very tolerant to **Poast Plus**; **Poast Plus** may be applied over the top of small, nonbearing trees or as a directed spray on larger trees.

DO NOT apply **Poast Plus** with another pesticide whose label cautions against use with oil adjuvants.

In almond, only almond hulls may be fed to animals.

16/20

Nonbearing Fruit and Nut Crops

For nonbearing areas, always add 2 pints of oil concentrate per acre.

The nonbearing crops that **Poast Plus**® herbicide may be applied to are:

- Avocado
- Date
- Fig
- Olive
- Plum
- Pomegranate
- Prune

To minimize the potential for tree injury, direct the spray away from the leaves as much as possible.

Set Aside Conservation Reserve Land, Fallow Acreage

Broadleaf Cover Crops: The growth of broadleaf cover crops such as alfalfa, clover, lespedeza, trefoils, and vetches will not be affected by **Poast Plus**.

Grass Cover Crops: Most seeded grass crops such as oats, Sudangrass, tall fescue, orchardgrass, bromegrasses, ryegrass, or timothy will be injured or killed by **Poast Plus**; therefore, **DO NOT** use **Poast Plus** if injury to these grass cover crops is undesirable.

Seeded grass cover crops may be injured or killed.

Restrictions and Limitations (partial list): **DO NOT** harvest or graze cover crops other than alfalfa, clover, birdsfoot trefoil, or sainfoin treated with **Poast Plus**.

This use is applicable only for the Midwest, South, and northeast areas or east of the Rocky Mountains (see maps in **Table 1**).

For alfalfa cover crops, **DO NOT** apply **Poast Plus** within 7 days of grazing, feeding, or cutting for (undried) forage, or within 14 days of cutting alfalfa for (dry) hay.

For alfalfa cover crops, **DO NOT** apply more than a total of 9.75 pints of **Poast Plus** per acre in one season.

Poast Plus may be applied in a tank mix with one of the following herbicides:

- Clarity®
- Marksman®
- glyphosate (e.g. Roundup®)
- 2,4-D

Interseeded Cover Crops

Poast Plus Activity on the Cover Crop

Grass cover crops controlled or suppressed by this use include wheat, oats, and barley, or any grass crop for which **Poast Plus** is labeled. **Poast Plus** will selectively control grass cover crops in seedling nongrass or broadleaf field, forage, or vegetable crops without injury. In addition, **Poast Plus** will control any annual grasses that have emerged since planting. The slow-dying grass can

provide a protective mulch for the primary crop seedlings for up to 3 weeks after applying **Poast Plus**.

Apply **Poast Plus** to cereals that are 3-4" in height (before tillering). **DO NOT** allow cereals to exceed this height as excessive competition and lack of control may occur.

Noncrop Areas

Deciduous Trees, Nonfood Crop Areas, Fallow Land

Deciduous trees, Nonfood crop areas, Fallow land:

Poast Plus may be used in noncrop areas including rights-of-ways, roadsides and other paved areas, along fences and hedgerows, public buildings, recreation areas, industrial sites, storage yards, airports, electric transformer stations, pipeline pumping stations, sewage disposal areas, on potting and top soils, uncultivated agricultural areas, and general indoor or outdoor sites.

Poast Plus is not recommended for use on red sprangle-top in California, Arizona or western New Mexico.

Notice to user: Due to variability within species and in application techniques, neither the manufacturer nor the seller has determined whether or not **Poast Plus** can be safely used on all varieties and species of nonbearing food crops and other nonfood crops under all conditions. Therefore, determine if **Poast Plus** can be used safely before broad use in the following manner:

On a small test area, apply the recommended rate of **Poast Plus** on nonbearing or nonfood crop species or varieties under the conditions expected to be encountered. Any adverse conditions should be visible within 7 days.

FINE FESCUE GROWN FOR TURF SEED:

The fine fescues tolerant to **Poast Plus** at all stages of growth are Creeping Red (*Festuca rubra*), Chewings (*Festuca nigrescens*), and Hard Fescue (*Festuca ovina*). Tall Fescue (*Festuca arundinacea*) is **NOT** tolerant to **Poast Plus**. Avoid all direct or indirect contact with any desired grass plant.

For control of annual ryegrass, downy brome, German velvetgrass, and Colonial and Highland bentgrasses, apply **Poast Plus** when the fine fescue is semi-dormant (generally November 1 through March 15). Application of **Poast Plus** at other times of the year will generally result in reduced control of these problem grass weeds. See recommendations for timing and rates in the **Application Rate Table** below for more specific instructions.

Restrictions and Limitations

- Apply **Poast Plus** only to fine fescue varieties classified as Creeping Red, Chewings, or Hard Fescue.
- **DO NOT** apply **Poast Plus** to tall fescue (*Festuca arundinacea*) because injury will occur.
- **DO NOT** graze treated field and **DO NOT** feed treated fescue screenings or hay to livestock.

17/20

- **DO NOT** apply if rainfall is expected within one hour following application, as grass control will be unsatisfactory.
- **Poast Plus® herbicide** does **NOT** control annual bluegrass (*Poa annua*) or rattail fescue (*Festuca myuros*).
- Make no more than 2 applications of **Poast Plus** to fine fescue grown for turf seed in one use season.
- **DO NOT** apply **Poast Plus** to fine fescue by air.

Ground Equipment

Thorough spray coverage of foliage is essential. Use a minimum of 10 gallons of water per acre with a minimum of 40 psi at the nozzle. Increase water volume to 20 gallons per acre and increase pressure to a minimum of 60 psi if grass foliage is dense. Use standard high-pressure pesticide hollow cone or flat fan nozzles. **DO NOT** use flood or whirl chamber nozzles. **DO NOT** use selective application equipment such as recirculating sprayers, wiper applicators, or shielded applicators. Refer to the main **Poast Plus** label for additional information pertaining to ground application methods.

Addition of Oil Concentrate

Always add a nonphytotoxic oil concentrate to the spray solution at 2 pints/acre. Refer to the main **Poast Plus** label for specific instructions pertaining to use of oil concentrates.

Application Rate Table for Use of Poast Plus in Fine Fescue Grown for Turf Seed

Grass Species	Application Time (when fine fescue is semi-dormant)	Rate of Poast Plus per Acre*
Annual Grasses		
Annual Ryegrass (<i>Lolium multiflorum</i>)	4-8"	2¼ pints
Downy Brome (<i>Bromus tectorum</i> , also called cheatgrass)	2-6"	3¾ pints
Late fall applications after maximum germination occurs produce the best results.		
Perennial Grasses		
German Velvetgrass (<i>Holcus mollis</i>)	2-4"	3 - 3¾ pints
Colonial and Highland Bentgrasses (<i>Agrostis tenuis</i>)	2-4"	2¼ - 3¾ pints
The higher rate of Poast Plus is recommended for use on well-established grass weeds.		
* If regrowth occurs or new plants emerge, make a second application at the same rate and time.		

TALL FESCUE GROWTH SUPPRESSION:

Apply **Poast Plus** to actively growing tall fescue after it has 4-6 inches of new growth, before the emergence of seedheads and before conifer bud break. Applications made from July 1 to mid-August may be less effective, especially if day temperatures reach 90° F. Tall fescue must be 1-year old before the first application of **Poast Plus**.

Adequate coverage of the leaf surface is necessary for absorption of this herbicide. Thus, for optimum control, **DO NOT** mow tall fescue turf for 30 days before or 14 days after applying **Poast Plus**.

Rate: Apply 1.5-1.875 pints of **Poast Plus** per acre. For greater fescue suppression, up to 2.5 pints of **Poast Plus** per acre can be used. Because of environmental differences at application and growth differences of tall fescue, control may exceed or fall short of that desired. Begin treating crops with **Poast Plus** at the minimum recommended rate and adjust rates as local conditions and experience dictate. Additional applications may be made if extended growth suppression is desired.

ORCHARD FLOOR MIDDLES:

Growth Management In Orchard Floor Middles

Poast Plus and 2,4-D dimethylamine can be used in a tank mix for growth management in orchard floor middles to reduce the number of mechanical mowings needed during a season. **Poast Plus** and 2,4-D dimethylamine can be safely applied for growth management in the following cool season grasses and mixtures: Kentucky bluegrass, perennial ryegrass, and tall fescue. Some degree of discoloration of the turf may occur. However, the turf will regrow and green up as effects of the treatment wear off. Make one application per season from the following options:

- **Poast Plus** and 2,4-D dimethylamine can be applied during the spring or summer when growth management is desired. **DO NOT** apply during bloom or within 3 days of a mowing.
- An optimal timing for application is after sod green up in the spring (before any mowing) or 3 days after the initial mowing of the season is made.
- A prebloom treatment is recommended as any broadleaf weeds such as dandelions can be controlled before they hamper fruit pollination. This treatment will provide 5-8 weeks of growth management depending on the sod makeup (e.g. grass species, amount of broadleaf weeds present, etc.), environmental conditions and the desired maintenance height of the middles.

See section III. **Additives** and **Mixing Order** for details.

Tank Mix Specific Restrictions

Make no more than 1 application of this tank mix per growing season.

DO NOT apply if rainfall or irrigation is expected within 6 hours after application as growth management effects will probably be unsatisfactory.

DO NOT apply to a grass sod that is less than 2 years old.

DO NOT apply to newly established orchards. Trees must be at least 1-year old and in vigorous condition.

DO NOT apply this tank mix within 14 days of harvest of apples and pears.

DO NOT apply this tank mix within one year of harvest of nonbearing plums. Not registered for use in California.

Vegetable Crops

Allow a minimum of 14 days between sequential applications.

Always add oil concentrate at 2 pints per acre. However, when the temperature exceeds 90°F and the relative humidity is 60% or greater, or any time the temperature exceeds 100° F regardless of the humidity, **Poast Plus® herbicide** plus adjuvants should be used with caution due to potential leaf injury.

Brassica Vegetables

Mustard greens may be harvested 14 days after the last application. All other brassica vegetable crops may be harvested no sooner than 30 days after the last application.

Curcubit Vegetables

Cantaloupe may be harvested 3 days after the last application. All other curcubit vegetable crops may be harvested no sooner than 14 days after the last application.

Fruiting Vegetables

Peppers may be harvested 7 days after the last application. All other fruiting vegetable crops may be harvested no sooner than 20 days after the last application. Tomato waste may be fed to animals.

Head and Petiole Vegetables

In Florida, celery may be harvested 14 days after the last application. All other head and petiole vegetable crops may be harvested no sooner than 30 days after the last application.

Horseradish

Not for use in California.

Potato and Tomato

In case of heavy infestations of quackgrass in potato, use 3¾ pints of **Poast Plus** per acre followed by 2¼ pints per acre sequentially if needed.

Potato and tomato wastes may be fed to animals.

Tank mixes: Poast Plus may be applied in a tank mix with other herbicides, such as metribuzin-containing products, in potato and tomato. The most restrictive labeling applies to tank mixes. Refer to section **IV. General Tank Mixing Information** for further instructions.

Specific Restrictions for Tank Mixing with Metribuzin-containing Products

This tank mix not applicable to California.

DO NOT apply this tank mix to sweet potato or yams. Apply only if there have been at least 3 successive days of sunny weather before application or crop injury may occur.

DO NOT add UAN solution or AMS to a **Poast Plus** + metribuzin tank mix.

DO NOT use this tank mix if grasses to be controlled include rhizome Johnsongrass, quackgrass, Bermudagrass, wirestem muhly, volunteer corn or cereal, shattercane, red rice, or itchgrass.

Apply only to russeted or white-skinned varieties of potato that are not early maturing.

DO NOT apply this tank mix within 60 days of potato harvest.

DO NOT treat transplanted tomatoes within 14 days of transplanting. Tomatoes must have recovered from transplant shock and new growth must be evident.

DO NOT treat seeded tomatoes until plants have reached the 5-6 leaf stage.

Sweet Potato

Eastern US includes AL, FL, GA, LA, MS, NC, SC, TN, TX and VA.

Western US includes AZ, CA, ID, NV, OR and WA.

Rhubarb

Rhubarb grown only in IL, IN, MI, MN, and WI may be harvested up to **15 day PHI**.

Aircraft application not registered.

18/20

19/20

Weeds listed in this label:	
Common Name	Scientific Name
Barnyardgrass (Watergrass)	<i>Echinochloa crus-galli</i>
Bermudagrass (Wiregrass)	<i>Cynodon dactylon</i>
Crabgrass, Large	<i>Digitaria sanguinalis</i>
, Smooth	<i>Digitaria ischaemum</i>
Cupgrass, Southwestern	<i>Eriochloa gracillis</i>
, Woolly	<i>Eriochloa villosa</i>
Fescue, Tall	<i>Festuca arundinacea</i>
Foxtail, Giant (Pigeongrass)	<i>Setaria faberi</i>
, Green	<i>Setaria viridis</i>
, Yellow	<i>Setaria glauca</i>
Goosegrass	<i>Eleusine indica</i>
Itchgrass	<i>Rottboellia exaltata</i>
Johnsongrass	<i>Sorghum halepense</i>
Junglerice	<i>Echinochloa colonum</i>
Lovegrass	<i>Eragrostis</i> sp.
Millet, Wild Proso	<i>Panicum miliaceum</i>
Muhly, Wirestem	<i>Muhlenbergia frondosa</i>
Oats, Tame	<i>Avena sativa</i>
, Wild	<i>Avena fatua</i>
Orchardgrass	<i>Dactylis glomerata</i>
Panicum, Browntop	<i>Panicum fasciculatu</i>
, Fall	<i>Panicum dichotomiflorum</i>
, Texas	<i>Panicum texanum</i>
Quackgrass	<i>Agropyron repens</i>
Red Rice	<i>Oryza sativa</i>
Ryegrass, Annual	<i>Lolium multiflorum</i>
, Perennial	<i>Lolium perenne</i>
Sandbur, Field	<i>Cenchrus incertus</i>
Shattercane/Wildcane	<i>Sorghum bicolor</i>
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>
Sprangletop, Red	<i>Leptochloa filiformis</i>
Stinkgrass	<i>Eragrostis cilianensis</i>
Volunteer Barley	<i>Hordeum vulgare</i>
Corn	<i>Zea mays</i>
Oats	<i>Avena sativa</i>
Rye	<i>Secale Cereale</i>
Wheat	<i>Triticum aestivum</i>
Witchgrass	<i>Panicum capillare</i>

Crops

This product may be used on the following crops:

Alfalfa	Juneberry
Apricot	Leafy Vegetables
Artichoke	Lentil
Asparagus	Lingonberry
Avocado*	Mint
Beans	Nectarine
Beet, Garden	Olive*
Birdsfoot Trefoil	Peach
Blueberry	Peanut
Brassica Crops	Peas, Dry & Succulent
Bulb Vegetables	Pistachio
Caneberries	Plum*
Canola/Crambe	Pome Fruits
Carrot	Pomegranate*
Cherry	Potato
Citrus	Prune*
Clover	Root & Tuberous
Corn (Poast Protected™)	Vegetables
field and sweet	Safflower
Cotton	Sainfoin
Cranberry	Salal
Cucurbits	Soybean
Date*	Strawberry
Fescue, Tall	Sugar Beet
Fig*	Sunflower
Flax	Tobacco
Fruiting Vegetables	Tree Nuts
Grape	Other Nonbearing &
Head & Petiole Vegetables	Nonfood Areas
Horseradish	

*** Nonbearing crop only.**

Look inside for complete restrictions and limitations and application instructions.

20
/20

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Basagran, Clarity, Dash, Frontier, Galaxy, Guardsman, Marksman, Outlook, Poast Plus, Pursuit, Raptor, and Storm are registered trademarks of BASF.

Poast Protected is a trademark of BASF.

Betamix, Betanex, Liberty and **LibertyLink** are registered trademarks of Bayer Crop Science GmbH.

Bronate, Buctril and **Sencor** are registered trademarks of Bayer AG.

Classic, Reliance, Staple, STS, Synchrony and **UpBeet** are registered trademarks of E.I. duPont de Nemours and Company.

Dual, Dual II, Flexstar, Reflex and **Touchdown** are registered trademarks of Syngenta Crop Protection, Inc.

FirstRate, Stinger are registered trademarks of Dow AgroSciences LLC.

Harness, Roundup and **RoundUp Ready** are registered trademarks of Monsanto Company.

Cobra, Resource and **Stellar** are registered trademarks of Valent U.S.A. Corporation.

Sundance is a registered trademark of Rosen's, Inc.

Laddok is a registered trademark of Micro Flo Company.

Blazer is a registered trademark of United Phosphorus, Inc.

© 2006 BASF Corporation
All rights reserved.

007969-00088.20070420.NVA 2006-04-026-0270
Supersedes: NVA 2005-04-026-0229

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709


The Chemical Company