



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460**

FEB 24 1994

**ROBERT RHODE
BASF CORP.
AGRICULTURAL PRODUCTS
BOX 13528
RESEARCH TRIANGLE PARK, NC 27709**

**OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES**

**Subject: Label Amendment Submission of 10/13/93 Response to PR Notice 93-7
EPA Reg. No. 7969-97
PIX CONCENTRATE PLANT REGULATOR**

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- **BEFORE** selling or distributing any product bearing the final printed labeling
- AND**
- **WITHIN** one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on Paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

Robert Rohde
BASF CORP
AGRICULTURAL PRODUCTS
BOX 13528
RESEARCH TRIANGLE PARK NC 27709

Comment for: EPA Reg Nr.7969-97
PIX CONCENTRATE PLANT REGULATOR

The following specific comments pertain to your WPS
labeling submission concerning the product
cited above:

The "Precautionary Statements" and the "Hazards to Humans and Domestic
Animals" headings are missing.

BASF

ACCEPTED
with COMMENTS 40 13
in EPA Letter Dated 5

FEB 24 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

7969-97

October 13, 1993

Pix[®] Concentrate

plant regulator

Active Ingredient*

Mepiquat chloride:

N,N-dimethylpiperidinium chloride	22.5%
Inert ingredients	77.5%
TOTAL	100.0%

*Equivalent to 2.01 pounds active ingredient per gallon.

EPA Reg. No. 7969-97

Agricultural Use Requirements
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

KEEP OUT OF REACH OF CHILDREN CAUTION

Harmful if swallowed or absorbed through the skin. Causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

Practical Treatment

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

Environmental Hazards

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment waste waters.

Net contents 1 gallon

BASF Corporation

P.O. Box 13528 Research Triangle Park, North Carolina 27709-3528

BEST AVAILABLE COPY

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- waterproof gloves and shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions, for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

BEST AVAILABLE COPY

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will create a mist or spray that is inhaled or absorbed through the skin or clothing. Only pesticides are to be applied in the treated area during application.

For any requirements specific to your State or other consulting agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS) at 29 CFR part 170. This Standard contains additional information for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions concerning the restrictions on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, waterproof gloves, and shoes plus socks.

BEST AVAILABLE COPY

In Case of Emergency

In case of large-scale spillage regarding this product, call:

CHEMTREC. 800-424-9300
BASF CORPORATION. . . . 800-832-HELP

In case of medical emergency regarding this product, call:

1. Your local doctor for immediate treatment.
2. Your local poison control center (hospital).
3. BASF CORPORATION. . 800-832-HELP

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple-rinse container (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

Pix® Concentrate plant regulator is a foliar applied plant regulator which modifies the cotton plant in several beneficial ways. It is the only such compound which allows the grower to manage the cotton plant for short-season production leading to reduced risk of yield and quality loss due to delayed and prolonged harvest. The use of Pix Concentrate will also result in several or all of the following: height reduction and more open canopy; better early boll retention and/or larger bolls; less boll rot; improved defoliation; reduced trash and lower ginning costs; better harvest efficiency; and darker green leaf color. Most of these effects often favorably influence the yield potential of the cotton plant.

On both short-staple and Pima cotton, the grower has the option of single, dual, or up to four low-rate multiple applications, which greatly facilitates his management flexibility. One of the major factors affecting cotton growth and development is soil moisture. If significant drought stress occurs when Pix Concentrate is applied, or occurs after a full rate (1.40 -2.80 fl oz/A) is applied, results may not be optimal. The low-rate multiple option gives the producer the ability to discontinue usage of Pix Concentrate if any significant stresses occur after an earlier application; in such a case, the total quantity of Pix Concentrate used over a season may be reduced. If stress is relieved, the grower has the option of continuing treatments with Pix Concentrate. In addition, the rate range indicated allows the grower to tailor his usage of Pix Concentrate to the degree of vegetative vigor in a given field. In areas where insecticides/miticides or foliar fertilizers are frequently applied, the timings are such that tank-mixing is often possible. (See Restrictions and limitations) Fields should be carefully scouted and Pix Concentrate should not be applied if plants are under severe stress from weather factors, mite, insect or nematode damage, disease stress, herbicide injury, or fertility stress. In the absence of these stresses, up to four low-rate multiple applications could be made each season, generally using 0.35-0.70 fl oz/A each time. After the first application (at matchhead square in the absence of stress), the rate and timing of subsequent application(s) will depend on vegetative vigor; under good growing conditions, additional treatments (0.35-0.70 fl oz/A) should be made at 7-14 day intervals. However, if new growth at any time is excessive, higher rates of Pix Concentrate (0.7-2.1 fl oz/A) can be used. If significant loss of squares and/or young bolls has occurred earlier due to insect pressure or other stresses, but now these stresses have been alleviated, the need for Pix Concentrate is increased - excess vegetative growth is likely because of poor fruit load. Total use per season should not exceed 4.2 fl oz/A.

Spray Volume

Water as Diluent

Ground and Aerial Application:

When applying Pix Concentrate with ground equipment, in all states but California, use a minimum of two gallons of water per acre; in California, use a minimum of five gallons per acre. For aerial application in all states but California, use a minimum of two gallons of water per acre; in California, use a minimum of five gallons per acre. Regardless of method or gallonage of application, thorough coverage of the cotton foliage is required.

Oil as Diluent

Ultra Low Volume (ULV) Aerial Application:

Only permitted in the following states: AL, AR, FL, GA, LA, MO, MS, NC, OK, SC, TN, and TX. A non-phytotoxic oil concentrate should be used which must contain either a petroleum or vegetable oil base and must meet the following criteria: (1) be non-phytotoxic; (2) contain only EPA-exempt ingredients; and (3) be successful in local experience. The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers which provide good mixing quality. If the oil does not contain an emulsifier, one must be added during mixing at a volume equal to 3% of the final volume of the mixing tank. Do not apply Pix Concentrate ULV without using emulsifiers. It is recommended that vegetable oil concentrates be highly refined.

Mixing Directions

Oil as Diluent:

While agitating, pour approximately one-half of the required volume of oil (total final spray solution at least 2 pts/A) into the mixing tank, followed by the emulsifier (if the oil does not already contain one) at approximately 3% of the final spray tank volume, and then simultaneously pour in the Pix Concentrate and the remainder of the oil.

Only moderate agitation should be employed while mixing and transporting.

Rain-safe Period/Compatibility

Rain-safe Period

If rain is expected within 8 hours, use of a high quality, EPA-exempt surfactant is recommended, which will result in a rain-safe period of 4 hours.

Compatibility

Pix Concentrate has an aqueous base, and as such is compatible with most insecticides and miticides. If in doubt, test compatibility by adding a teaspoon of the insecticide or miticide to a pint of ready-to-use spray solution of **Pix Concentrate**. You may combine **Pix Concentrate** with foliar fertilizers if your prior experience has shown the combination to be compatible and non-injurious under your conditions.

Restrictions and Limitations

Significant insect or mite damage after application(s) of **Pix Concentrate** is likely to eliminate or greatly reduce the benefits derived from **Pix Concentrate**.

Do not apply a single application of 1.4-2.8 fl oz **Pix Concentrate** to cotton that is drought stressed; i.e., stressed due to lack of soil moisture. If using the low-rate multiple option, discontinue use until the moisture stress is alleviated.

Do not apply more than 4.2 fl oz of **Pix Concentrate** per acre per season.

Do not apply **Pix Concentrate** within 30 days of harvest.

Do not graze or feed cotton forage to livestock within 30 days of application, or after applying **Pix Concentrate** in oil as a ULV application by air.

Do not tank mix with other products other than those mentioned under **Compatibility**.

Do not apply this product through any type of irrigation system.

Time and Rate of Application: Short-staple and Long-staple (Pima) Cotton

The times and rates of application have been carefully researched and the Directions for use should be observed as specified below.

Single or Dual Applications

Geographic Area	Time of Application	Rate per Acre
AL, AR, AZ, CA, FL, GA, LA, MO, MS, NM, NC, SC, TN, VA	First application: Apply when cotton is actively growing and is between 20" and 30" tall, provided cotton is not more than 7 days beyond early bloom stage (5-6 blooms per 25 row feet). If cotton is 24" tall and has no blooms, apply Pix® Concentrate. Use 1.4 fl oz per acre on cotton where excessive vegetative growth is not likely to be a problem, and 2.8 fl oz per acre in areas tending to have excessive vegetative growth. See Restrictions and limitations.	1.4 to 2.8 fl oz
	Second application for control of excessive vegetative growth: If the cotton field has a history of vigorous growth, and/or conditions after the first application of Pix Concentrate favor vigorous growth, apply a second application 2 to 4 weeks after the first application.	1.4 fl oz
OK, TX (except Rio Grande Valley)	Areas where excessive vegetative growth is not a problem First application: Apply when cotton is in early bloom stage (5-6 blooms per 25 row feet) and actively growing. If no blooms are present and the cotton is 20" tall and actively growing, apply Pix Concentrate. See Restrictions and limitations.	1.4 fl oz
	Second application: If conditions after the first application of Pix Concentrate favor vigorous growth, apply a second application 2-4 weeks after the first application.	1.4 fl oz
OK, TX	Areas where excessive vegetative growth is a problem First application: Apply when cotton is actively growing and between 20" and 30" tall, provided cotton is not more than 7 days beyond early bloom stage (5-6 blooms per 25 row feet). If cotton is 24" tall and has no blooms, apply Pix Concentrate.	2.8 fl oz
	Second application for control of excessive vegetative growth: If cotton field has a history of vigorous growth, and/or conditions after the first application of Pix Concentrate favor vigorous growth, apply a second application 2-4 weeks after the first application.	1.4 fl oz

Refer to Conversion Chart on Page 4.

Low-rate Multiple Applications

Geographic Area (States)	Time of Application	Average Historical Use Rate = ½ pt/A of Pix Equivalent	Average Historical Use Rate = 1 pt/A (or fully irrigated)
AL, AR, AZ, CA, FL, GA, LA, MO, MS, NC, NM, OK, SC, TN, TX, VA	First application: Optimal results will be achieved when plants are in the matchhead square* stage of growth.	0.35 fl. oz.	0.7 fl. oz.
	Second application: 7-14 days later, or when regrowth occurs.	0.35-0.7 fl. oz.*	0.7-1.4 fl. oz.*
	Third application: 7-14 days later, or when regrowth occurs.	0.35-0.7 fl. oz.*	0.7-1.4 fl. oz.*
	Fourth application: 7-14 days later, or when regrowth occurs.	0.35-0.7 fl. oz.*	0.7-1.4 fl. oz.*

*Use higher rate if previous application was not made or if growing conditions are conducive to vigorous growth.

*matchhead square is when the first square of a typical cotton plant is 1/8-1/4" in diameter. The first application should go on when 50% of the plants have one or more matchhead squares.

Do not apply more than 4.2 fl. oz. per acre per season.

Conversion Chart: Pix Concentrate and Pix*	
fl. oz. Pix Concentrate	fl. oz. Pix
0.35	2 (¼ pt)
0.70	4 (¼ pt)
1.40	8 (½ pt)
2.80	16 (1 pt)

*1 oz of Pix Concentrate = 5.7 oz of Pix

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of **BASF CORPORATION ("BASF")** or the Seller. All such risks shall be assumed by the Buyer. BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for use, subject to the inherent risks referred to above. **BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.** BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Additional Information

For additional information concerning this label and use of Pix Concentrate call BASF's "COMMSERV" at 1-800-367-8896.

*Pix is a registered trademark of BASF AG.
©199 BASF Corporation.*