
BASF

Not intended for use in California





Postemergence Flowable Herbicide

For postemergence use on residential and nonresidential turf.

Active Ingredients:

Sodiurn salt of bentazon* [3-(isopropyl)-1H-2,1,3-benzothiadiazin	
-4(3H)-one,2,2-dioxide]	19.1%
Atrazine* (2-chloro-4-ethylamino-6-isopropylamino-s-triazine)	17.5%
Inert ingredients	63.4%
Total	
*Equivalent to 1.66 pounds per gallon each of bentazon and atrazine.	
EPA Reg. No. 7969-103	

KEEP OUT OF REACH OF CHILDREN. DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Corrosive. Causes eye damage and skin imitation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed.

Statement of practical treatment

If in eyes, immediately flush eyes with plenty of water. Get medical attention. If on skin, immediately flush skin with plenty of water. Get medical attention if imitation persists. May cause allergic skin response.

See inside booklet for complete **Precautionary Statements**, **Directions For Use**, and **Conditions of Sale and Warranty**.


Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the **Directions For Use** section for information about this standard.

Shake well before using.

Net Contents: 1 gallon

BASE Corporation
P.O. Box 13528, Research Triangle Park, NC, 27709-3528



Specimen Label

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swalfowed.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- · Long-sleeved shirt and long pants
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eyewear
 Discard clothing and other
 absorbent materials that have been
 drenched or heavily contaminated
 with this product's concentrate. Do
 not reuse them, Follow manufacturer's instructions for clearing/maintaining PPE. If no such instructions
 or washables, use detergent and
 not water. Keep and wash PPE
 separately from other laundry.

Engineering Controls Statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations Users should:

- Wash hands before eating, drinking, chewing gum, usir:g tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Atrazine, which is present in this product, can travel (seep or leach) through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply Prompt^a herbicide to sand and loarny sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further infor-

mation on the type of soil in your area and the location of groundwater.

This pesticide is toxic to aquatic invertebrates. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with the labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eyewear

Non-agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter treated areas without protective clothing until sprays have dried.

, ک

General Information

Prompt^a herbicide is intended for the postemergence control of a broad spectrum of broadleaf weeds in turf. Prompt does not control grasses. **Prompt** is effective mainly through contact action; therefore, weeds must be thoroughly covered with spray. Large leaf canopies shelter smaller weeds and prevent adequate spray coverage. The desired plants recommended for use in this labeling are tolerant to **Prompt** when used as directed. Very slight leaf speckling of desired plants may occur but these plants generally outgrow this condition.

Timing of Applications

Apply **Prompt** early postemergence when weeds are small and actively growing and before weeds reach the maximum size listed in the **Application Rate Table**. Early application to weeds produces the most beneficial effect on weed control, allows use of the lower rate (depending on weed species), and makes it easier to obtain thorough spray coverage. Delay in application which permits weeds to exceed the maximum size stated will result in inadequate control.

Water Volume and Spray Pressure

Apply recommended rates of Prompt as follows:

Ground Equipment
Use a minimum of 20 gallons of the water per broadcast acre and a suminimum of 40 psi pressure (measured at the boom, not at the plump or in the line). When weed foliage is dense use up to 50 gallons of water and up to 80 psi. Use standard, high-pressure pesticide hollow cone or flat fan nozzles for best weed coverage.

3 4 x

Special Information on Irrigation In irrigated areas, it may be necessary to irrigate prior to treatment with Prompt* herbicide to ensure active weed growth. Weeds growing under drought conditions usually are not satisfactorily controlled, so it will be necessary to irrigate these areas before application or wait until adequate rainfall has been received.

NON-FOOD USE — Turf

For use in Bermudagrass, St. Augustinegrass, zoysia and centipedegrass. Apply no earlier than 10 days after sprigging or plugging. On newly sprigged turfgrass and hybrid bermudagrass, temporary slowing of growth and yellowing may occur following application.

Special Directions for Other Weed Problems Canada Thistle

Apply 31/2 pints of **Prompt** per acre when plants are from 8-10 inches tall to the bud stage. Make a follow-up application 7-10 days later using the same rate of **Prompt**, or use 2 pints of **Basagran* herbicide** per acre.

Yellow Nutsedge

A single application of **Prompt** at 31/2 pints per acre will provide suppression of yellow nutsedge. For control of yellow nutsedge apply 31/2 pints of **Prompt** per acre when plants are 1 to 4 inches tall. Make a follow-up application 7-10 days later using the same rate of **Prompt** or 11/2 to 2 pints of **Basagran** per acre.

Field Bindweed

For suppression of field bindweed apply 31/2 pints of **Prompt** per acre when field bindweed vines are 8 to 10 inches long.

Restrictions and Limitations
Do not apply Prompt when desired

plants are under stress from prolonged cold, wet weather, poor fertility, or other factors.

Do not apply **Prompt** if desired plants show injury (leaf phytotoxicity and/or plant stunting) produced by any other prior pesticide applications, because this injury may be enhanced and/or prolonged. Do not apply **Prompt** during prolonged periods of drought or during unseasonably cold weather as unsatisfactory weed control may result.

Do not graze treated area or feed treated forage to livestock. Rainfall or overhead irrigation soon after application (within 8 hours) may nullify the effectiveness of **Prompt**.

Do not apply this product through any type of imgation system. Groundwater contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material. This product may not be mixed, loaded, or used within 50 feet of all wells including abandoned wells, drainage wells and sink holes.

Storage and Disposal

Store above 15° F.
Do not contaminate water, food or feed by storage or disposal.
Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Application Rate Table for Bermudagrass, St. Augustinegrass, Zoysia, and Centipedegrass

Application Rates for Weed Growth Stages**						
Weeds Controlled	2 pints per acre		2 ¹ /2 pints per acre		3 ¹ /2 pints per acre	
Annuals	Leaf Stage	Maximum Height (in)	Leaf Stage	Maximum Height (in)	Leaf Stage	Meximum Height (in
Annual Morningglory Beggarticks Bristly Starbur Cocklebur Common Lambsquarters Common Ragweed Corn Speedwell Cudweed Dayflower Glant Ragweed Hopclover Jimsonweed Ladysthumb Oldfield Toadflax Oxalis Pennsylvania Smartweed Prickly Sida or Teaweed Redroot Pigweed Small Hopclover Smallflower Morningglory Smooth Pigweed Spurred Anoda Tall Waterhemp Velvetleaf Venice Mallow Wild Buckwheat Wild Sunflower Perennials	2.4 2.6 	3 4	Up to 4 2-10** Up to 8 Up to 5 5 Up to 4 Up to 6 Up to 10 Up to 4 Up to 6 Up to 6 Up to 6 Up to 5 Up to 5	4 85432 4 49 9 6 46 254346	4-6 Up to 6 Up to 4 2·10°· 8·12 4·7 6 Up to 6 4-6 6-10 10-14 Up to 6 10-14 Up to 6 Up to 6 6-8 4-6 6-10 4-6 4-6 4-6 4-6	662 855546281283123 26 348 588
Canada Thistle Field Bindweed Yellow Nutsedge		S	iee Special Dire iee Special Dire iee Special Dire	ictions on page	- 3	

Triple-rinse container (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. In Case of Emergency In case of large-scale spillage regarding this product: avoid contact, isolate area and keep out animals and unprotected persons. Confine spill and call: CHEMTREC 800-424-9300 BASF Corporation 800-832- HELP

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment,
- Your local poison control center (hospital),
- BASE Corporation 800-832-HELP.

Appendix of Scientific Names

The following are scientific names for the weeds listed in this label. For specific recommendations on control of these weeds, refer to the **Application Rate Table.**

Broadleaf Weeds

Common Name	Scientific Name		
Beggarticks	Bidens frondosa		
Bristly Starbur	Acanthospermum hispidum		
Canada Thistle	Circium arvense		
Cocklebur	Xanthium strumarium		
Common Lambsquarters	Chenopodium album		
Common Ragweed	Ambrosia artemisiifolia		
Corn Speedwell	Veronic arvensis		
Cudweed	Gnaphalium spp.		
Dayflower	Commelina spp.		
Field Bindweed	Convolvulus arvensis		
Giant Ragweed	Ambrosia trifida		
Hopolover	Trifolium aureum		
Hopdover, Small	Trifolium dubium		
Jimsonweed	Datura stramonium		
Ladysthumb	Polygonum persica ia		
Morningglory, Annual	Ipomea spp.		
Oldfield Toadflax	Linaria canadensis		
Oxalis	Oxalis stricta		
Pennsylvania Smartweed	Polygonum pensylvanicum		
^{∴ つ} rickly Sida or Teaweed	Sida spinosa		
łedroot Pigweed	Amaranthus retroflexus		
Morningglory, Smallflower	Jacquemontia tamnifolia		
Smooth Pigweed	Amaranthus hybridis		
Spurred Anoda	Anoda cristata		
Tall Waterhemp	Amaranthus tuberculatus		
Velvetleaf	Abutiton theophrasti		
Venice Mallow	Hibiscus trionum		
Wild Buckwheat	Polygonum convolvulus		
Wild Mustard	Sinapis arvensis		
Wild Sunflower	Helianthus annua		
Yellow Nutsedge	Cyperus esculentus		

Conditions of Sale and Warranty The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended con sequences may result, because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRAN-TY OF FITNESS OR MER-CHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRAN-TY. IN NO CASE SHALL BASE OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Basagran is a registered trademark of BASF AG. Prompt is a registered trademark of BASF Corporation. The Specialty Products Logo is a trademark of BASF Corporation.

© 1994 BASF Corporation NVA 1294/PR 4200-0638

BASE CorporationP.O. Box 13528
Research Triangle Park, NC 27709





Postemergence Flowable Herbicide

For postemergence use on residential and nonresidential turf.

Active Ingredients: Sodium salt of bentazon* [3-(isopropyl)-1H-2,1,3-benzothiadiazin

Inert ingredients 63.4%

*Equivalent to 1.66 pounds per gallon each of bentazon and atrazine.

EPA Reg. No. 7969-103

EPA Est. No. 11773-IA-1

KEEP OUT OF REACH OF CHILDREN.

DANGER/PELIGRO

- il usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you o not understand the label, find someone to read it to you in detail.)

corrosive. Causes eye damage and skin imtation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Statement of practical treatment

If in eyes, immediately flush eyes with plenty of water. Get medical attention. If on skin, immediately flush skin with plenty of water. Get medical attention if irritation persists. May cause allergic skin response.

See inside booklet for complete Precautionary Statements, Directions For Use, and Conditions of Sale and Warranty.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions For Use section for information about this standard.


SHAKE WELL BEFORE USING

Het contents 1 gallon

BASF Corporation

PO Box 13528, Research Triangle Park, NC 27709-3528

ACO60194 ICN8003



Precautionary Statements

Hazards to Humans (and Domestic Animals)

Corrosice. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swal-'owed

Personal Protective Equipment (PPE)

Applicators and other handlers must wear

- Long-sleeved shirt and long pants
- Waterproof gloves
- Chemical-resistant footwear plus socks.
- Protective eyewear

Discard 60thing and other absorbent materials that have been drenched or heavily contaminated with this croduct is concentrate. Do not reuse them, Follow manufacturer's instructions for cleaning/maintaining PPE. If no seen instructions for washables, use detergent and hot water. Keep and wash PPE separately from other aurdry

Engineering Controls Statement:

Whith handlers use closed systems, enclosed cabs, or aircraft in a mariner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet
- Remove closting immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
- Remote PPE immediately after handling this product. Wash the outside of gloves before removing. As such as passible, wash thoroughly and change into clean clothing.

Environmental Hazards

Afragree, which is present in this product, can travel, seep or leach) through soil and can enter groundwate which may be use tias drinking water. Atrazine has been found in groundwater. Users are advised not to is a , Prompt® herbicide to sand and loamy sand sols where the water table (groundwater) is close to the suffice and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can proto first or ofermation on the type of soil in your area and the location of groundwater

This performs to a great invertebrates. For terrestrial uses, do not apply directly to water or to areas where sight in water is present or to intertidal areas below the mean high water mark. Do not apply where or Doinct apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water tive earing integrit prodisposal of wastes

Directions For Use

It is a violation of Federal law to use this product in a manner inco this product in a way that will contact workers or other persons, e ed handlers may be in the area during application. For any require sult the agency responsible for pesticide regulation

Agricultural use requirements

Use this product only in accordance with its labeling and with CFR part 170. This Standard contains requirements for the profarms, forests, nurseries, and greenhouses, and handlers of a requirements for training, decontamination, notification, and e specific instructions and exceptions pertaining to the stateme tive equipment (PPE), and restricted-entry interval. The require this product that are covered by the Worker Protection Standa

Do not enter or allow worker entry into treated areas during the r

PPE required for early entry to treated areas that is permitted unithat involves contact with ariything that has been treated, such a

- Coveralls
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective evewear

Non-agricultural Use Requirements

The requirements in this box apply to uses of this product that a Protection Standard for agricultural pesticides (40 CFR Part 170) used to produce agricultural plants on farms, tilrests, nurseries,

Do not enter treated areas without protective clothing until spray

General Information

compt is intended for the posternergence control of a broad spe es not control grasses. Prompt is effective mainly through conf bughly covered with spray. Large leaf canopies shelter smaller we The desired plants recommended for use in this labeling are tolera slight leaf speckling of desired plants may occur but these plants

Timing of Applications

Apply Prompt early postemergence when weeds are small and ac maximum size listed in the Application Rate Table.

Early application to weeks produces the most beneficial effection idepending on weed species), and makes it easier to obtain thoro which permits weeds to exceed the maximum size stated will resu



Water Volume and Spray Pressure

Apply recommended rates of Prompt as follows:

Ground Equipment

Use a minimum of 20 gats, of water per broadcast acre and a minimum of 40 psi pressure (measured at the boom, not at the pump or in the line). When weed foliage is dense use up to 50 gats, of water and up to 80 psi pressure. Use standard high pressure pesticide hollow cone or flat fan nozzles for best weed coverage.

Special Information on Irrigation

In irrigated areas, it may be necessary to irrigate prior to treatment with **Prompt** to ensure weeds are growing actively. Weeds growing under drought conditions usually are not satisfactority controlled, so it will be necessary to irrigate these areas before application or wait until adequate rainfall has been received.

NON-FOOD USE

T

For use in St. August negrass, zoysia and centipedegrass, Applying earlier than 10 days after sprigging or plugging

Special Directions for Other Weed Problems

Canada Thistle

Apply 37 pints of **Prompt** per acre when plants are from 8-10 inches tall to the bud stage. Make a follow-up application 7-10 days, after using the same rate of **Prompt**, or use 2 pints of **Basagran® herbicide** per acre **Yellow Nutsedge**.

A single application of **Prompt** at 3+ pints per acre will provide suppression of vellow nutsedge. For control of the low nutsedge apply 3+ pints of **Prompt** per acre when plants are 1 to 4 inches tall. Make a follow-up application 2-10 days later using the same rate of **Prompt** or 1 (to 2 pints of **Basagran** per acre. **Field Bindweed**

For suppression of the 1 contineed apply 3 counts of **Prompt** per aure when field bindweed vices are 8 to 10 months on p

Restrictions and Limitations

Or in trace, Prompt when desired plants are under stress from prolonged cold, wet weather, poor fertility in their factors.

Eximit acculingment to bis red blants snow injury ileaf phytotoxicity and or plant sturiting) produced by any interstrict past one list illustrations it eliable this injury may be enhanced and or prokingled.

Extra lact . Prompt during the orded periods of drought or during unseasonably cold weather as unsais to the week of the day west.

Contributed season feed treated forage to is estock.

Bair fail or overhead masticn soor lafter application (within 8 hours) may no lify the effectiveness of **Prompt.** Do not account is credited to reach any type of inigation system.

Carrier Collatering Itamin at immay beine duced by diking and flooring of permanent liquid bulk storage sites to this of intermedice in stern.

This find is the wind frem lend in eaded for used within 30 feet of all wells including abandoned wells. This is expensely that he wells are the release.

Application Rate Table for St. Augustinegrass, Zoysia, and Centipedegrass

Application Rates for Weed Growth Stages**						_	
Weeds Controlled	2 pints per acre		2% pints per acre		3½ pints per acre		_
Annuals	Leaf Stage	Maximum Height(in)	Leaf Stage	Maximum Height(in)	Leaf Stage	Maximum Height(in)	_
Annual Morningglory		_	Up to 4	4	4-6	6	-
Beggarticks		~			Up to 6	6	
Bristly Starbur	_			_	Up to 4	2	
Cocklebur	2-4**	3	2-10"	8	2-10**	2	
Common Lambsquarters	1 1	_	Up to 8	5	δ-12	8	i
Common Ragweed	_	_	Up to 5	ă	4.7	5	i
Corn Speedwell		_	6	3	9	5	1
Cudweed		1	š	2	8	2	
Dayflower	!		<u> </u>		Up to 6	3	
Giant Ragweed	i _ /	~	Up to 4	4	4.6	4	i
Hopclover	_	_ !	Op 10 4	-	4.0	6 2 8	i
Jimsonweed	2-4	3	Up to 6	1	<u> </u>	2	İ
Ladysthumb	2-6	ă	Up to 10	9	6-10		1
Oldfield Toagflax	-		Optoro	9	10-14	12	Ì
Oxalis			- 1	- i	6	8	i
Pennsylvania Smartweed	2-6	4	Up to 10	9	6	3	!
Prickly Sida or Teaweed	1 -	•	OPIGIO	9	10-14	12	İ
Redroot Pigweed		_		_	Up to 6	3	1
Small Hopclover	1	_	Up to 10	6	1	_	į
Smallflower Morningglory	1 = 1	_		- !		2	,
Smooth Pigweed	i — I	~]	Up to 4	4	4 6	6	1
Spurred Anoda		-	Up to 10	6	-	_	i
Tall Waterhemp	· ·	_		— j	Up to 6	3	i
Velvetleaf	2.4	3	Up to 8	2	6 9	4	:
Venice Mallow	'	3	Up to 6	5	りる	8	
Wild Buckwheat	_		Up to 8	4	-		
Wild Mustard		-	Up to 4	3	: -	5	
.Vild Sunflower	_	-	Up to 6	4 .	٠, ٠	8	
Lind Somower			Up to 5	6,	: •	8	


Perennials Canada Thistle Field Bindweed Foolow Nutsedge See Special Directions on page 4 See Special Directions on page 4 See Special Directions on page 4

Storage and Disposal

Store above 151 F

Do not contaminage water, food or feed by storage or disposal. Pesticide wastes are in the initiatious improper disposal of excess pesticide, spray mixture or misate is a violation of Ferdinal will fitness wastes cannot be disposed of by use according to fabel instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidation in the entire container (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary land fill or by incineration, or, if allowed by state and local authorities, by burning, if burned stay out of amoke

[&]quot;Do not treat earlier transleaf stage shown and do not count cotyledon leaves



In Case of Emergency

In case of large-scale spillage regarding this product: avoid contact, isolate area and keep out animals

and unprotected persons. Confine spill and call:

CHEMTREC BASE Corporation

800-424-9300 800-832- HELP

In case of medical emergency regarding this product, call:

- 1 Your local doctor for immediate treatment,
- 2 Your local poison control center (hospital),
- 3. BASE Corporation 800-832-HELP.

Appendix of Scientific Names

The following are scientific names for the weeds listed in this label. For specific recommendations on control of these weeds, refer to the **Application Rate Table**.

Broadleaf Weeds

Common Name	Scientific Name		
Beggarticks	Bidens frondosa		
Bristly Starbur	Acanthospermum hispidum		
Canada Thist e	Circium arvense		
Cockiebur	Xanthium strumarium		
Common Lambsc Litters	Chenopodium album		
Common Ragwer	Ambrosia arternisiifolia		
Corn Speed ve i	Veronic arvensis		
Cudweed	Gnaphalium spp		
Dayflower	Commelina spp		
Field Bindweed	Convolvulus arvensis		
Giant Ragweed	Ambrosia trifida		
Hopciover	Trifolium aureum		
Hopciover, Sma	Trifolium dubium		
Jimsonweed	Datura stramonium		
Lädysthumb	Polygonum persicaria		
Morhinggion, Arinua	Ipomea spp		
Oidfie d'Toadflax	Linaria canadensis		
Oxads	Oxalis stricta		
Pentisy van a Smartweed	Polygonum pensylvanicum		
Prickly Sida or Teaweed	Sida spinosa		
Redroot Pigweed	Amaranthus retroflexus		
Morninggier, Smalfiblier	Jacquemontia famnifolia		
Smooth Pigweed	Amaranthus hybridis		
Spurred Anoda	Anoda cristata		
Tall Watertlemp	Amaranthus tuberculatus		
Velvet eaf	Abutiton theophras 'i		
Venice Malina	Hibiscus trionium		
% d B ic+wheat	Polygonum convolvulus		
하 네 Mustard	Sinapis arvensis		
# 35 m 0 k v	Heiranthus annua		
Yelfow Mutsedge	Cyperus esculentus		

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The circctions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASE CORPO-BATION ("BASE") or the Seller All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONBEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Bisagran is a registered trademark of BASF AG

Prompt is a registered trademark of BASE Corporation

The Specialty Products Logo is a trademark of BASE Corporation

⇒ 1994 BASE Corporation

6

7

 \wedge