



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460 MAR 1 7 1994

Abraham J. Tobia BASF CORP AGRICULTURAL PRODUCTS BOX 13528 RESEARCH TRIANGLE PARK, N.C. 27709

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Subject:

Label Amendment Submission of 11/17/93 in Response to PR Notice 93-7

EPA Reg. No. 7969-105 POLYRAM 80 DF

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling AND
- WITHIN one year from date of this acceptance.

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely

Jim Tompkins, Deputy Chief Registration Support Branch Registration Division (7505W)

Attachment

Page 3

BASF CORP POLYRAM 80 DF 7969-105 11/17/93 Original Submission

The following statement on your proposed label is not required for this product: 'Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate." You may delete it or leave it on the label.

The requirement under the heading "Personal Protective Equipment" must state: "Applicators and other handlers must wear:". Cross out your lead-in statement and replace it with correct one. EXCEPTION: If you have an entirely separate PPE section specifying what "Mixers and loaders must wear", the requirement under the heading "Personal Protective Equipment" must state: "Applicators and other handlers (other than mixers and loaders) must wear:". Cross out your lead-in statement and replace it with the correct one.

Delete the crossed-out statements on your proposed label. They are redundant statements or phrases.

Correct the typographical errors circled on your proposed label.

Polyram® 80 DF fungicide

For agricultural or commercial use only.

Active Ingredients:

[ethylenebis (dithiocarbamate)]zinc with 1 part by weight (16.1%) ethylenebis(dithiocarbamic acid) bimolecular and	
trimolecular cyclicanhydrosulfides and disulfides	80.0%
Inert Ingredients	
Zinc expressed as metallic - 14.0%	
TOTAL	100.0%

EPA Reg. No. 7969-105

EPA Est. No. 7969-WG-01

KEEP OUT OF REACH OF CHILDREN CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. See next page for additional precautionary statements.

Statement of Practical Treatment

If swallowed: Dilute by giving two glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person. If inhaled: Remove subject to fresh air; if not breathing give artificial respiration, preferably by mouth; get medical attention. If on skin: Wash with plenty of soap and water; get medical attention. If in eyes: Flush with plenty of water; consult a physician if irritation persists.

Agricultural Use Requirements


Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

See the attached booklet for complete Precautionary Statements, Directions for Use, and Conditions of Sale and Warranty.

Net weight 30 Pounds

BASF Corporation

PO Box 13528 Research Triangle Park, North Carolina 27709-3528



5 11

Precautionary Statements (continued)

Personal Protective Equipment

Applicators and other handlers must wear:

- Coveralls over longsleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Mixers and Loaders must wear:

- Coveralls over longsleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear
- 👾 Chemical-resistant apron/ Discard clothing or other absorbent materials that have been drenched or heavily contaminated, with product's concnetrate. not reuse them. Follow manufacturer's instructions for cleaning/maintining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. During application, human flaggers must be in enclosed cabs.

Engineering Controls Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Apply this product only as specified on the label.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Storage and Disposal

Pesticide Storage: Do not contaminate water, food or feed by storage or disposal.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact vour State Pesticide or **Environmental Control** Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out

6811

of smoke.

In Case of Emergency

In case of large-scale spillage regarding this product, call:

CHEMTREC 800-424-9300 BASF Corp 800-832-HELP

In case of medical emergency regarding this product, call:

- 1. Your local doctor for immediate treatment
- 2. Your local poison control center (hospital)
- 3. BASF Corp 800-832-HELP

All applicable directions, restrictions, precautions and conditions of sale and warranty are to be followed.

This labeling must be in the possession of the user at the time of application.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses. and handlers of agricultural pesticides. it contains requirements for training. decontamination.notification. and emergency assistance.

It also contains specific emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls overlong-sleeved shirt and long pants
- Waterproof gloves
- Shous plus socks

If this product is used on a crop no other product containing a different EBDC active ingredient may be used on the same crop during the same growing season.

Apples

General Information

Polyram 80 DF fungicide is effective for the control of apple scab and apple rusts. Polyram 80 DF is a contact fungicide; therefore, thorough coverage of plant parts to be protected is essential for effective disease control.

Time and Rate of Application

it is recommended that Polyram 80 DF be used in an Integrated Pest Management program. Use either the "pre-bloom" or "extended application" schedule specified in Application Rates and Timings table. Do not combine or integrate the two treatment schedules. Use onlv on а protective schedule.

Method of Application

Rates of Polyram 80 DF per 100 gallons are based on an application of 300 gallons per acre of dilute spray. Refer to Application Rates and Timing table if other than 300 gallons per acre is applied.

Apply Polyram 80 DF as a spray using sufficient water to obtain thorough coverage.

Ground equipment: Apply a minimum of 20 gallons per acre.

Air equipment: Apply a minimum of 10 gallons per acre.

Mixing Instructions

Maintain agitation to keep the material in suspension and apply with properly calibrated spray equipment.

Tank mixtures: Polyram 80 DF is compatible with most of the apple pesticides with which it might normally be used, including spray oils.

When used with oil. Kelthane, or Diazinon. Polyram 80 DF should be thoroughly mixed with water in the spray tank and tank filled nearly full of water before the other material is Do not combine added. Polyram 80 DF with fixed coppers or highly alkaline materials such as Bordeaux mixture or lime sulfur. Observe all cautions and limitations on labeling of all products used in mixtures.

Restrictions and Limitations for Apples

Do not combine or integrate the "pre-loom" and "extended application" schedules.

Do not graze livestock in treated areas.

Pre-bloom Use

Do not exceed 6 pounds of **Polyram 80 DF** per acre per application.

Do not apply more than 4 applications of Polyram 80 DF per season.

Do not apply Polyram 80 DF after bloom.

Do not apply more than 24 pounds of Polyram 80 DF per acre per year.

Extended Application or Use in Tank Mixtures

Do not apply more than 3 pounds of **Polyram 80 DF** per acre per application.

Do not apply more than 7 applications of **Polyram 80 DF** per season.

Do not apply more than 21 pounds of Polyram 80 DF per acre per year.

Do not apply within 77 days of harvest.

Potatoes

General Information

Polyram 80 DF fungicide is effective for the control of early blight and late blight.

Polyram 80 DF is a contact fungicide, therefore thorough coverage of plant parts to be protected is essential for effective disease control.

Time and Rate of Application

It is recommended that Polyram 80 DF be used in an Integrated Pest Management program.

Begin application when plants are 4 to 6 inches high by applying 2 pounds per acre. (Application intervals - 5 to 10 days. Refer to the Application Rates and Timings table.) The addition of a surfactant, that based on local experience, has been shown to cause phytotoxic effect, to spray solutions will improve performance. Also vine kill should occur 14 days before harvest.

Method of Application

Rates of Polyram 80 DF for potatoes are based on an application of 100 gallons per acre of dilute spray.

Apply Polyram 80 DF as a spray using sufficient water to obtain thorough coverage. Refer to the Application Rates and Timings table if other than 100 gallons per acre is applied.

Ground equipment: Apply a minimum of 15 gallons per

acre.

Air equipment: Apply in a minimum of 5 gallons per acre.

Mixing Instructions

Maintain agitation to keep the material in suspension and apply with properly calibrated equipment.

Tank Mixtures: Polyram 80 DF is compatible with most of the potato pesticides with which it might normally be used. Do not apply Polyram 80 DF with fixed coppers or highly alkaline materials such as Bordeaux mixture or lime sulfur. Observe all cautions and limitations on labeling of all products used in mixtures.

Chemigation

Apply by chemigation only to potatoes using only sprinkler irrigation systems. Do not apply chemigation to any other crop or to this crop using any other type of irrigation system.

Apply this product only through one of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation systems. Crop injury, lack of

effectiveness, illegal 10 pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration. you should contact State Extension specialists. Service equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. person knowledgeable of the chemigation system responsible for its operation, or under the supervision of the responsible person. shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional valve. check vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent watersource contamination from backflow. The pesticide injection pipeline must contain а functional, automatic. quick-closing check valve to prevent the flow of fluid back toward the injection The pump. pesticide injection pipeline also must contain functional, normally-closed, solenoid-operated valve located on the intake side of

pump injection the and connected to the system interlock to prevent fluid from being withdrawn from the vlaque tank when the irrigation system is either automatically or manually shut down. The system must contain a functional interlocking control to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump motor must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where a pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Agitation must be maintained in supply tank.

Restrictions and limitations for Potatoes

Do not apply more than 2 pounds of Polyram 80 DF per acre per application.

9:11

Do not apply more than 14 pounds of Polyram

80 DF per acre per season.

Do not exceed 7 applications of Polyram 80 DF per season.

A minimum preharvest interval of 3 days is required in CT, DE, FL, MA, ME, MI, NH, NY, OH, PA, RI, VT, and WI, and at least 14 days elsewhere.

Roses

General Information

Polyram 80 DF fungicide is effective for the control of Cercospora Leaf Spot, Rust, and Black Spot of roses. Polyram 80 DF is a contact fungicide, therefore, thorough coverage of plants to be protected is essential for effective disease control.

Time and Rate of Application

It is recommended that Polyram 80 DF be used in an Integrated Pest Management program. Spray Polyram 80 DF when first leaves unfold; repeat weekly as needed.

During periods continuously rainy weather shorten application period to twice a week. Refer to Application Rates and Timings table for use rates of Polyram 80 DF on roses.

Method of Application

Rates of Polyram 80 DF for roses is expressed on application of 100 gallons per acre of dilute spray. Refer to the Application Rates and Timings table if other than 100 gallons per acre is applied.

Treatment: When using knapsack sprayers or other suitable small area spraye quipment mix 6 teaspoonfuls per 1 gallon of water. Apply to the foliage on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

Restrictions and Limitations for Roses

Do not exceed 2 applications of Polyram 80 DF per week.

Application Rates and Timing							
Crop	Diseases Controlled	Pounds of DF per App	_ •	Remarks (Also refer to Directions For Use)	Restriction		
Apples Use either the "pre-bloom" or "extended application" schedule. Do not combine or integrate the two-	Scab Rusts	2.0 Rates are based on 300 GPA Do not exc product per applica	acre per	Pre Bloom Use: Begin applications at % to % inch green tip and continue on a 7-to 10-day schedule through bloom.	Do not apply more than 4 applications per season. Do not apply after bloom. Do not apply more than 24 lbs. per acre per year. Do not graze livestock in treated areas.		
treetment schedules.		1.0 Rates are based on 300 GPA Do not exc product per applica	acre per	Extended Application Schedule or for Use in Tank Mixtures: It is recommended that this product be used in an Integrated Pest Management Program. For implementation of IPM programs or for use as a resistence management tool: applications based on tree-row volume. Begin applications at % to % inch green tip and continue applications on a 7- to 10-day schedule through the second cover spray or to within 77 days of hervest.	Do not apply more than 7 applications per season. Do not apply within 77 days of harvest. Do not apply more than 21 lbs. per acre per year. Do not graze livestock in treated areas.		
Potatoes	Early Blight Late Blight	2.0 2.0 Rates are based on 100 GPA Do not exceed 2 lb product per acre per application		Begin application when plants are 4 to 6 inches high by applying 2 lb. per acre. (Application intervals - 5 to 10 days.) The addition of a surfactant, that based on local experience, has been shown to cause no phytotoxic effect, to spray solutions will improve performence. It is recommended that this product be used within an Integrated Pest Management Program. Also vine kill should occur 14 days before harvest.	Do not apply more than 14 lbs. per acre per season. Do not exceed 7 applications per season. A minimum prehervest interval of 3 days is required in CT, DE, FL, MA, ME, MI, NH, NY, OH, PA, RI, VT and WI, and at least 14 days elsewhere.		
Roses	Cercospora Leaf Spot Rust Black Spot	1.0 to 1.5 Rates are based on 100 GPA Do not exce product per applica	acre per	Spray when first leaves unfold; repeat weekly as needed. During periods of continuously rainy weather, shorten application period to twice a week. Start off at 1.0 ib per acre; increase to 1.5 ib per acre if disease pressure increases.	Do not exceed 2 applications per week.		

Conditions of Sale and Warranty

The Directions of use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this Crop injury, product. other ineffectiveness or unintended consequences may result because of such factors weather as conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF

CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer. BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in Use. **Directions** for the subject to the inherent risks referred to above. BASF OTHER MAKES NO EXPRESSED OR IMPLIED WARRANTY OF FITNESS OR **MERCHANTABILITY OR ANY** IMPLIED OTHER IN NO CASE WARRANTY. **BASF** OR THE SHALL SELLER BE LIABLE **FOR** CONSEQUENTIAL, SPECIAL,


INDIRECT **DAMAGES** RESULTING FROM THE USE HANDLING OF THIS OR PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Polyram is a registered trademark of BASF Aktiengessellschaft.

• 1993 BASF Corporation

BASE Corporation
PO Box 13528
Research Triangle Park, NC 27709







UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

MAR 1 7 1994

Abraham J. Tobia
BASF CORP
AGRICULTURAL PRODUCTS
BOX 13528
RESEARCH TRIANGLE PARK, N.C. 27709

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Subject:

Label Amendment Submission of 11/17/93 in Response to PR Notice 93-7

EPA Reg. No. 7969-105 POLYRAM 80 DF

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling AND
- WITHIN one year from date of this acceptance.

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,

Jim Tompkins, Deputy Chief Registration Support Branch Registration Division (7505W)

Attachment

BASF CORP POLYRAM 80 DF 7969-105 11/17/93 Original Submission

The following statement on your proposed label is not required for this product: "Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate." You may delete it or leave it on the label.

The requirement under the heading "Personal Protective Equipment" must state: "Applicators and other handlers must wear:". Cross out your lead-in statement and replace it with correct one. EXCEPTION: If you have an entirely separate PPE section specifying what "Mixers and loaders must wear", the requirement under the heading "Personal Protective Equipment" must state: "Applicators and other handlers (other than mixers and loaders) must wear:". Cross out your lead-in statement and replace it with the correct one.

Delete the crossed-out statements on your proposed label. They are redundant statements or phrases.

Correct the typographical errors circled on your proposed label.

Polyram® 80 DF fungicide

For agricultural or commercial use only.

Active Ingredients:

A mixture of 5.2 parts by weight (83.9%) of ammoniates of	
[ethylenebis (dithiocarbamate)]zinc with 1 part by weight	
(16.1%) ethylenebis(dithiocarbamic acid) bimolecular and	
trimolecular cyclicanhydrosulfides and disulfides	80.0%
Inert Ingredients	20.0%
Zinc expressed as metallic - 14.0%	
TOTAL 1	00.0%

EPA Reg. No. 7969-105

EPA Est. No. 7969-WG-01

KEEP OUT OF REACH OF CHILDREN CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. See next page for additional precautionary statements.

Statement of Practical Treatment

If swallowed: Dilute by giving two glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person. If inhaled: Remove subject to fresh air; if not breathing give artificial respiration, preferably by mouth; get medical attention. If on skin: Wash with plenty of soap and water; get medical attention. If in eyes: Flush with plenty of water; consult a physician if irritation persists.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

See the attached booklet for complete Precautionary Statements, Directions for Use, and Conditions of Sale and Warranty.

Net weight 30 Pounds

BASF Corporation

PO Box 13528 Research Triangle Park, North Carolina 27709-3528

MAR 1 7 1994

ter the Federal Insecticide dicide, and Rodemicide Assembled, for the positive marcal junger EPA Neg. N.

Precautionary Statements (continued)

Personal Protective Equipment

مان کیائی Applicators and other handlers must wear:

- Coveralls over longsleeved shirt and long pants
- · Waterproof gloves
- Shoes plus socks

Mixers and Loaders must wear:

- Coveralls over longsleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear Protective
- Chemical-resistant apron/ Discard clothing or other absorbent materials that have been drenched or heavily contaminated with product's conchetrate. Do not reuse them. manufacturer's instructions for cleaning/maintining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. During aerial application, human flaggers must be in enclosed cabs.

Engineering Controls Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Apply this product only as specified on the label.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Storage and Disposal

Pesticide Storage: Do not contaminate water, food or feed by storage or disposal.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact State Pesticide vour or **Environmental Control** Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for quidance.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out

of smoke.

In Case of Emergency

In case of large-scale spillage regarding this product, call:

CHEMTREC 800-424-9300 BASF Corp 800-832-HELP

In case of medical emergency regarding this product, call:

- 1. Your local doctor for mmediate treatment
- 2. Your local poison control center (hospital)
- 3. BASF Corp 800-832-HELP

All applicable directions, restrictions, precautions and conditions of sale and warranty are to be followed.

This labeling must be in the possession of the user at the time of application.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a that will contact workers or other persons. either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. lt contains training, requirements for decontamination, notification, and emergency assistance.

It-also-contains-specifie emergency-assistance- It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

If this product is used on a crop no other product containing a different EBDC active ingredient may be used on the same crop during the same growing season.

Apples

General Information

Polyram 80 DF fungicide is effective for the control of apple scab and apple rusts. Polyram 80 DF is a contact fungicide; therefore, thorough coverage of plant parts to be protected is essential for effective disease control.

Time and Rate of Application

It is recommended that Polyram 80 DF be used in an Integrated Pest Management Use either the program. "pre-bloom" or "extended application" schedule specified in Application Rates and Timings table. Do not combine or integrate the two treatment schedules. only on а protective schedule.

Method of Application

Rates of Polyram 80 DF per 100 gallons are based on an application of 300 gallons per acre of dilute spray. Refer to Application Rates and Timing table if other than 300 gallons per acre is applied.

Apply Polyram 80 DF as a spray using sufficient water to obtain thorough coverage.

Ground equipment: Apply a minimum of 20 gallons per acre.

Air equipment: Apply a minimum of 10 gallons per acre.

Mixing Instructions

Maintain agitation to keep the material in suspension and apply with properly calibrated spray equipment.

Tank mixtures: Polyram 80 DF is compatible with most of the apple pesticides with which it might normally be used, including spray oils.

When used with oil. Kelthane, Diazinon, ٥r Polyram 80 DF should be thoroughly mixed with water in the spray tank and tank filled nearly full of water before the other material is added. Do not combine Polyram 80 DF with fixed coppers or highly alkaline materials such as Bordeaux lime sulfur. mixture or Observe all cautions and limitations on labeling of all products used in mixtures.

Restrictions and Limitations for Apples

Do not combine or integrate the "pre-loom" and "extended application" schedules.

Do not graze livestock in treated areas.

Pre-bloom Use

Do not exceed 6 pounds of **Polyram 80 DF** per acre per application.

Do not apply more than 4 applications of Polyram 80 DF per season.

Do not apply Polyram 80 DF after bloom.

Do not apply more than 24 pounds of Polyram 80 DF per acre per year.

Extended Application or Use in Tank Mixtures

Do not apply more than 3 pounds of Polyram 80 DF per acre per application.

Do not apply more than 7 applications of Polyram 80 DF per season.

Do not apply more than 21 pounds of Polyram 80 DF per acre per year.

Do not apply within 77 days of harvest.

Potatoes

General Information

Polyram 80 DF fungicide is effective for the control of early blight and late blight.

Polyram 80 DF is a contact fungicide, therefore thorough coverage of plant parts to be protected is essential for effective disease control.

Time and Rate of Application

It is recommended that **Polyram 80 DF** be used in an Integrated Pest Management program.

Begin application when plants are 4 to 6 inches high by applying 2 pounds per acre. (Application intervals - 5 to 10 days. Refer to the Application Rates and Timings table.) The addition of a surfactant, that based on local experience, has been shown to cause пο phytotoxic effect, to spray solutions will improve performance. Also vine kill should occur 14 days before harvest.

Method of Application

Rates of Polyram 80 DF for potatoes are based on an application of 100 gallons per acre of dilute spray.

Apply Polyram 80 DF as a spray using sufficient water to obtain thorough coverage. Refer to the Application Rates and Timings table if other than 100 gallons per acre is applied.

Ground equipment: Apply a minimum of 15 gallons per

acre.

Air equipment: Apply in a minimum of 5 gallons per acre.

Mixing Instructions

Maintain agitation to keep the material in suspension and apply with properly calibrated equipment.

Tank Mixtures: Polyram 80 DF is compatible with most of the potato pesticides with which it might normally be used. Do not apply Polyram 80 DF with fixed coppers or highly alkaline materials such as Bordeaux mixture or lime sulfur. Observe all cautions and limitations on labeling of all products used in mixtures.

Chemigation

Apply by chemigation only to potatoes using only sprinkler irrigation systems. Do not apply chemigation to any other crop or to this crop using any other type of irrigation system.

Apply this product only through one of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation systems. Crop injury, lack of

effectiveness. Of illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about vou calibration. should State Extension contact specialists. Service equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safetv devices for public water systems are in place. person knowledgeable of the chemigation system responsible for its operation, or under the supervision of the responsible person, shall shut the system down make necessarv and adjustments should the need arise.

The system must contain a functional check valve. vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent watersource contamination from backflow. The pesticide pipeline must injection contain a functional. automatic, quick-closing check valve to prevent the flow of fluid back toward the injection The pump. pesticide injection pipeline must contain also functional, normally-closed, solenoid-operated valve located on the intake side of

injection the gump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain a functional control interlocking to automatically shut off the pesticide injection when the water pump motor stops. The irrigation line or water pump motor must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where a pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Agitation must be maintained in supply tank.

Restrictions and limitations for Potatoes

Do not apply more than 2 pounds of Polyram 80 DF per acre per application.

Do not apply more than 14 pounds of Polyram 80 DF per acre per season.

Do not exceed 7 applications of Polyram 80 DF per season.

A minimum preharvest interval of 3 days is required in CT, DE, FL, MA, ME, MI, NH, NY, OH, PA, RI, VT, and WI, and at least 14 days elsewhere.

Roses

General Information

Polyram 80 DF fungicide is effective for the control of Cercospora Leaf Spot, Rust, and Black Spot of roses. Polyram 80 DF is a contact fungicide, therefore, thorough coverage of plants to be protected is essential for effective disease control.

Time and Rate of Application

It is recommended that Polyram 80 DF be used in an Integrated Pest Management program. Spray Polyram 80 DF when first leaves unfold; repeat weekly as needed.

During periods continuously rainy weather shorten application period to twice a week. Refer to Application Rates and Timings table for use rates of Polyram 80 DF on roses.

Method of Application

Rates of Polyram 80 DF for roses is expressed on application of 100 gallons per acre of dilute spray. Refer to the Application Rates and Timings table if other than 100 gallons per acre is applied.

Spot Small or Area Treatment: When using knapsack sprayers or other suitable small area spray equipment mix teaspoonfuls per 1 gallon of water. Apply to the foliage on a spray-to-wet basis. Spray coverage should be uniform and complete. not spray to the point of runoff.

Restrictions and Limitations for Roses

Do not exceed 2 applications of Polyram 80 DF per week.

Application Rates and Timing						
Crop	Diseases Controlled	Pounds of Polyram DF per Application		Remarks	Restriction	
		100 gal	Acre	(Also refer to Directions For Use)		
Apples Use either the "pre- bloom" or "extended application" schedule. Do not combine or integrate the two-	Scab Rusts	2.0 Rates are based on 300 GPA Do not exceproduct per application	acre per	Pre Bloom Use: Begin applications at % to % inch green tip and continue on a 7-to 10-day schedule through bloom.	Do not apply more than 4 applications per season. Do not apply after bloom. Do not apply more than 24 lbs. per acre per year. Do not graze livestock in treated areas.	
treatment schedules.		1.0 Rates are based on 300 GPA Do not exceproduct per application	ecre per	Extended Application Schedule or for Use in Tank Mixtures: It is recommended that this product it e used in an integrated Pest Management Program. For implementation of IPM programs or for use as a resistance management tool: applications based on tree-row volume. Begin applications at ¼ to ½ inch green tip and continue applications on a 7- to 10-day schedule through the second cover spray or to within 77 days of harvest.	Do not apply more than 7 applications per season. Do not apply within 77 days of harvest. Do not apply more than 21 lbs. per acre per year. Do not graze livestock in treated areas.	
Patatoes	Early Blight Late Blight	2.0 Rates are based on 100 GPA Do not exc product per applica	ecte per	Begin application when plants are 4 to 6 inches high by applying 2 lb. per acre. (Application intervals - 5 to 10 days.) The addition of a surfactant, that based on local experience, has been shown to cause no phytotoxic effect, to spray solutions will improve performance. It is recommended that this product be used within an Integrated Pest Management Program. Also vine kill should occur 14 days before hervest.	Do not apply more than 14 lbs. per acre per season. Do not exceed 7 applications per season. A minimum preharvest interval of 3 days is required in CT, DE, FL, MA, ME, MI, NH, NY, OH, PA, RI, VT and Wi, and at least 14 days elsewhere.	
Roses	Cercospora Leaf Spot Rust Black Spot	1.0 to 1.5 Rates are based on 100 GPA Do not exce product per applica	acre per	Spray when first leaves unfold; repeat weekly as needed. During periods of continuously rainy weather, shorten application period to twice a week. Start off at 1.0 % per acre; increase to 1.5 % per acre if disease pressure increases.	Do not exceed 2 applications per week.	

Conditions of Sale and Warranty

The Directions of use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness other or unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF

CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer. BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in Directions for the subject to the inherent risks referred to above. BASF MAKES NO OTHER EXPRESSED OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE **FOR** CONSEQUENTIAL, SPECIAL,

INDIRECT **DAMAGES** OR RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it. subject to the foregoing Conditions of Sale and Warranty which may varied only by agreement in writing signed by a duly authorized representative of BASF.

Polyram is a registered trademark of BASF Aktiengessellschaft.

• 1993 BASF Corporation

BASF Corporation
PO Box 13528
Research Triangle Park, NC 27709

