

7969-106

10-8-2003

4/9

OCT 8 2003

Mr. David W. Haughey
BASF Corporation
26 Davis Drive
Research Triangle Park, N.C. 27709

SUBJECT: Amendment/Master Label
Nexter Miticide/Insecticide
EPA Registration Number: 7969-106
Application Dated: August 27, 2003


The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, is acceptable. A stamped copy of the label is enclosed for your records.

Sincerely,



Linda Arrington, Acting
Product Manager (03)
Insecticide Branch
Registration Division 7505c

BASF



Nexter[®]

miticide/insecticide

Active Ingredient:

[2-*tert*-butyl-5-(4-*tert*-butylbenzylthio)-4-chloropyridazin-3(2*H*)-one]75.0%

Inert Ingredients:25.0%

Total:100.0%

Plus water-soluble packaging

EPA Registration Number 7969-106

EPA Est. No.

U.S. Patent Number: 4,877,787

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside booklet for complete **First Aid, Precautionary Statements, Directions For Use, and Conditions of Sale and Warranty.**

Net contents: 2.06 pounds
(five 6.6-ounce water-soluble bags)

Product of Japan; formulated in the United States with U. S. and imported ingredients

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).	

Precautionary Statements

Hazards to Humans and Domestic Animals

WARNING. May be fatal if inhaled. Do not breathe dust or spray mist. For handling activities, use dust/mist filtering respirator (MSHA/NIOSH approval numbers prefix TC-21C), or a NIOSH approved respirator with a N, P, R, or HE prefilter. Wear long-sleeved shirt and long pants, socks and shoes and waterproof gloves. Harmful if swallowed or absorbed through skin. Avoid contact with skin. Remove contaminated clothing and wash before reuse. Causes moderate eye irritation. Do not get in eyes or on clothing. Wear goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves
- Protective eye wear
- For handling activities, use dust/mist filtering respirator (MSHA/NIOSH approval numbers prefix TC-21C), or a NIOSH approved respirator with a N, P, R, or HE prefilter.
- Chemical resistant headgear for overhead exposure.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Keep out of lakes, ponds, or streams. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Do not apply when weather conditions favor drift from target area. Drift or runoff from treated areas may be hazardous to fish in adjacent sites. This product is highly toxic to bees. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. Application early in the morning or at dusk is suggested.

Endangered Species Concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of federal law.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions and **Conditions of Sale** and **Warranty** are to be followed.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours**. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eye wear
- For handling activities, use dust/mist filtering respirator (MSHA/NIOSH approval numbers prefix TC-21C), or a NIOSH approved respirator with a N, P, R, or HE prefilter.
- Chemical resistant headgear for overhead exposure.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage:

Store in a cool, dry place. This package contains water-soluble bags inside a foil liner (overwrap). The

water-soluble bags dissolve in water and the contents will disperse. If all the water-soluble bags are not used, carefully reseal the overwrap. Each overwrap contains five water-soluble bags. Do not remove the water-soluble bags from the overwrap except for immediate use. If exposed to moisture, the water-soluble bags may break.

Pesticide Disposal:

Pesticide wastes are acutely hazardous. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal:

The outer case and inner overwrap packaging of the water-soluble bag should be incinerated or disposed of in a sanitary landfill, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Do not re-use the empty packaging.

In Case of Spill

In case of large-scale spillage regarding this product, call:

CHEMTREC	800-424-9300
BASF Corporation	800-832-HELP

Steps to be taken in case material is released or spilled:

Wear the personal protective equipment specified on the label. Recover the material for re-use according to label whenever possible. Sweep and/or shovel up the spilled material into an appropriate closed container. Avoid the creation of dusty conditions. Remove and wash clothing and personal protective equipment prior to re-use. Keep the spill out of all sewers and open bodies of water.

I. General Information

This package contains **Nexter® miticide/insecticide**, a 75% wettable powder, in water-soluble bags. **Nexter** is a selective contact miticide/insecticide that controls pests in apples, almonds, apricots, cherries, citrus groves and non-bearing citrus nursery beds or greenhouses, cranberries, grapes, nectarines, peaches, pears, pistachio, plums, prunes and tree nut group crops when used at recommended rates. (**Refer to Table 1. Nexter Application Rate Table.**) Nexter provides knockdown and residual control. A good performance evaluation can be made 4-7 days after treatment. For optimum results, **Nexter** should be applied as pest populations build and prior to reaching economic thresholds.

Mite Resistance Management

Naturally occurring strains of mites and insects listed on this label may not be effectively controlled due to reduced sensitivity. If insensitive strains are present in a field, use a product with a different mode of action to ensure control.

Nexter use should be alternated with other miticides as part of a mite management program to minimize resistance. Repeated use of the same miticide has been documented to result in the buildup of resistant strains of mites. To limit the potential for **Nexter** insensitivity development, do not make more applications than those specified under the maximum per-season in **Table 2. Crop-Specific Restrictions and Limitations**. Consult with your local or state extension personnel for advice on miticide use and selection.

Table 1. Nexter® miticide/insecticide Application Rate Table¹

Pests Controlled	Rate per Acre
European red mite	4.4 - 5.2 oz.
Southern red mite	4.4 - 7.0 oz
Apple Rust Mite Blackmargined aphid Broad mite Citrus Bud mite Citrus red mite Citrus rust mite Eastern grape leafhopper (nymphs) False spider mite (Citrus flat mite) Grape leafhopper (nymphs) Peach silver mite Pear rust mite Pink Citrus rust mite Silverleaf whitefly ² Sixspotted mite Sweet potato whitefly ² Texas citrus mite Variegated leafhopper (nymphs) Virginia creeper leafhopper (nymphs) White apple leafhopper (nymphs) ² Willamette spider mite Yellow pecan aphid	5.2 - 10.67 oz
McDaniel spider mite Pacific spider mite Pear psylla	6.6 - 10.67 oz.
Twospotted spider mite	8.8 - 10.67 oz.
Pests Suppressed	Rate per Acre
Apple aphid Brown citrus aphid Citrus root weevil	5.2 - 10.67 oz

¹ Allow a minimum of 30 days between sequential applications of Nexter in crops that allow more than 1 application per season. For rates above 5.2 ounces per acre on citrus, apply Nexter on a 90 day interval.

² Silverleaf whitefly, White apple leafhopper (nymphs), and Sweet potato whitefly are only considered to be suppressed in California.

Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial sprayer cleaner

according to the manufacturer's directions and by triple rinsing the equipment before and after applying this product.

II. Application Instructions

Nexter may be applied by ground equipment using either diluted or concentrated sprays. Apply recommended rates of **Nexter** as instructed by **section VI. Crop-Specific Information**. Spray the last 3 rows windward of surface water using nozzles on only one side with the spray directed away from surface water. Avoid spraying over the tops of trees/vines by adjusting or turning off the top nozzles. Shut off the nozzles on the side away from the grove when spraying the outside row. Shut the nozzles off when turning at the ends of the rows and when passing tree/vine gaps in rows.

Coverage

Apply **Nexter** in sufficient water to ensure thorough coverage of foliage and fruit. Thorough coverage is required for optimum control. Spraying alternate rows may reduce **Nexter** performance. **Nexter** must be applied to each row for optimum control. To achieve adequate coverage, use proper spray pressure, nozzles, nozzle spacing, volume per acre, and tractor speed. Consult spray nozzle and accessory guide for information pertaining to proper equipment calibration.

Ground Application (Broadcast)

Water Volume: Use 100-400 gallons of spray solution per broadcast acre for optimal performance. In Florida, a minimum of 20 gallons of water per acre in citrus may be used.

III. Additives

In general, no additives or adjuvants are necessary for effective use of **Nexter**. However, the use of additives may be considered for certain conditions such as obtaining better spray distribution, adhesion or penetration of product on to leaf or plant surfaces. Consult a local BASF representative or local agricultural authorities or a consultant for more information concerning additives.

IV. General Tank Mixing Information

The phytotoxic potential of **Nexter** has been assessed on a wide variety of plants with no phytotoxicity observed. However, all varieties and cultivars have not been tested with possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Therefore, before using **Nexter**, test the product on a sample of the crop to be treated to ensure that a phytotoxic response will not occur as a result of applications.

Compatibility Test for Mix Components

Before mixing components, always perform a compatibility jar test.

For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Only use water from the intended source at the source temperature.

Add components in the sequence indicated in the **Mixing Order (see below)** using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

Mixing Order

- 1) **Water.** Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
- 2) **Agitation.** Maintain constant agitation throughout mixing and application.
- 3) **Products in PVA bags.** Place any product contained in water-soluble PVA bags such as

Nexter® miticide/insecticide into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.

- 4) **Water-dispersible products** (such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions).
- 5) **Water-soluble products.**
- 6) **Emulsifiable concentrates** (such as oil concentrate when applicable).
- 7) **Water-soluble additives** (such as AMS or UAN when applicable).
- 8) **Remaining quantity of water.** Maintain constant agitation during application.

Carefully remove the recommended number of water-soluble bags from the inner overwrap packaging and carefully place them into the spray water in the mixing tank. Reseal the outer package making sure that no moisture contacts the water-soluble bags. Do not open the water-soluble bags. Allow the bags to completely dissolve. Use the maximum agitation while mixing **Nexter** in the spray tank.

A defoaming agent may also be necessary. Do not attempt to dissolve the water-soluble bags directly in diesel oils or summer spray-type oils. The bags are water-soluble, not oil soluble.

Boron will prevent the water-soluble bags from dissolving. If boron-containing products are to be used, the water-soluble bags containing **Nexter** must be dissolved completely before the boron-containing product can be added to the spray tank. If boron-containing products have been used in previous applications, thoroughly wash the spray tank before using **Nexter**. Always reseal the overwrap package to protect the remaining unused bags.

V. Restrictions and Limitations

- **Preharvest Interval (PHI):** Refer to **Table 2. Crop-Specific Restrictions and Limitations**
- **Restricted Entry Interval (REI): 12 hours.**
- Do not apply **Nexter** by air.
- Except for cranberries, do not apply through any type of **irrigation** equipment.
- Do not use less than 100 gallons of water per acre except the following:
 - 50 gallons on grapes
 - 20 gallons on citrus grown in Florida
- Do not apply **Nexter** to apricots and cherries in California or to cranberries in New York State.
- **Drift:** Do not apply **Nexter** when weather conditions favor drift to surface water. Do not apply within 110 feet upwind of surface water or when windspeed is above 8 mph. Do not apply during a temperature inversion.
- Allow a minimum of 30 days between sequential applications of **Nexter** in crops that allow more than 1 application per season. For rates above 5.2 ounces per acre on citrus, apply **Nexter** on a 90 day interval.
- **Nexter** is not for sale, distribution, or use in Nassau and Suffolk counties in New York State. In the remainder of the state, read and follow all applicable directions, restrictions and precautions on this label.

Table 2. Crop-Specific Restrictions and Limitations

Crop	Minimum Time from Application to Harvest (PHI)(days)	Maximum Rate per Acre per Application (oz.)	Maximum Number of Applications per Season ³	Aircraft Application
Apples	25	10.67	1	No
Apricots ¹	300	10.67	2	No
Cherries ¹	300	10.67	2	No
Citrus	7	10.67	2	No
Cranberries ²	21	10.67	2	No
Grapes	7	10.67	2	No
Nectarines	7	10.67	2	No
Peaches	7	10.67	2	No
Pears (including Oriental)	7	10.67	1	No
Pistachio	7	10.67	2	No
Plums	7	10.67	2	No
Prunes	7	10.67	2	No
Tree Nuts Group (Almond, Beech, Brazil, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory, Macadamia, Pecan, Black Walnut and English Walnut)	7	10.67	2	No

¹ Do not Apply Nexter to apricots and cherries in California.

² Use Nexter in cranberries in CT, DE, ME, MA, NH, NJ, RI and VT only.

³ Allow a minimum of 30 days between sequential applications of Nexter in crops that allow more than 1 application per season. For rates above 5.2 ounces per acre on citrus, apply Nexter on a 90 day interval.

VI. Crop-Specific Information

Apples

See **Table 1** for application rates for specific pests. Apply **Nexter® miticide/ insecticide** in 100 - 400 gallons of water per acre. **Nexter** must be applied to each row for maximum coverage. Use the higher rate of **Nexter** to ensure adequate concentration in mature orchards with dense foliage.

Citrus

Apply 5.2-10.67 ounces of **Nexter** in sufficient water to achieve thorough coverage. For rates above 5.2 ounces per acre, apply Nexter on a 90-day interval. Use the higher rate of Nexter to ensure adequate concentration in full size trees with dense foliage.

When combining **Nexter** with summer oils, use a minimum of 5 gallons of oil and 6.6 ounces of Nexter per acre.

- **In Florida Only.** **Nexter** may be applied in low volume application equipment with a minimum water volume of 20 gallons of water per acre. It is the user's responsibility to ensure thorough spray coverage in these low volume applications.

Cranberries

Nexter is registered for use on cranberries in the states of Connecticut, Delaware, Maine, Massachusetts, New Hampshire, New Jersey, Rhode Island and Vermont.

Nexter is a selective contact miticide/insecticide that controls southern red mite in cranberries when used at the recommended rates. Complete spray coverage of both upper and lower leaf surfaces is essential for optimal performance.

Applications should be made either early season (mid-May to mid-June) or after fruit set (mid-July through August). Do not apply **Nexter** when bees are actively foraging. The preharvest interval of 21 days must be observed.

Nexter may be applied by chemigation or by ground equipment. Sufficient water volume is necessary to obtain complete coverage of the spray target. Apply 3.5 - 7.0 ounces of **Nexter** in no less than 100 gallons and no more than 600 gallons of water per acre. If using chemigation, use an injection system protected by backflow equipment.

Chemigation: Apply this product only through solid set or hand-move sprinkler systems. Do not apply this product through any other type of irrigation system. Lack of effectiveness can result from non-uniform distribution of treated water. Use only in sprinklers

that apply uniformly and have appropriate check valves. When application of pesticide is complete, thoroughly flush the injection system and sprinkler lines with a minimum volume of water for complete rinse-out. The system must contain a functional check valve or appropriate gooseneck pipe loop, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or, in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a Venturi injector on the discharge side of the pump, or a metering pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Chemigation Systems Connected to Public Water Systems:

Public water system means a system for the provision of piped water to the public for human consumption, if such a system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days of the year. Chemigation systems connected to public water systems must contain a functional reduced-pressure-zone (RPZ) backflow preventer, or the functional equivalent, in the water supply line upstream from the point of pesticide introduction. As an additional option to the RPZ, the water from a public water system can be discharged into a reservoir tank prior to pesticide introduction. There should be a complete physical break (air gap) of at least twice the inside diameter of the fill pipe between the outlet end of the fill pipe and the top of the overflow rim of the reservoir tank.

Grapes

See **Table 1** for application rates for specific pests. Apply **Nexter® miticide / insecticide** in 50 - 400 gallons of water per acre. **Nexter** must be applied to each row for maximum coverage. Use the higher rate of **Nexter** to ensure adequate concentration in mature vineyards with dense foliage.

Pears

Pear (Including oriental)

See **Table 1** for application rates for specific pests. Apply **Nexter** in 100-400 gallons of water per acre. **Nexter** must be applied to each row for maximum coverage. Use the higher rate of **Nexter** to ensure adequate concentration in mature orchards with dense foliage.

In pears, applications may be made early from pink through petal-fall to control eggs, early pear psylla instars and motile mites. **Nexter** is also effective when applied after petal fall as mite populations begin to build.

Stone Fruits

Apricots, Cherries, Nectarines, Peaches, Plums and Prunes

See **Table 1** for application rates for specific pests. Apply **Nexter** in 100 - 400 gallons of water per acre. **Nexter** must be applied to each row for maximum coverage. Use the higher rate of **Nexter** to ensure adequate concentration in mature orchards with dense foliage. For best control, pest populations must be building with primarily immature stages present at the time of application.

Cherries and apricots are treated after spring harvest and have a 300-day PHI. **Nexter** cannot be used on these two crops in California.

Tree Nuts Group and Pistachio

See **Table 1** for application rates for specific pests. Apply **Nexter** in 100 - 400 gallons of water per acre for tree nuts and pistachios. Use the higher rate of **Nexter** to ensure adequate concentration in full-sized trees with dense foliage.

For best control, pest populations must be building with primarily immature stages present at the time of application. **Specifically for almonds:** Applications may be made early from shuck split through midsummer.

Crops

This product can be used on the following crops:

Almond	Chestnut	Nectarines
Apple	Chinquapin	Peaches
Apricots	Citrus	Pear
Beech	Cranberries	Pecan
Black walnut	English Walnut	Pistachio
Brazil nut	Filbert	Plums
Butternut	Grapes	Prunes
Cashew	Hickory	
Cherries	Macadamia	

Look inside for complete **Restrictions and Limitations** and **Application Instructions**.

Pests Listed in This Label

Broad mite	Family: Tarsonemidae <i>Polyphagotarsonemus latus</i>
False spider mite	Family: Tenuipalpidae <i>Brevipalpus phoenicis</i>
Citrus flat mite	<i>Brevipalpus lewisi</i>
Apple rust mite	Family: Eriophyidae <i>Aculus schlectendali</i>
Citrus bud mite	<i>Aceria sheldoni</i>
Citrus rust mite	<i>Phyllocoptruta oleivora</i>
Peach silver mite	<i>Aculus fockeui</i>
Pear rust mite	<i>Eptimerus pyri</i>
Pink citrus rust mite	<i>Aculops pelekassi</i>
Citrus red mite	Family: Tetranychidae <i>Panonychus citri</i>
European red mite	<i>Panonychus ulmi</i>
McDaniel spider mite	<i>Tetranychus mcdanieli</i>
Pacific spider mite	<i>Tetranychus pacificus</i>
Sixspotted mite	<i>Eotetranychus sexmaculatus</i>
Southern red mite	<i>Oligonychus ilicis</i>
Texas citrus mite	<i>Eutetranychus banksi</i>
Twospotted spider mite	<i>Tetranychus urticae</i>
Willamette spider mite	<i>Eotetranychus willamettei</i>
Citrus root weevil	Family: Curculionidae <i>Pachnaeus litus</i>
Apple aphid	Family: Aphididae <i>Aphis pomi</i>
Blackmargined aphid	<i>Monella caryella</i>
Brown citrus aphid	<i>Toxoptera citricida</i>
Yellow pecan aphid	<i>Monelliopsis pecanis</i>
Sweet potato whitefly	Family: Aleyrodidae <i>Bemisia tabaci</i>
Silverleaf whitefly	<i>Bemisia argentifolii</i>
Pear psylla	Family: Psyllidae <i>Cacopsylla pyricola</i>
Eastern grape leafhopper	Family: Cicadellidae <i>Erythroneura cornes</i>
Grape leafhopper	<i>Erythroneura elegantula</i>
Variegated leafhopper	<i>Erythroneura variabilis</i>
Virginia creeper leafhopper	<i>Erythroneura ziczac</i>
White apple leafhopper	<i>Typhlocyba pomaria</i>

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer. BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

U.S. Patent Number: 4,877,787

Nexter is a registered trademark of Nissan Chemical Industries, Ltd.

© 2003 BASF Corporation

All rights reserved

007969-00106.20031212.NVA 2003-04-055-0240.pdf

Supersedes NVA 2003-04-055-0046

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

BASF

Agricultural Products