



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 28, 2016

Craig D. Kleppe
Product Registration Manager
BASF Corporation
26 Davis Drive
P.O. Box 13528
Research Triangle Park, NC 27709

Subject: Label Amendment – Clarifying use sites etc.
Product Name: Basagran 5L Herbicide
EPA Registration Number: 7969-112
Application Date: May 2, 2016
Decision Number: 517131

Dear Mr. Kleppe:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 7969-112
Decision No. 517131

with FIFRA section 6. If you have any questions, please contact Shanta Adeeb by phone at 703-347-0502, or via email at adeeb.shanta@epa.gov.

Sincerely,

A handwritten signature in black ink, reading "Kathryn V. Montague". The signature is written in a cursive style with a long horizontal flourish extending from the end.


Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure



We create chemistry

Group 6 Herbicide



Basagran® 5L

Herbicide

[Alternate Trade Name] **Rezult® B herbicide**

For postemergence use in alfalfa grown for seed production, beans, clover grown for seed, corn, nonbearing foods crops, peanut, peas, peppermint, rice, sorghum, soybean, and spearmint

Active Ingredient:

sodium salt of bentazon* 53.0%

Other Ingredients: 47.0%

Total: 100.0%

* Equivalent to 5.0 pounds per gallon bentazon [3-(1-methylethyl)-1H,2,1,3-benzothiadiazin-4(3H)-one 2,2-dioxide] formulated as a soluble liquid

EPA Reg. No. 7969-112

EPA Est. No.

**KEEP OUT OF REACH OF CHILDREN
DANGER/PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID: If in eyes: Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes. Call a poison control center or doctor for treatment advice. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. **DO NOT** give anything to an unconscious person. **HOTLINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor or going for treatment. **In case of emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).** **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

See inside for complete **Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

Net Contents:

BASF Corporation
26 Davis Drive, Research Triangle Park, NC 27709

Precautionary Statements

Hazards to Humans and Domestic Animals

DANGER. Corrosive. Causes irreversible eye damage. **DO NOT** get in eyes or on clothing. Avoid contact with skin. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear (goggles, face shield, safety glasses)

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For terrestrial uses, **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Bentazon, the active ingredient in this product, is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

It is a violation of federal law to use any pesticide in a manner that results in the death of an endangered species or in adverse modification of their habitat.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in possession of the user at time of herbicide application.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Observe all directions, precautions, and restrictions in this label and the labels of products used in combination with **Basagran® 5L herbicide**. Use of **Basagran 5L** not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **48 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

DO NOT allow this product to freeze.

Pesticide Disposal

Wastes resulting from this product must be disposed of on-site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity $\leq$ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity $>$ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(continued)

STORAGE AND DISPOSAL (continued)

Container Handling (continued)

Refillable Container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

In Case of Emergency

In case of large-scale spill of this product, avoid contact, isolate area, and keep out animals and unprotected persons. Confine spill and call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment.
- Your local poison control center (hospital).
- BASF Corporation 1-800-832-HELP (4357)

Steps to take if this material is released or spilled:

- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

Product Information

Basagran® 5L herbicide is intended for selective postemergence control of certain broadleaf weeds and sedges in alfalfa grown for seed, beans, clover grown for seed, corn, nonbearing food crops, peanut, peas, peppermint, rice, sorghum, soybean, and spearmint.

Basagran 5L does not control grasses.

Mode of Action

Basagran 5L is a benzothiadiazinone herbicide and a photosynthesis PS II inhibitor belonging to the herbicide mode-of-action **Group 6** (WSSA)/**Group C₃** (HRAC).

Basagran 5L is effective mainly through contact action; therefore, weeds must be thoroughly covered with spray.

Crop Tolerance

All labeled crops are tolerant to **Basagran 5L**. Leaf speckling or bronzing may occur, but plants generally outgrow this condition within 10 days. New growth is normal and crop vigor is not reduced.

Application Instructions

Apply **Basagran 5L** postemergence to small, actively growing weeds early in the season for the most effective control as broadcast, band, or spot spray applications. Delaying application permits weeds to exceed the specified maximum size and will prevent adequate control.

Apply the specified rates of **Basagran 5L** to actively growing weeds before they reach the maximum sizes listed in **Table 1**, **Table 3**, and **Table 4**.

When applying **Basagran 5L** sequentially, wait a minimum of 5 days before re-treating.

Rainfast period - Rainfall or overhead irrigation within 4 hours after application may reduce the effectiveness of **Basagran 5L**.

Irrigation

In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth because weeds growing under drought conditions usually are not satisfactorily controlled.

Spray Coverage

Thorough spray coverage is required for optimum control of emerged weeds. Dense leaf canopies shelter smaller weeds and can prevent adequate spray coverage. Early season application to small weeds makes thorough spray coverage easier to obtain.

Cultivation

DO NOT cultivate within 5 days before applying or within 7 days after applying **Basagran 5L**. Timely cultivation after 7 days may help provide season-long weed control.

Application Methods and Equipment

Avoiding spray drift at the application site is the responsibility of the applicator. The spray system and weather-related factors determine the potential for spray drift. The applicator is responsible for considering these factors when making application decisions to avoid spray drift onto nontarget areas.

Aerial Application

Use a minimum spray volume of 5 gallons of water per acre. To obtain uniform coverage and to reduce the potential of spray drift to nontarget areas from aerial application, the following measures must be followed:

- **Boom Length** - The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the fixed wingspan or 90% of rotor-blade diameter.
- **Nozzle Type** - Use low-drift straight stream nozzles (D-8 or larger).
- **Nozzle Orientation** - Nozzles must always point backward parallel with the airstream and never be pointed downward more than 45 degrees.
- **Application Height** - Without compromising aircraft safety, apply at a height of 10 feet or less above the crop canopy or tallest plants.
- **DO NOT** apply when wind speed is greater than 10 miles per hour, or during periods of temperature inversions or stable atmospheric conditions.
- **DO NOT** apply **Basagran** by air if sensitive crop species (such as cotton, sugar beet, sunflower, or okra) are within 200 feet downwind.

Ground Application

Use a minimum spray volume of 10 gallons of water per acre to ensure adequate spray coverage. Use higher spray volume (up to 20 gallons of water per acre) to improve spray coverage when crop and weed foliage is dense. Apply using spray nozzles that deliver **medium-to-coarse spray droplets** as defined by ASABE standard S-572.1 and as shown in the nozzle manufacturer's spray catalogs. **DO NOT** use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles or selective application equipment such as recirculating sprayers or wiper applicators. **DO NOT** use brass nozzles because of the corrosive effects of nitrogen additives.

Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions, followed by triple rinsing the equipment before and after applying **Basagran 5L**.

Table 1. Application Rates and Weeds Controlled in All Crops Except Rice*

Weeds Controlled (includes ALS-resistant, glyphosate-resistant, and triazine-resistant biotypes)	Basagran® 5L herbicide (pints/A)					
	0.8**		1.2		1.6	
	Weed Leaf Stage	Maximum Height (inches)	Weed Leaf Stage	Maximum Height (inches)	Weed Leaf Stage	Maximum Height (inches)
Anoda, spurred	—	—	Up to 6	3	6 to 8	4
Balloonvine	—	—	2 to 4	2	4 to 6	3
Beggarticks	—	—	Up to 6	6	6 to 8	8
Bindweed, field ⁸	See the Special Directions for Problem Weeds section.					
Bindweed, hedge ⁸	See the Special Directions for Problem Weeds section.					
Buckwheat, wild	—	—	Up to 4	3	4 to 6	5
Canada thistle ⁸	See the Special Directions for Problem Weeds section.					
Cocklebur ^{1,8}	2 to 4	4	2 to 6	6	6 to 10	10
Croton, tropic	—	—	Up to 2	2	2 to 4	4
Dayflower	—	—	Up to 6	4	6 to 10	8
Devilsclaw ²	—	—	—	—	Up to 6	3
Eclipta	—	—	Up to 6	2	Up to 6	2
Galinsoga ²	—	—	—	—	Cotyledon to 6	2
Groundsel, common	—	—	—	—	—	3
Jimsonweed	Up to 4	4	Up to 6	6	6 to 10	10
Ladysthumb	Up to 4	4	Up to 6	6	6 to 10	10
Lambsquarters, common ^{2,3}	Up to 4	1	Up to 6	1.5	Up to 6	2
Marshelder	—	—	Up to 4	2	Up to 8	4
Mayweed/Dogfennel	—	—	—	2	—	3
Morningglory ⁵	—	—	4	4	6	6
Morningglory, cypressvine ⁵	—	—	4	4	4	4
Morningglory, smallflower ⁵	—	—	4	4	4	4
Mustard, wild	Up to 4	2	Up to 6	4	6 to 10	8
Nightshade, hairy ⁷	—	—	—	—	2 to 6	4
Nutsedge, yellow ⁸	See the Special Directions for Problem Weeds section.					
Poinsettia, wild ²	—	—	Up to 6	4	4 to 8	6
Purslane, common	—	—	Up to 4	1	4 to 6	2
Radish, volunteer	—	—	2 to 6	4	6 to 10	10
Ragweed, common ²	—	—	—	—	4 to 6	3
Ragweed, giant ³	—	—	—	—	Up to 4	6
Redweed	—	—	4 to 6	6	6 to 10	8
Senna, coffee ²	—	—	—	—	Up to 1 pinnate	2
Sesbania ²	—	—	—	—	3 to 5	3
Shepherdspurse ⁴	—	—	Up to 6	4	6 to 10	8
Sida, prickly (Teaweed)	—	—	Up to 6	3	6 to 8	4

(continued)

Table 1. Application Rates and Weeds Controlled in All Crops Except Rice* *(continued)*

Weeds Controlled (includes ALS-resistant, glyphosate-resistant, and triazine-resistant biotypes)	Basagran® 5L herbicide (pints/A)					
	0.8**		1.2		1.6	
	Weed Leaf Stage	Maximum Height (inches)	Weed Leaf Stage	Maximum Height (inches)	Weed Leaf Stage	Maximum Height (inches)
Smartweed, Pennsylvania	Up to 4	4	Up to 6	6	6 to 10	10
Starbur, bristly	—	—	Up to 4	2	4 to 6	3
Sugar beet, volunteer	—	—	2 to 4	—	4 to 8	—
Sunflower, wild	Up to 2	3	Up to 4	5	4 to 6	8
Velvetleaf ^{6,8}	Up to 4	2	Up to 4	2	4 to 6	5
Venice mallow	Up to 4	2	Up to 6	2	6 to 10	4

*For **Basagran 5L** use rates and weeds controlled in rice, refer to **Table 3. Application Rates and Weeds Controlled in Rice - Flooded Fields** and **Table 4. Application Rates and Weeds Controlled in Rice - Drained Fields** in the **Crop-specific Information** section.

** This rate not applicable in California.

¹ **DO NOT** treat earlier than leaf stage shown, and **DO NOT** count cotyledon leaves.

² Use crop oil concentrate (COC) or COC plus urea ammonium nitrate (UAN).

³ For regrowth or new germination, a second application of **Basagran 5L** may be necessary.

⁴ **DO NOT** treat rosette before seed stalk appears.

⁵ **Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, and Virginia** - Make a second application 5 to 14 days later. For all other states, apply 1.6 to 2.4 pints of **Basagran 5L** per acre to annual morningglory plants not larger than 4 true leaves. Control may be partial or inconsistent.

⁶ Always use UAN or ammonium sulfate (AMS) as spray additive.

⁷ **Basagran 5L** does not control black nightshade or Eastern black nightshade.

⁸ See the **Special Directions for Problem Weeds** section.

Special Directions for Problem Weeds

Field and Hedge Bindweed - Kentucky, Illinois, Indiana, Michigan, and Ohio. Make a single application of 1.6 to 2.4 pints of **Basagran 5L** per acre when weeds are up to a maximum of 10-inches tall for suppression only.

Canada Thistle - Apply 1.6 pints of **Basagran 5L** per acre when weeds are from 8-inches tall to the bud stage. Make a second application at the same rate 7 to 10 days later.

Yellow Nutsedge - Apply 1.2 to 1.6 pints of **Basagran 5L** per acre when weeds are up to a maximum of 8-inches tall. If regrowth occurs, make a second application at the same rate 7 to 10 days later. For control in California, apply 1.6 pints per acre and make second application 10 to 14 days later.

Treatment for Late Rescue of Cocklebur - Make a single application of 1.6 to 2.4 pints of **Basagran 5L** per acre to weeds up to 24-inches tall. For improved activity, apply 1.2 pints of **Basagran 5L** per acre initially, and repeat application 10 to 14 days later.

Treatment for Late Rescue of Velvetleaf - Make a single application of 2.4 pints of **Basagran 5L** per acre plus spray additives (an oil concentrate and UAN, refer to **Additives** section for more details) to weeds up to 12-inches tall. For improved activity, apply 1.2 pints of **Basagran 5L** per acre plus spray additives (an oil concentrate and UAN or AMS, refer to **Additives** section for more details), followed by a second application at the same rate in 5 to 7 days.

Additives

To achieve consistent and optimum weed control, one of or a combination of the following additives listed in **Table 2**.

Additive Rates must be added to the spray tank.

Additives may cause some leaf burn, but new growth is normal and crop vigor is not reduced. The potential for leaf burn is increased when relative humidity and temperature are high. When an adjuvant (or a specific adjuvant product, such as a drift control agent) is to be used with this product, BASF recommends the use of a Chemical Producers and Distributors Association (CPDA) certified adjuvant. See **Table 2. Additive Rates**. Applicators must use the additive(s) that will provide the best performance suited for their geography, target weed, and environmental conditions.

Table 2. Additive Rates

Additive/Adjuvant	Rate (per volume of spray solution)
Crop oil concentrate (COC) ¹	1 gal/100 gals (1% volume/volume [v/v])
Methylated seed oil concentrate (MSO) ^{1,2} (MSO adjuvant must contain at least 60% methylated seed oil. Poor performance may occur with adjuvants containing less than 60% methylated seed oil.)	1 gal/100 gals (1% v/v)
Nonionic surfactant (NIS) (Use NIS containing at least 80% active ingredient. Organosilicone surfactant may be used in place of NIS.)	1 to 2 quarts/100 gals (0.25% to 0.50% v/v)
Additive/Nitrogen Fertilizer ³	Rate (per volume of spray solution)
Ammonium sulfate (AMS) (use only spray-grade dry AMS)	8.5 to 17 lbs/100 gals (1% to 2% weight/volume [w/v])
Urea ammonium nitrate (UAN) ⁴ (recommended liquid fertilizers include 28% N, 32% N, or 10-34-0)	1.25 to 2.5 gals/100 gals (1.25% to 2.5% v/v)

¹ Petroleum-based or vegetable seed-based crop oil concentrate may be used. The oil concentrate must be nonphytotoxic, contain only EPA-exempt ingredients, provide good mixing quality in a jar test, and be successful in local experience. The exact composition of suitable products will vary; however, petroleum-oil or vegetable-oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils are more satisfactory than unrefined vegetable oils. To determine the suitability of oil concentrates with **Basagran® 5L herbicide**, conduct a jar test (see **Tank Mixing Information** section).

² MSO is recommended when weeds are under moisture or temperature stress.

³ AMS and UAN are not for use in California.

⁴ UAN may be added in place of other spray additives to improve control of cocklebur, devilscrow, Pennsylvania smartweed, velvetleaf, Venice mallow, wild mustard, and wild sunflower. **Basagran 5L** plus nitrogen solution will not provide adequate control of common ragweed and common lambsquarters. If these weeds or other weeds requiring oil concentrate are present in addition to velvetleaf, then oil concentrate should also be used.

Tank Mixing Information

Basagran 5L may be tank mixed with additives and/or one or more other pesticide products registered for use in a given crop according to the specific tank mixing instructions in this label and respective product labels if the product labels do not prohibit such mixing. The most restrictive labeling applies to tank mixes. Refer to the companion product label(s) to determine the specific use rates, crop growth stage, weeds controlled and weed growth stage. Always perform a mixing test to check the compatibility of **Basagran 5L** with all potential tank mix partners. See **Crop-specific Information** for more details.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Basagran 5L** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers not specified on this label.

Compatibility Test with Oil Concentrate

Before mixing components, always perform a compatibility jar test. Use only water from the intended source at the source temperature. For a spray volume of 20 gallons per acre, use 3.3 cups (800 mL) of water. For other spray volumes, adjust proportionately. Add 2 teaspoons each of herbicide and oil concentrate for each 1.6 pints per acre of label rate.

Add components in the following sequence, gently mixing between additions:

1. **Basagran 5L**
2. Tank mix product, if used
3. Oil concentrate

Cap jar, invert 10 cycles, let stand for 15 minutes.

Evaluate. An ideal tank mix combination will be uniform. The suitability of the oil concentrate is questionable if any of the following are observed:

- **Free oil at the surface** - Film or globules
- **Flocculation** - Fine particles which may be suspended in the liquid or found as a precipitated layer at the bottom of the jar
- **Clabbering** - Thickening texture (coagulated) resembling yogurt or a curd-like texture as with cottage cheese

Mixing Order

Maintain constant agitation throughout mixing and application until spraying is complete.

1. **Water** - Fill tank 1/2 to 2/3 full with clean water and start agitation.
2. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.

3. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
4. **Water-soluble additives** (including NIS, and nitrogen fertilizers such as AMS or UAN)
5. **Water-dispersible products** (such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
6. **Water-soluble products** (such as **Basagran® 5L herbicide**)
7. **Emulsifiable concentrates** (such as COC or MSO)
8. **Remaining quantity of water**

Application Restrictions

- **Maximum seasonal use rate** - **DO NOT** apply more than a **total of 3.2 pints** (2.0 pounds of bentazon) of **Basagran 5L** per acre, per season in all crops.
- **DO NOT** apply more than a **total of 2.0 pounds of bentazon** ai (from all sources) per acre, per season.
- **DO NOT** apply to weeds under stress such as lack of moisture, herbicide injury, mechanical injury, or cold temperatures, or unsatisfactory control may result.
- **DO NOT** apply to crops subjected to stress conditions such as hail damage, flooding, drought, injury from other herbicides, or widely fluctuating temperatures, or crop injury may result.
- **DO NOT** apply to crops that show injury (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications because this injury may be enhanced or prolonged.
- **DO NOT** apply through any type of irrigation system.

Crop-specific Information

This section provides use directions for **Basagran 5L** in specific crops. Read product information, mixing, application, weeds controlled, and additive instructions in the preceding sections of the label. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Alfalfa Grown for Seed

For use ONLY in Idaho, Montana, Nevada, Oregon, Washington, and Wyoming.

Seed alfalfa is tolerant to postemergence applications of **Basagran 5L**, however some leaf speckling, leaf bleaching or whitening, and temporary stunting may occur under certain conditions. Applications made at or after flower bud formation may reduce seed yields.

Apply **Basagran 5L** in the spring as a broadcast postemergence application at rates up to 1.6 pints per acre to alfalfa with at least 2 trifoliate leaves, but with no

flower bud formation. If needed for additional control of troublesome weeds, make a second application 5 to 14 days later at the same use rate.

The addition of oil concentrate (see **Additives** section for details) to **Basagran 5L** on seed alfalfa may result in slight leaf burn or temporary stunting, but all new growth is normal. The potential for leaf burn is increased when relative humidity and temperature are high. A few oil concentrates have exhibited excessive leaf burn. Refer to your supplier of **Basagran 5L** for information concerning successful local experience before purchasing any oil concentrate.

Crop-specific Restrictions

- **DO NOT** apply **Basagran 5L** if alfalfa shows injury (leaf phytotoxicity and/or plant stunting) produced by any other prior herbicide applications because this injury may be enhanced and/or prolonged.
- For use only in fields of alfalfa grown for seed production.
- **DO NOT** use on alfalfa produced for livestock feed.
- **DO NOT** use **Basagran 5L**-treated alfalfa seed for sprouting.

Special Crop Use Restrictions

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/nonfood use. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for alfalfa. Consequently, no portion of this alfalfa seed crop, including but not limited to, forage or stubble, green chop, hay, pellets, meal, whole seed, cracked seed straw, or seed screenings, may be used or distributed for human food or animal feed purposes.

Any alfalfa seed from a field treated with this product must bear a specific tag or conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading, with the following statement: "Not for human or animal consumption". All seed screenings from seed processing shall be disposed of in such a manner that the screenings cannot be distributed or used for human food or animal feed purposes.

Beans, Dry and Succulent

Bean types tolerant to **Basagran 5L** are adzuki, black turtle soup, cranberry, faba/fava, great northern, kidney, large lima, navy, pink, pinto, red, small lima, snap, and white beans.

Not for use on adzuki beans in California.

Apply **Basagran 5L** to beans after the first trifoliate leaf has fully expanded. Even at the tolerant stages, yellowing,

bronzing, speckling or burning of leaves may occur under certain conditions. This temporary injury is generally outgrown without delaying podset or maturity or reducing yield. Applying **Basagran® 5L herbicide** with an oil concentrate (COC or MSO) may increase crop injury (i.e. increase leaf burn) and may reduce yields, especially for snap bean.

State-specific Instructions for Georgia and South Carolina. **DO NOT** apply **Basagran 5L** as a solo treatment to dry and succulent beans and snap beans or severe crop injury may occur. **Basagran 5L** may be applied from 4.8 to 12.8 fluid ounces per acre to dry and succulent beans and snap beans only when tank mixed with **Raptor® herbicide**, **Pursuit® herbicide** or **Varisto® herbicide**. Refer to the **Raptor**, **Pursuit** and **Varisto** labels for additional use directions or restrictions.

Crop-specific Restrictions

- **DO NOT** apply **Basagran 5L** to beans before the first trifoliate leaf is fully expanded or severe crop injury may occur.
- **DO NOT** apply **Basagran 5L** to garbanzo beans or lupines at any stage of growth, or severe crop injury may occur.
- **DO NOT** apply **Basagran 5L** to dry or succulent beans within 30 days of harvest.

Tank Mixes

Basagran 5L may be tank mixed in beans with one or more of the following herbicide products. Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

- **Outlook® herbicide**
- **Poast® herbicide**
- **Pursuit**
- **Raptor**
- clethodim
- **Reflex® herbicide**
- **Varisto**

Basagran 5L may be tank mixed in succulent beans with one or more of the following herbicide products. Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

- **Poast**
- **Pursuit**
- **Raptor***
- clethodim
- **Reflex** (snap bean only)
- **Varisto**

*Only in states that **Raptor** is labeled for use on succulent lima bean and snap bean (see **Raptor** label)

Clover Grown for Seed

For use ONLY in Oregon and Washington.

Clover is tolerant to postemergence applications of **Basagran 5L**; however, some leaf burning may occur under certain conditions. Clover plants generally outgrow this condition within 10 days. Apply **Basagran 5L** with the appropriate additive and rate (see **Additives** section and **Table 2**).

Apply **Basagran 5L** in the spring as a broadcast postemergence application at rates up to 1.6 pints per acre. If needed for additional control of troublesome weeds, make a second application 5 to 14 days later at the same use rate.

Crop-specific Restrictions

- **DO NOT** graze livestock or harvest forage or hay for livestock feed for at least 36 days after treatment in Oregon and Washington.

Tank Mixes

Basagran 5L may be tank mixed in clover grown for seed with one or more of the following herbicide products. Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

- **Raptor**
- **Varisto**

Corn

Apply **Basagran 5L** postemergence to corn (corn in this label refers to field, sweet, popcorn, and corn grown for seed or silage). Before applying **Basagran 5L** to seed corn, verify the selectivity of **Basagran 5L** on your inbred line with your local seed company (supplier) to help avoid potential injury to sensitive inbreds.

Crop-specific Restrictions

- **DO NOT** graze treated corn fields for at least 12 days after the last treatment with **Basagran 5L**.

Tank Mixes

Basagran 5L may be tank mixed in corn with one or more of the following herbicide products. Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

- **Outlook**
- **Status® herbicide**
- atrazine
- glyphosate (e.g. **Roundup® herbicide**)

Peppermint and Spearmint

Peppermint and spearmint are tolerant to **Basagran® 5L herbicide**; however, some leaf burning may occur under certain conditions, such as when plants are growing very actively and have extensive new, succulent tissue. Mint plants generally outgrow this condition within 10 days.

Crop-specific Restrictions

- **DO NOT** apply **Basagran 5L** to peppermint or spearmint within 20 days of harvest.

Tank Mixes

Basagran 5L may be tank mixed in peppermint and spearmint with one or more of the following herbicide products. Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

- **Poast® herbicide**
- **Buctril® herbicide**
- **Sinbar® herbicide**
- **Stinger® herbicide**

Peas, Dry and Succulent

Pea types tolerant to **Basagran 5L** are Austrian winter, dry field, garden, English, marrowfat, and southern peas.

Not for use on blackeyed peas in California.

Apply **Basagran 5L** to peas after three pairs of leaves (or 4 nodes) are present. Even at the tolerant stages, yellowing, bronzing, speckling, or burning of leaves may occur under certain conditions. The addition of an oil-based additive, such as a HSOC, COC, or MSO, may increase the likelihood of crop injury. This temporary injury is generally outgrown without delaying podset or maturity or reducing yield.

In western irrigated areas, avoid applying **Basagran 5L** during prolonged periods (for 2 to 5 days) of cold weather (day temperature below 75° F and night temperature below 55° F) because weed control may be nullified.

State-specific Instructions for Georgia and South Carolina. **DO NOT** apply **Basagran 5L** as a solo treatment to dry and succulent peas or severe crop damage may occur. **Basagran 5L** may be applied from 4.8 to 12.8 fluid ounces per acre to dry and succulent peas only when tank mixed with **Raptor® herbicide**, **Pursuit® herbicide** or **Varisto® herbicide**. Refer to the **Raptor**, **Pursuit** and **Varisto** labels for additional use directions or restrictions.

Crop-specific Restrictions

- **DO NOT** apply **Basagran 5L** to dry peas within 30 days of harvest.
- **DO NOT** apply **Basagran 5L** to succulent peas within 10 days of harvest (30 days in California).

- **DO NOT** apply **Basagran 5L** to peas under stress from root rot.
- **DO NOT** apply **Basagran 5L** to garbanzo beans or to lupines at any stage of growth, or severe crop injury may occur.

Crop-specific Precautions

- Infurrow treatments of insecticides or nematicides may predispose the peas to injury from **Basagran 5L**.

Tank Mixes

Basagran 5L may be tank mixed in peas with one or more of the following herbicide products. Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

- **Pursuit**
- **Raptor**
- MCPA
- **Reflex® herbicide**
- **Thistrol® herbicide**
- **Varisto**

The tank mix of **Basagran 5L** and **Thistrol** is for use in the northeastern states of Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont, and in the mid-Atlantic states of Delaware, Maryland, Virginia, and in the Pacific northwestern states of Idaho, Oregon, and Washington. Apply this tank mix after the 3-leaf stage (4-node stage) of peas, but not later than 3 nodes before pea flowering.

NOTICE TO USER: Because of variability among pea cultivars and in application techniques, neither the manufacturers nor the sellers have determined whether or not the tank mix of **Basagran 5L** and **Thistrol** can be safely used on all pea crops under all conditions. Therefore, determine if the tank mix of **Basagran 5L** and **Thistrol** can be used safely before large-scale use.

For improved control of pigweed species and common lambsquarters, apply a tank mix of **Basagran 5L** and MCPA.

Tank Mix Restrictions

- **DO NOT** apply the tank mix to peas when temperatures exceed 90° F.
- **DO NOT** apply the tank mix to peas after pea flower buds appear.
- Tank mixes not applicable on succulent peas in California.

Tank Mix Precautions

- Other crops, in particular beans, cotton, grapes, tomatoes, and ornamental plants, may be severely injured by off-target spray drift of **Thistrol**.

Peanut

Apply **Basagran® 5L herbicide** from peanut cracking through pegging.

Crop-specific Restrictions

- **DO NOT** graze treated peanut fields for at least 50 days after the last **Basagran 5L** treatment.
- **Basagran 5L** treated peanut hay and forage may be fed to livestock.

Crop-specific Precautions

- Infurrow treatments of insecticides and nematicides may predispose peanuts to injury from **Basagran 5L**.

Tank Mixes

Basagran 5L may be tank mixed in peanut with one or more of the following herbicide products. Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

- **Cadre® herbicide**
- **Outlook® herbicide**
- **Poast® herbicide**
- **Pursuit® herbicide**
- 2,4-DB amine
- **Cobra® herbicide**
- **ET® herbicide/defoliant**
- **Gramoxone Inteon® herbicide**
- **Ultra Blazer® herbicide**

Apply the tank mix of **Basagran 5L** and **Gramoxone Inteon** at the ground crack stage of peanuts to control an early flush of weeds. Apply a second application up to 28 days after ground crack stage. Always add NIS at the recommended rates to the tank mix of **Basagran 5L** and **Gramoxone Inteon**. **DO NOT** use an oil concentrate or any other oil-based additive with the tank mix of **Basagran 5L** and **Gramoxone Inteon**.

Basagran 5L may be tank mixed with foliar fungicides such as **Headline® SC fungicide** and **Priaxor® fungicide**.

Tank Mix Restrictions

- **DO NOT** include UAN solution or AMS with the tank mix of **Basagran 5L**, **Poast**, and **Ultra Blazer**.
- **DO NOT** add oil concentrate, UAN, or any other additives to the tank mix of **Basagran 5L** and 2,4-DB.
- Use only amine formulations of 2,4-DB.
- Tank mixes not applicable in California.

Rice

For rice grown in the Mississippi Delta region, Texas, and the states throughout the southeastern US.

Apply **Basagran 5L** early postemergence in rice, at the use rates specified in and before weeds exceed the maximum size listed in **Table 3** and **Table 4**.

Rice straw may be fed to livestock.

Alternate Flooding Culture

Arkansas, Louisiana, Mississippi, and Texas. Weed growth stages generally correspond to rice that is tillering (stooling) and occur before the permanent flood.

Basagran 5L must be applied when there is no water on the field and 24 hours or more before flooding. If **Basagran 5L** cannot be applied until after flooding, see directions under **Continuous Flooding Culture**.

Continuous Flooding Culture

In states using continuous flooding culture, or when treating after the permanent flooding, treatment must be made only when weeds are above the surface of the water. Weeds submerged at the time of application will not be adequately controlled. For early treatment, water may be partly or completely drained to expose more weed growth to spray applications of **Basagran 5L**. **DO NOT** raise water level for at least 24 hours after application or unsatisfactory control may result. **DO NOT** use ground equipment to apply to flooded fields because splashing will wash **Basagran 5L** off weed leaf surfaces and ineffective control may result.

Crop-specific Restrictions

- **DO NOT** use **Basagran 5L** on rice fields in which the commercial cultivation of catfish or crayfish is practiced.
- **DO NOT** use water containing **Basagran 5L** residues from rice cultivation to irrigate crops used for food or feed unless **Basagran 5L** is registered for use on these crops.
- **DO NOT** apply more than 3.2 pints of **Basagran 5L** per acre per season whether one or two rice crops (including ratoon) are grown that season.

Tank Mixes

Basagran 5L may be tank mixed in rice with one or more of the following herbicide products. Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

- **Beyond® herbicide**
- **Facet® L Herbicide**
- **Newpath® herbicide**
- propanil
- **Command® herbicide**
- **Grasp® herbicide**
- **League™ herbicide**
- **Londax® herbicide**
- **Permit® herbicide**
- **Permit Plus® herbicide**
- **RiceBeaux® herbicide**
- **Strada® herbicide**
- **Ultra Blazer**

Tank Mix Restrictions

- Apply the tank mix of **Basagran® 5L herbicide** and **Londax® herbicide** within 7 days of establishing permanent flood.
- Apply the tank mix of **Basagran 5L** and propanil only to drained fields.
- **DO NOT** use oil concentrate with the tank mix of **Basagran 5L** and propanil. Add propanil to the tank mix of **Basagran 5L** based on active ingredient (ai) of formulation used. Test propanil products for physical tank mix compatibility with **Basagran 5L**.

Table 3. Application Rates and Weeds Controlled in Rice - Flooded Fields

Weeds Controlled	Basagran 5L Application Rates for Weed Growth Stages ¹ (pints/A)			
	1.2		1.6	
	Maximum Height Above Soil (inches)	Height Range Above Water Level (inches)	Maximum Height Above Soil (inches)	Height Range Above Water Level (inches)
Cocklebur	10	3 to 6	15	6 to 10
Dayflower	6	3 to 5	10	5 to 8
Flatsedge, annual or rice	See the Special Directions for Problem Weeds in Rice section.			
Redstem	4	2 to 3	8	4 to 6
Smartweed	6	2 to 5	10	5 to 8
Water plantain, arrowhead	—	—	7	5 to 6
Water plantain, common	—	—	7	5 to 6
Yellow nutsedge	6	4 to 5	10	6 to 8

¹ If a second weed flush develops after the first application, re-treat according to this rate table.

Table 4. Application Rates and Weeds Controlled in Rice - Drained Fields

Weeds Controlled	Basagran 5L Application Rates for Weed Growth Stages ¹ (pints/A)			
	1.2		1.6	
	Weed Leaf Stage	Maximum Height (inches)	Weed Leaf Stage	Maximum Height (inches)
Cocklebur	2 to 10	10	10 to 15	15
Dayflower	2 to 10	6	10 to 15	10
Ducksalad	—	—	6 to 10	6
Eclipta	4 to 6	2	4 to 6	2
Flatsedge, annual or rice	See the Special Directions for Problem Weeds in Rice section.			
Gooseweed	4 to 6	4	6 to 10	8
Redstem	Up to 6	4	6 to 10	8
Redweed	4 to 6	6	6 to 10	8
Smartweed	2 to 10	6	10 to 15	10
Spikerush	2 to 6	6	6 to 8	8
Water plantain, arrowhead	—	—	Up to 4	7
Water plantain, common	—	—	Up to 4	7
Yellow nutsedge	4 to 6	6	6 to 8	10

¹ If a second weed flush develops after the first application, re-treat according to this rate table.

Special Directions for Problem Weeds in Rice

Annual or Rice Flatsedge (*Cyperus iria*) - For suppression, apply 1.2 to 1.6 pints of **Basagran® 5L herbicide** per acre when weeds are up to a maximum of 8-inches tall. If regrowth occurs, make a second application at the same rate 7 to 10 days later.

Rice Grown in California

Apply **Basagran 5L** early postemergence in rice at the use rates specified in **Table 5**.

Rice straw may be fed to livestock.

Alternate Flood Culture

Weed growth stages generally correspond to rice that is tillering (stooling) and occur before the permanent flood. **Basagran 5L** must be applied when there is no water on the field and 24 hours or more before flooding. If **Basagran 5L** cannot be applied until after flooding, see directions under **Continuous Flood Culture**.

Continuous Flood Culture

In continuous flood culture, or when treating after the permanent flooding, treatment must be made only when weeds are above the surface of the water. Weeds submerged at the time of application will not be adequately controlled. For early treatment, water may be partly or completely drained to expose more weed growth to spray applications of **Basagran 5L**. **DO NOT** raise water level for at least 24 hours after application or unsatisfactory control

may result. Application of **Basagran 5L** with ground equipment to flooded fields can cause splashing that will wash **Basagran 5L** off weed leaf surfaces and ineffective control may result.

Crop-specific Restrictions

- **DO NOT** apply more than 3.2 pints of **Basagran 5L** per acre per season.
- **DO NOT** release paddy water from field if **Basagran 5L** has been applied during the growing season. Tail water may be recirculated within the field or other rice fields similarly treated with **Basagran 5L**. Draining water from any **Basagran 5L**-treated field during the growing season is not permitted.

Tank Mixes

Basagran 5L may be tank mixed in rice with, but not limited to, the following herbicide products. Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

- propanil

Tank Mix Restrictions

- Apply the tank mix of **Basagran 5L** and propanil only to drained fields.
- **DO NOT** use oil concentrate with the tank mix of **Basagran 5L** and propanil. Add propanil to the tank mix of **Basagran 5L** based on active ingredient (ai) of formulation used. Test propanil products for physical tank mix compatibility with **Basagran 5L**.

Table 5. Application Rates and Weeds Controlled in Rice Grown in California

Weeds Controlled	Basagran 5L Application Rates for Weed Growth Stages¹ (pints/A)			
	1.2		1.6	
	Application in Drained Fields Before Permanent Flooding			
	Weed Leaf Stage	Maximum Height (inches)	Weed Leaf Stage	Maximum Height (inches)
Ducksalad	—	—	6 to 10	6
Redstem	Up to 6	6	6 to 10	8
Weeds Controlled	Application to Flooded Fields			
	Maximum Height Above Soil (inches)	Height Range Above Water Level (inches)	Maximum Height Above Soil (inches)	Height Range Above Water Level (inches)
Redstem	4	2 to 3	8	4 to 6
Water plantain, common	—	—	7	5 to 6
Yellow nutsedge	6	4 to 5	10	6 to 8

¹ If a second weed flush develops after the first application, re-treat according to this rate table.

Sorghum, Grain and Forage

Apply **Basagran® 5L herbicide** postemergence to sorghum (sorghum in this label refers to grain and forage sorghum). Before applying **Basagran 5L** to sorghum, verify the selectivity of **Basagran 5L** on your inbred or hybrid line with your local seed company (supplier) to help avoid potential injury to sensitive inbreds and hybrids.

Not for use on forage sorghum in California.

Crop-specific Restrictions

- **DO NOT** apply more than 1.6 pints of **Basagran 5L** per acre per season in sorghum.
- **DO NOT** apply to sorghum that is heading or blooming.
- **DO NOT** graze treated sorghum fields for at least 12 days after the last treatment with **Basagran 5L**.

Tank Mixes

Basagran 5L may be tank mixed in sorghum with one or more of the following herbicide products. Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

- **Clarity® herbicide**
- **Facet® L herbicide**
- **Outlook® herbicide**
- atrazine

Soybean

Apply **Basagran 5L** postemergence to soybean; soybean is tolerant to **Basagran 5L** at all stages of growth. Slight leaf speckling and leaf bronzing may occur under certain conditions, but soybean generally outgrows these conditions within 10 days.

Crop-specific Restrictions

- **DO NOT** graze or cut treated soybean fields for forage or hay for at least 30 days after the last treatment of **Basagran 5L**.

Tank Mixes

Basagran 5L may be tank mixed in soybean with one or more of the following herbicide products. Other tank mix products may be used, provided the product is registered for the intended use. Always follow the most restrictive language on the tank mix products' labels.

- **Outlook**
- **Poast® herbicide**
- **Pursuit® herbicide**
- **Raptor® herbicide**
- 2,4-DB amine
- glyphosate (e.g. **Roundup® herbicide**)
- **Cobra® herbicide**
- **Flexstar® herbicide**
- **Reflex® herbicide**
- **Resource® herbicide**
- **Ultra Blazer® herbicide**
- **Varisto® herbicide**

Basagran 5L may be tank mixed with foliar fungicides and/or foliar insecticides (except malathion and **Sevin® herbicide**) if the application timings properly coincide.

The tank mixing of an insecticide with **Basagran 5L** may increase the potential for crop injury.

Tank Mix Restrictions

- **Basagran 5L, Ultra Blazer, and Poast.** Oil concentrate must be used with the tank mix of **Basagran 5L, Poast,** and **Ultra Blazer** in place of a spray surfactant.
- **Basagran 5L and 2,4-DB amine.** Use only amine formulations of 2,4-DB. Use no other adjuvant except UAN at 2 to 4 pints per acre with this tank mix. **DO NOT** apply more than 1 application of this tank mix per season. The use of this tank mix will cause soybean foliage injury (such as burning, bronzing, or crinkling) and may reduce yields. **DO NOT** use this tank mix on soybeans that show symptoms of disease such as *Phytophthora* root rot.
- **DO NOT** apply tank mixes in California.

Nonbearing Food Crops

Not for use on nonbearing food crops in California.

Basagran 5L may be used for selective postemergence weed control in the following nonbearing food crops: almonds, apples, apricots, avocados, blackberries*, blueberries, cherries, crabapples, dates, figs, grapes, grapefruit, lemons, limes, macadamias, nectarines, olives, oranges, peaches, pears, pecans, pistachios, plums, pomegranates, prunes, raspberries*, tangelos, tangerines, walnuts.

*Apply **Basagran 5L** at or before planting only.

Apply **Basagran® 5L herbicide** at 1.2 to 1.6 pints per acre as a postemergence directed spray away from the foliage of desired plants. **DO NOT** allow spray to contact green stems, bark, or foliage. A directed spray application should reduce the potential for leaf injury. However, some leaf speckling and leaf bronzing may occur under certain conditions. If needed, use a spray shield or wrap or cover the plants when spraying around very young trees or vines.

Crop-specific Restrictions

- **DO NOT** apply to nonbearing foods using aircraft or any aerial equipment that results in a broadcast spray application.
- **DO NOT** graze animals in treated orchards or fields.
- **DO NOT** use hay from treated areas for animal feed or bedding.
- **DO NOT** apply **Basagran 5L** to nonbearing food crops within one year of harvest.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Basagran, Beyond, Cadre, Clarity, Facet, Headline, Newpath, Outlook, Poast, Priaxor, Pursuit, Raptor, Rezult, Status, and Varisto are registered trademarks of BASF.

Buctril and Sevin are registered trademarks of Bayer Crop Science.

Cobra and Resource are registered trademarks, and **League** is a trademark of Valent USA Corp.

Command is a registered trademark of FMC Corporation.

ET is a registered trademark of Nichino America, Inc.

Flexstar, Gramoxone Inteon, and Reflex are registered trademarks of a Syngenta Group Company.

Grasp and Stinger are registered trademarks of Dow Agrosciences LLC.

Londax and Ultra Blazer are registered trademarks of United Phosphorus, Inc.

Permit and Permit Plus are registered trademarks of Nissan Chemical Industries, Ltd.

RiceBeaux is a registered trademark of RiceCo LLC.

RoundUp is a registered trademark of Monsanto.

Sinbar is a registered trademark of Tessenderlo Kerley, Inc.

Strada is a registered trademark of Isagro USA.

Thistrol is a registered trademark of Nufarm, Inc.

© 2016 BASF Corporation
All rights reserved.

007969-00112.20160502c.**NVA 2016-04-033-0088**
Supersedes: NVA 2014-04-033-0222

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry