



007969-00113-060499

Printed: 10:54:46 Friday, 25 Jun, 1999 # 47 / 3498

Systems Integration Group, Inc.

PM 25

7969-113

6/4/99

PS 1/2

Supplemental Label

lh - 4/8/99
NVA 98-4-82-0198
Vers. 1.3

Paramount®

herbicide

For use in Roadsides, Fence Lines, Rights-of-Ways and Non-crop Areas

EPA Registration Number 7969-113

All applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** on the EPA-registered label are to be followed. This labeling must be in the possession of the user at the time of application.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** are to be followed. This labeling must be in the user's possession during application.


In Case of Emergency

In case of large-scale spillage regarding this product, call:

CHEMTREC 800-424-9300
BASF Corporation 800-832-HELP

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment.
- Your local poison control center (hospital).
- BASF Corporation (800-832-HELP).



Application Instructions

Paramount® herbicide is a dry flowable herbicide for selective control or suppression of bindweeds, leafy spurge and certain other weeds in roadsides, fence lines, rights-of-ways and non-crop areas. **Paramount** is a herbicide with mainly systemic action. Uptake into the plant occurs through both the foliage and root system. Adequate soil moisture and/or light rain after application is important for soil uptake. Thorough coverage of foliage is important for consistent weed control.

Visual symptoms of weed control of **Paramount** may take more than three weeks following application to develop on perennial weeds (see **Table 1**). The full affect of **Paramount** may not be evident on perennial weeds for 3-6 months after application. For annual weeds, visual symptoms of weed control of **Paramount** may take up to two weeks following application to develop (see **Table 2**). These symptoms include initial twisting to stunting, reddening, and chlorosis about 14 days followed by necrosis and death about 21 days after application. **Paramount** may be applied to non-crop areas such as fence lines, roadsides, highway medians, utilities, railroad and pipeline rights-of-way. **Paramount** may be applied to non-cropland areas for the control of certain weeds in the Noxious Weed Control Programs, Districts or Areas including broadcast or spot treatments.

Apply **Paramount** to actively growing weeds. Refer to **Table 1. Application Rates and Timing - Perennial Weeds** and **Table 2. Application Rates and Timing - Annual Weeds** for recommended rates and timing.

Tank Mixing Information

Other registered products may be tank mixed with **Paramount**. Read and follow the applicable **Restrictions and Limitations and Directions For Use** on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

Table 1. Application Rates and Timing - Perennial Weeds

| Perennial Weeds | Stage or Timing | Rate Per Acre |
|---|---|---------------|
| Bindweed, field Bindweed, hedge | Fall application prior to frost, at least 4" long and growing | 8 ounces |
| Leafy Spurge | Suppression - apply at yellow bract (prebloom) or in fall prior to frost | 8 ounces |
| Sowthistle Thistle, Canada Thistle, Russian | Suppression - apply at rosette stage or bud stage; avoid application when seed stalk is bolting | 8 ounces |
| Clover | Small seeded legumes are controlled when they are actively growing | 5.3-8 ounces |

Table 2. Application Rates and Timing - Annual Weeds

| Annual Weeds | Maximum Size | Rate per Acre | Maximum Size | Rate per Acre |
|--|--------------|---------------|-----------------|---------------|
| Barnyardgrass Broadleaf signalgrass Foxtail, green | 4 leaves | 4 ounces | Up to tillering | 8 ounces |
| Foxtail, giant Foxtail, yellow | 2 leaves | 4 ounces | Up to tillering | 8 ounces |
| Crabgrass, large | - | - | 3 leaves | 8 ounces |
| Flax, volunteer | 5 inches | 4 ounces | 7 inches | 5.3 ounces |
| Bedstraw/cleavers | 3 whorls | 4 ounces | 5 whorls | 8 ounces |
| Morningglory | - | - | 3 leaves | 8 ounces |

All leaf stages listed above refer to true leaves.

Restrictions and Limitations

Do not allow **Paramount® herbicide** to contact or drift onto sensitive crops. Severe injury may result if this occurs.

Do not graze or harvest hay from treated areas.

Do not apply to water or to areas where surface water is present.

Do not apply to irrigation ditches or areas that act as a channel for water entering cropland.

Paramount is a registered trademark of BASF Corporation.

©BASF Corporation
All rights reserved
NVA 98-4-82-0198

BASF Corporation
P.O. Box 13528
Research Triangle Park, NC 27709

BASF