

Paramount®

herbicide

For weed control in fallow systems, grass grown for seed, pre-plant wheat, pre-plant and in-crop sorghum, and non-crop areas in the following states: CO, ID, KS, MO, MT, ND, NE, NM, NV, OK, OR, SD, UT, WA, WY, and designated counties in TX.

Active Ingredients:

3,7-Dichloro-8-quinolinecarboxylic acid

75.0%

Inert Ingredients:

25.0%

Total

100.0%

EPA Reg. No. 7969-113

EPA Est. No.: 65387-AR-001

Keep out of reach of children.

CAUTION

See inside label for complete **Precautionary Statements, Statements of Practical Treatment, Directions For Use, and Conditions of Sale and Warranty.**

ACCEPTED
APR 11 2002

7969-113

Net contents: Five 1-pound water-soluble bags

BASF Corporation

P.O. Box 13528, Research Triangle Park, NC 27709

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).</p>	

Precautionary Statements

Hazards to Humans and Domestic Animals

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or spray mist. Causes moderate eye injury. May cause allergic skin response.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Wash thoroughly with soap and water after handling. Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not apply directly to water, areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of rinsate.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions and precautions are to be followed. This labeling must be in the user's possession during application

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours**. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Protective eyewear
- Shoes plus socks

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Container Storage: Do not store box under wet conditions. Handle box carefully when stored at temperatures less than 50° F to avoid breaking the water-soluble bags. II.

Container Disposal: Completely empty box into application equipment. Then dispose of packaging in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burning, stay out of smoke.

Pesticide Disposal: Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.

In Case of Spill

In case of large-scale spillage regarding this product, call:

CHEMTREC 800-424-9300
BASF Corporation 800-832-HELP

I. General Information

Paramount® herbicide is intended for use in fallow systems, grass grown for seed, pre-plant wheat, pre-plant and in-crop sorghum, and non-crop areas. **Paramount** herbicide is formulated as a dry flowable designed for dilution with water and spraying in common agricultural spray equipment. It is packaged in pre-measured, one pound, water-soluble bags that are added to the mixing vat or directly into the spray tank. When used as directed, **Paramount** will provide suppression or control of weed species listed in **Table 1**.

Table 1. Target Weeds

Weeds Controlled	Weeds Suppressed*
<u>Annual Grasses</u>	<u>Annual Broadleaves</u>
Barnyardgrass	Kochia
Crabgrass, large	Lambsquarters,
Foxtail , giant	Common
, green	Ragweed ,common
, yellow	,giant
Signalgrass, broadleaf	Sunflower ,wild
	Thistle ³ ,Russian
<u>Annual Broadleaves</u>	Velvetleaf
Bedstraw, catchweed	
(cleavers)	<u>Perennial Broadleaf</u>
Clovers	Dandelion
Lettuce , prickly	Sowthistle ³ , perennial
Morningglory spp.	Spurge ² , leafy
Flax , volunteer	Thistle ³ , Canada
<u>Perennial Broadleaf</u>	
Bindweed ¹ , field	
, hedge	

* For improved control, add a tank mix partner that is active on listed species.

¹ Refer to the section entitled "**Field and Hedge Bindweed Control Recommendations**" for use directions.

² Use 8.0 – 16.0 ounces of **Paramount** per acre in non-crop areas for suppression and annual growth control, but do not exceed a total of 16.0 ounces of **Paramount** per acre per calendar year. Apply **Paramount** at yellow bract (pre-bloom) or in the fall prior to the first killing frost. For best performance on this species, tank mix 8.0

ounces per acre of **Paramount** with 4-6 ounces per acre of **Distinct**[®] herbicide.

³ Use 8.0 ounces of **Paramount** per acre for suppression and annual growth control, but do not exceed a total of 16.0 ounces of **Paramount** per acre per calendar year. Apply **Paramount** at rosette stage or bud stage. Avoid application when seed stalk is bolting. For best performance on this species, tank mix 8.0 ounces per acre of **Paramount** with 4-6 ounces per acre of **Distinct**[®] herbicide.


Field and Hedge Bindweed Control Recommendations

For most effective bindweed control, apply **Paramount** herbicide in the fall just prior to the first killing frost. Bindweed plants should be actively growing and at least 4 inches long. If tillage is a part of local post-harvest practices, allow a minimum of 30 days for bindweed plants to re-grow after tillage prior to application. For best long-term bindweed control, make yearly applications of **Paramount** at 5.3 – 8.0 ounces per acre in the fall. Use the higher rate for dense populations or large plants.

Paramount Application Area

For the application regions for **Paramount**, see Figure 1.

Figure 1. Application Region for Paramount



Paramount may be used in the following counties of Texas: Archer, Armstrong, Bailey, Baylor, Borden, Briscoe, Brown, Callahan, Carson, Castro, Chlidress, Clay, Cochran, Coke, Coleman, Collin, Collingsworth, Concho, Cooke, Cottle, Crosby, Dallam, Dawson, Deaf Smith, Denton, Dickens, Donley, Fisher, Floyd, Foard, Garza, Glasscock, Gray, Grayson, Hale, Hall, Hansford, Hardeman, Hartley, Haskell, Hemphill, Hockley, Hutchinson, Jack, Jones, Kent, King, Know, Lamb,

Lipscomb, Lubbock, Lynn, McCulloch, Montague, Moore, Motley, Nolan, Chiltree, Oldham, Parmer, Potter, Randall, Roberts, Runnels, Schackelford, Scurry, Sherman, Sterling, Stonewall, Swisher, Taylor, Terry, Throckmorton, Wheeler, Wichita, Wilbarger, Wise, Yoakum, and Young.

Be sure to obtain and follow all Texas state requirements for Paramount uses.

Mode of Action

Paramount is a systemic herbicide with plant uptake occurring through both the foliage and roots. Resultant herbicide symptoms on susceptible plants include twisting, stunting, reddening and chlorosis. For annual plants, symptoms may take up to two weeks after application to develop with death occurring in about three weeks. For perennial weeds, symptoms may not be evident for several weeks after application and full effect may not be evident for 3 to 6 months.

Coverage

When making postemergence applications, weeds must be thoroughly covered with spray because foliar uptake of **Paramount** by the target weed is important for optimum control. Large leaf canopies shelter smaller weeds and can prevent adequate spray coverage.

Cleaning Spray Equipment

Clean spray equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product.

II. Application Instructions

Based on the uses described in this label, **Paramount** should be applied by ground application equipment. **Paramount** may be applied using aerial application equipment only when permitted by supplemental labeling. The supplemental labeling must be in the possession of the user at the time of aerial application.

Paramount may be applied as either a broadcast or spot spray application. Applications must be made to actively growing weeds.

For most broadleaf weeds, the most effective control will result from applying **Paramount** early, when weeds are small. Delaying application permits weeds to exceed the maximum size and may prevent adequate control.

In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth.

Ground Application (Broadcast)

Water Volume: Use 5-30 gallons of water per broadcast acre. When weed foliage is dense, higher spray volumes may be required.

Spray Pressure: Use a maximum of 30 psi (measured at the boom, not at the pump or in the line).

Application Equipment: Use only nozzles that will produce uniform spray patterns and thorough coverage, spaced up to 20 inches apart. Select nozzles designed to produce minimal amounts of fine spray particles. Do not use controlled droplet applicator (CDA) nozzles as erratic coverage can cause inconsistent weed control. Do not use selective application equipment such as recirculating sprayers or wiper applicators. Drift reduction nozzles such as Delavan® Raindrop Drift Reduction Flat Spray Tip, RF Tips, XR Tee Jet™ Extended range Flat Spray Tips, or other brands of comparable capabilities are recommended.

A methylated seed oil or crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:

- be non-phytotoxic,
- contain only EPA-exempt ingredients,
- provide good mixing quality in the jar test, and
- be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils.

For additional information, see **Compatibility Test for Mix Components**.

For bindweed control in Oklahoma, New Mexico and the designated counties of Texas, the use of methylated seed oil plus AMS is mandatory with Paramount when it is applied alone.

Nitrogen Fertilizer Source:

• **Urea ammonium nitrate (UAN):** Commonly referred to as 28%, 30%, or 32% nitrogen solution. Do not use brass or aluminum nozzles when spraying UAN.

• **Ammonium sulfate (AMS):** AMS may be substituted for UAN. Use high-quality AMS (spray grade) to avoid plugging of nozzles. Other sources of nitrogen are not as effective as those mentioned. BASF does not recommend applying AMS if applied in less than 10 gallons per acre because of potential problems with precipitation in reduced volumes. Use AMS only if it has been demonstrated to be successful in local experience. Because most nitrogen solutions are mildly corrosive to galvanized, mild steel, and brass spray equipment, rinse the entire spray system with water soon after use. Use high-quality AMS to avoid plugging spray nozzles. The AMS must be readily soluble in water and contain no insoluble materials. Local sources of high-quality, fine, feed-grade AMS may be better than fertilizer grade. Low-quality AMS may contain material that will not readily dissolve, which could result in nozzle tip plugging. To determine AMS quality, perform a jar test adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If any un-dissolved sediment is observed, pre-dissolve the AMS in water and filter before adding it to the spray tank. If the AMS is added directly to the spray tank, add slowly while agitating. Adding the mix to quickly may clog outlet lines.

III. Spray Additives

To achieve consistent weed control, the use of spray additive(s) with **Paramount** is required. The recommended spray additive with **Paramount** is methylated seed oil. The use of crop oil concentrate with **Paramount** is also permitted. A nitrogen fertilizer source (AMS or UAN) can be added to enhance efficacy, but cannot be used in place of methylated seed oil or crop oil concentrate. Refer to **Table 2. Spray Additive Rate Per Acre** for spray additive rates.

Table 2. Spray Additive Rate Per Acre

Spray Additive	Ground Application
Methylated Seed Oil	1.0 – 2.0 pints ²
Crop Oil Concentrate	2.0 pints
AMS ¹	2.5 pounds
UAN solution ¹	2.0 – 4.0 quarts

¹ optional
² For best grass control, use at least 1.5 pints per acre of methylated seed oil

Methylated Seed Oil or Crop Oil Concentrate:

Nonionic Surfactant:

Alternatively, an 80% active nonionic spray surfactant may only be used when **Paramount** is tank mixed with other products that restrict the use of oil additives. However, the use of nonionic surfactant may result in

reduced weed control with **Paramount**. The standard label recommendation for nonionic surfactant is 1 quart per 100 gallons of water (0.25% vol./vol.). Applications with nonionic surfactant require the addition of a nitrogen fertilizer source.

IV. General Tank Mixing Information

Read and follow the applicable **Restrictions and Limitations** and **Directions For Use** on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

Tank Mix Partners/Components

Use the following tank mixes to achieve control of the weeds listed as suppressed in **Table 1**. The following herbicides may be tank mixed with **Paramount** herbicide according to the specific tank mixing instructions in this label and respective product labels. For all recommended tank mixes, use a rate of 5.3 - 8.0 ounces per acre of **Paramount**.

BASF does not recommend using tank mixes other than those listed on BASF labeling. Physical incompatibility, reduced weed control, or crop injury may result from mixing **Paramount** with other pesticides, additives, or fertilizers. Local agricultural authorities may be a source of information when using other than BASF recommended tank mixes.

- **2,4-D**
- **Atrazine**
- **Buctril®** (bromoxynil)
- **Buctril®/Atrazine** (bromoxynil + atrazine)
- **Clarity®** (dicamba)
- **Cyclone®** (paraquat)
- **Distinct®** (diflufenzopyr + dicamba)
- **Fallowmaster®** (glyphosate dicamba)
- **Frontier®** (dimethenamid)
- **Guardman® Max** (dimethenamid-P + atrazine)
- **Landmaster®** (glyphosate + 2,4-D)
- **Marksman®** (dicamba + atrazine)
- **Outlook™** (dimethenamid-P)
- **Peak®** (prosulfuron)
- **Roundup® RT** (glyphosate)
- **Roundup® Ultra** (glyphosate)
- **Weedmaster®** (dicamba + 2,4-D)

Compatibility Test for Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

- 1) **Water:** For 20 gallons per acre spray volume, use 3-1/3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
- 2) **Products in PVA bags:** (such as **Paramount®** herbicide) Cut an opening in the water-soluble PVA bag just large enough to use a teaspoon for measuring

purposes. Use the opened water-soluble PVA bag first when preparing spray solution. For **Paramount** at the 5.3 ounce rate, use 1 teaspoon; for the 8.0 ounce rate, use 1.5 teaspoons. Cap the jar and invert 10 cycles.

- 3) **Water-dispersible products** such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions.
- 4) **Water-soluble products:** Cap the jar and invert 10 cycles.
- 5) **Emulsifiable concentrates:** (methylated seed oil or crop oil concentrate when applicable). Cap the jar and invert 10 cycles.
- 6) **Water-soluble additives** (AMS or UAN when applicable): Cap the jar and invert 10 cycles.
- 7) Let the solution stand for 15 minutes.
- 8) **Evaluate** the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. Do not use any spray solution that could clog spray nozzles.

Mixing Order

- 1) **Water.** Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
- 2) **Agitation.** Maintain constant agitation throughout mixing and application.
- 3) **Inductor.** If an inductor is used, rinse it thoroughly after each component has been added.
- 4) **Products in PVA bags:** (such as **Paramount**). Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
- 5) **Water-dispersible products** (such as **Paramount**, dry flowables, wettable powders, suspension concentrates, or suspo-emulsions).
- 6) **Water-soluble products.**
- 7) **Emulsifiable concentrates** (such as oil concentrate when applicable).
- 8) **Water-soluble additives** (such as AMS or UAN when applicable).
- 9) **Remaining quantity of water.** Maintain constant agitation during application.

V. General Restrictions and Limitations

- **Maximum seasonal use rate:** Do not apply more than a total of 16 ounces of **Paramount®** herbicide per acre, per calendar year.
- **Restricted Entry Interval (REI): 12 hours.**
- **Crop Rotation Restrictions:** In case of crop failure, only Spring or Winter wheat and grain sorghum may be immediately replanted. Do not plant any other crop other than Spring or Winter wheat or grain sorghum for 309 days (10 months) following application. For alfalfa, clover, dry beans, flax, peas, lentils, safflower, Solanaceous crops listed below, sunflower, and sugar beet, do not replant for 24 months and conduct a bioassay prior to planting any of these crops.
- Do not apply to weeds or grasses under stress due to lack of moisture, herbicide injury, mechanical injury or cold temperatures, as unsatisfactory control may result.
- Do not apply to crops subjected to stress conditions such as hail damage, flooding, drought, injury from other herbicides, or widely fluctuating temperatures, as crop injury may result.
- Do not use selective application equipment such as re-circulating sprayers, wiper applicators, or shielded applicators.
- **Rain fast period: Paramount** is rain fast 6 hours after application.
- **Wind Speed for Ground Applications:** Do not apply **Paramount** when wind is blowing more than 10 mph.
- Do not apply **Paramount** using aerial equipment unless permitted by supplemental labeling.
- Do not apply through any type of irrigation equipment.

Drift:

- Do not allow **Paramount** to drift onto other desirable plants, especially sensitive crops belonging to the following plant families:
 1. Solanaceae [tomato, potato, tobacco, eggplant, peppers (Capsicum), among others]
 2. Umbelliferae (celery, parsley, carrots, among others)
 3. Leguminosae (alfalfa, green bean, among others)
 4. Convolvulaceae (sweet potato, among others)
 5. Chenopodiaceae (spinach, sugar beet, among others)
 6. Malvaceae (okra, among others)
 7. Cucurbitaceae (watermelon, cantaloupe, squash, pumpkin, among others)
 8. Compositae (lettuce, sunflowers, among others)
 9. Linaceae (flax)
- Do not allow spray containing **Paramount** to drift onto areas where tomatoes are to be planted, have been planted, or onto emerged tomatoes as severe injury will occur.
- Do not use **Paramount** in tank mixes not specified on this label or BASF technical bulletins.
- Do not premix **Paramount** with fungicides, herbicides, insecticides, additives, or fertilizers as contamination of mixing equipment and movement of **Paramount** to off-site mixing areas can occur

VI. Crop Specific Information

Table 3. Use Rate of Paramount per Package

Use rate of Paramount (ounces/acre)	Acres treated with a single 1-pound water soluble bag of Paramount
5.3	3
8.0	2

Grass Grown For Seed

For use in the following grasses grown for seed:

Cool Season Grasses:

- Bromegrass; smooth, meadow, smooth X meadow cross
- European Dunegrass
- Fescue; fine, tall
- Junegrass
- Kentucky Bluegrass
- Quackgrass
- Needlegrass; green
- Orchardgrass
- Ryegrass; annual, indian, perennial
- Wheatgrass; bluebunch, crested, fairway, fairway X crested cross, intermediate, pubescent, siberian, slender, tall, thickspike, western, bluebunch X quack cross
- Wildrye; altai, basin, beardless, dahurian, mammoth, russian

Warm Season Grasses:

- Bermudagrass
- Bluestem; big, little, sand
- Gramma; blue, side-oats
- Sandreed; prairie
- Switchgrass

Apply **Paramount** at 5.3 ounces per acre for control of annual grasses and broadleaf weeds (see **Table 1**). Apply **Paramount** for bindweed control after grass seed harvest and hay removal but before the first killing frost. Refer to the section entitled "**Field and Hedge Bindweed Control Recommendations**" for use directions.

Tank Mixing Information:

Other registered products may be tank mixed with **Paramount**. Read and follow the applicable **Restrictions and Limitations** and **Directions For**

Use on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

Fallow Systems or Pre-plant Wheat or Pre-plant Sorghum

Paramount can be applied in fallow areas or pre-plant wheat or pre-plant grain sorghum at 5.3 ounces per acre for control of annual grasses and broadleaf weeds (see **Table 1**). For bindweed control with **Paramount**, refer to the section entitled "**Field and Hedge Bindweed Control Recommendations**" for use directions.

Fallow Tank Mixes:

Other registered products may be tank mixed with **Paramount**. Read and follow the applicable **Restrictions and Limitations** and **Directions For Use** on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

- 2,4-D
- Clarity® (dicamba)
- Distinct® (diflufenzopyr + dicamba)
- Fallowmaster® (glyphosate + dicamba)
- Landmaster® (glyphosate + 2,4-D)
- Roundup® RT (glyphosate)
- Roundup® Ultra (glyphosate)

In-Crop Sorghum

Apply **Paramount** to grain sorghum at 5.3 - 8.0 ounces per acre from preemergence to postemergence (to 12 inch tall sorghum) for control of annual grasses and broadleaf weeds. For best annual grass control, **Paramount** should be applied at 5.3 - 8.0 ounces per acre in a tank mix with atrazine at 0.5-1.0 pound a.i. per acre when weeds are less than 2" tall. Do not use liquid fertilizer as a carrier for postemergence applications of **Paramount** to grain sorghum.

In Oklahoma, New Mexico, and in the designated counties in Texas, apply only 8.0 ounces of **Paramount** per acre to in-crop sorghum.

Table 4. Tank mixes with Paramount

Herbicide Tankmix Partner	Fallow and Pre-plant Wheat	Pre-plant Sorghum	Postemerge Sorghum
2,4-D	0.375-1.0 lb ai	0.375-1.0 lb ai	0.125-0.5 lb ai
Atrazine	-----	0.5-1.0 lb ai	0.5-1.0 lb ai
Clarity®	4-16 oz	4-16 oz.	8 oz
Fallowmaster®	22-44 oz	22-44 oz	-----
Landmaster®	32-54 oz	32-54 oz	-----
Peak®	-----	-----	0.25 oz
Roundup® Ultra and RT®	12-32 oz	12-32 oz	-----
Buctril®	-----	-----	16 oz
Buctril® / Atrazine	-----	-----	32 oz
Guardsman Max®	-----	-----	40 - 64 oz

Crop Specific Restrictions and Limitations

Do not allow livestock to graze in treated areas.

Do not harvest hay from treated areas within 309 days after application.

Do not feed treated grasses, forage, hay, silage, straw, seed nor seed screenings to livestock.

Do not apply to water or to areas where surface water is present.

Do not apply to irrigation ditches or areas that act as a channel for water entering cropland.

Noncrop Areas

(Roadsides, Fencelines and Rights-of Ways)

Paramount may be applied to non-crop areas such as fence lines, roadsides, highway medians, utilities, railroad and pipeline rights-of-way. **Paramount** may be applied to non-cropland areas for the control of certain weeds in the Noxious Weed Control Programs, Districts or Areas including broadcast or spot treatments. Use 5.3 – 8.0 ounces of **Paramount** per acre for control of annual weeds, or 8.0 – 16.0 ounces per acre for other perennial weeds (see **Table 1**), but do not exceed a total of 16.0 ounces of **Paramount** per acre per calendar year. For bindweed control with **Paramount**, refer to the section entitled “**Field and Hedge Bindweed Control Recommendations**” for use directions.

Non-Crop Tank Mixes:

Other registered products may be tank mixed with **Paramount**. Read and follow the applicable **Restrictions and Limitations** and **Directions For Use** on all products involved in tank mixing. The most restrictive labeling applies to tank mixes.

- **2,4-D**
- **Clarity®** (dicamba)
- **Distinct®** (diflufenzopyr + dicamba)
- **Roundup® RT** (glyphosate)
- **Roundup® Ultra** (glyphosate)

Crops:

This product can be used on the following crops:

- Fallow Systems
- Grass Grown for Seed
- Grain Sorghum
- Wheat (pre-plant)

Look inside for complete **Restrictions and Limitations and Application Instructions.**

Weeds listed in this label:

Common Name	Scientific Name
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bedstraw/cleavers	<i>Galium aparine</i>
Bindweed, field	<i>Convolvulus arvensis</i>
, hedge	<i>Calystegia sepium</i>
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>
Clover, crimson	<i>Trifolium incarnatum</i>
, red	<i>Trifolium pratense</i>
, white	<i>Trifolium repens</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Dandelion	<i>Taraxacum officinale</i>
Foxtail, giant	<i>Setaria faberi</i>
, green	<i>Setaria glauca</i>
, yellow	<i>Setaria viridis</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, common	<i>Chenopodium album</i>
Morningglory	<i>Ipomea spp.</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
, giant	<i>Ambrosia trifida</i>
Sunflower, wild	<i>Helianthus annuus</i>
Thistle, Canada	<i>Cirsium arvense</i>
, Russian	<i>Salsola iberica</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Volunteer Flax	<i>Linum sp.</i>

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. **BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.** BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Product of U.S. and Korea; formulated in the U.S. with U.S. and imported ingredients.

Clarity, Distinct, Frontier, Guardsman Max, Marksman, Paramount, and Weedmaster are registered trademarks and Outlook is a trademark of BASF Corporation. Buctril is a registered trademark of Rhone-Poulenc. Cyclone is a registered trademark of Zeneca Inc. Peak is a registered trademark of Novartis Corporation. Fallowmaster, Landmaster, and Roundup are registered trademarks of Monsanto Company. Delavan is a registered trademark of Garlock International Tee Jet is a trademark of Spraying Systems Company.

BASF Corporation
P.O. Box 13528
Research Triangle Park, NC 27709



© 2002 BASF Corporation
All rights reserved
NVA 2002-04-082-0002
Revised 1-10-02