

7969-130

09/25/2006

1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

SEP 25 2006

Ms. Rebecca Johnston
BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709-3528

Subject: Drive 75 DF Herbicide
EPA Registration Number 7969-130
Application dated August 2, 2006

Dear Ms. Johnston:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable, provided you make the following changes before you release the product for shipment.

- 1) Add an appropriate EPA Establishment Number to the label.
- 2) Revise the PRECAUTIONARY STATEMENTS to the following: "Harmful if swallowed, absorbed through skin, or inhaled. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation. Avoid breathing dusk or spray mist. May cause allergic response."
- 3) In the section STORAGE AND DISPOSAL, revise the subheading "Storage" to "Pesticide Storage."
- 4) On page 4, revise "recommendations" to "instructions" in the subsection "Additives in Spray Mix to Achieve Control."
- 5) On page 5, state the active ingredient equivalent to 2 lbs product in the restriction "DO NOT apply more than 2 pounds of Drive 75 DF per acre (or 0.73 oz per 1000 sq ft) in one year [X lbs a.i. per acre per year]." State the active ingredient equivalent throughout the label anywhere a product rate is listed.
- 6) On page 10, in the section CONDITIONS OF SALE AND WARRANTY, revise "should" to "must" in the first paragraph. Add "To the extent consistent with applicable law" before the statements "All such risks shall be assumed by the Buyer" and "BASF makes no other express or implied warranty of fitness or merchantability or any other express or implied warranty." Revise "To the extent permitted by law" to "To the extent consistent with applicable law" in the third paragraph.

2/12


Page 2

EPA Registration Number 7969-130

Submit one (1) copy of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling will supercede all previously accepted ones. A stamped copy of labeling is enclosed for your records.

If you have any questions, please contact Hope Johnson at 703-305-5410.

Sincerely,



James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505P)

3/12



The Chemical Company

DRIVE®

75 DF HERBICIDE

Active Ingredient:

quinclorac: 3,7-dichloro-8-quinolinecarboxylic acid 75.0 %

Inert Ingredients: 25.0 %

Total: 100.0 %

U.S. Patent No. 4,497,651

EPA Reg. No. 7969-130

EPA Establishment Number _____

KEEP OUT OF REACH OF CHILDREN.

CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

See inside booklet for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

ACCEPTED
with COMMENTS
in EPA Letter Dated

SEP 25 2006
Under the Federal Insecticide,
Fungicide and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

Net contents: _____

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

7969-130

FIRST AID

If swallowed

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- **DO NOT** induce vomiting unless told to do so by a poison control center or doctor.
- **DO NOT** give anything by mouth to an unconscious person.

If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If in eyes

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If inhaled

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

Note to physician: Probable mucosal damage may contraindicate the use of gastric lavage. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Causes moderate eye injury. May cause allergic skin response.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. For more options, refer to **Category A** on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as butyl rubber ≥ 14 mils, or natural rubber ≥ 14 mils, or neoprene rubber ≥ 14 mils, or nitrile rubber ≥ 14 mils
- Shoes plus socks

Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Keep out of lakes, ponds and streams. **DO NOT** apply directly to water, areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water by cleaning of equipment or disposal of rinsate.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with this labeling.

For use only by certified commercial applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

All applicable directions, restrictions and precautions are to be followed. This labeling must be in the possession of the user at time of application.

DO NOT apply this product by air or through any type of irrigation equipment.

BASF Corporation does not recommend or authorize the use of this product in manufacturing, processing or preparing custom blends with other products for application to turf.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the specified area during application.

For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, and water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material (such as nitrile, butyl, neoprene, and/or barrier laminate)
- Shoes plus socks

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouses.

Professional applications to residential and nonresidential turf are not within the scope of the Worker Protection Standard.

- **DO NOT enter or allow others to enter the treated area until sprays have dried.**

Storage and Disposal

DO NOT contaminate water, food, or feed by storage or disposal.

Storage: DO NOT store box under dampness or high humidity.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

Container Disposal: Dispose of empty packaging in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

In Case of Emergency

In case of large-scale spillage regarding this product, call:
 CHEMTREC 1-800-424-9300
 BASF Corporation 1-800-832-HELP (4357)

I. General Information

Drive® 75 DF herbicide may be applied postemergence to residential and nonresidential turfgrasses (Refer to **Table 1**) for the control of many broadleaf and grass weeds. Examples of such sites include, but are not limited to: grounds or lawns around residential and commercial establishments, multi-family dwellings, military and other institutions, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses and sod farms.

Drive 75 DF is absorbed by foliage and roots and translocated throughout the plant. The control symptoms exhibited by broadleaf weeds include leaf and stem curl or twisting, and chlorosis. Susceptible grasses demonstrate stunting, chlorosis, and gradual reddening followed by necrosis and death. Refer to **Tables 1, 2 and 3** for turfgrass tolerance and susceptible weed species.

II. Application Information

Apply Drive 75 DF to actively growing weeds as postemergence broadcast or spot sprays using the rate, turf species and growth stage indicated in **Tables 1, 2 and 3**. **DO NOT** exceed the labeled application rate or fail to comply with use restrictions listed in **VII. General Restrictions and Limitations**.

To achieve consistent weed control, use methylated seed oil. Refer to **Tables 1, 2 and 3** for rates.

Adding adjuvants may cause slight leaf burn, but new growth is normal, and turf vigor is not reduced. Avoid potential for leaf burn and turfgrass damage by delaying application when relative humidity and temperatures are high. Additional stress from low mowing heights may also increase the possibility of turf injury. Addition of chelated iron or sprayable soluble nitrogen fertilizer will reduce a slight yellowing that may occur on some turfgrass species.

Spot Applications: Postemergence spot applications may be made to susceptible weeds in turfgrass that is tolerant to **Drive 75 DF** (see **Tables 1 and 2**). Apply 0.367 ounces of **Drive 75 DF** per 1000 square feet of treated area. Spray coverage should be uniform and complete. See **Table 5** for **Spot Spray Mix Instructions**.

Mowing Information: DO NOT mow 2 days before or after applying **Drive 75 DF** to maximize weed control and minimize potential turf injury. **Clippings from the first three mowings should be left on the treated area.**

6/12

Irrigation and Rainfall: If soil moisture is not sufficient prior to **Drive® 75 DF herbicide** application, irrigation may improve weed control. For best results, **DO NOT** water or irrigate for 24 hours after application. If rainfall does not occur in 2 to 7 days after application, irrigation of at least one-half inch is required.

Extended Grass Control: To extend grass control, **Drive 75 DF** can be tank mixed with **Pendulum®** or **PRE-M® herbicide** to provide residual control of annual grasses. Consult the respective tank mix labels for additional weeds controlled.

Seeding/Overseeding/Sprigging:

The use of **Drive 75 DF** before or after seeding or overseeding a turf area will not significantly interfere with the turfgrass seed germination and growth of those grass types identified as tolerant or moderately tolerant in **Table 1**. Consult **Table 4** for timing of applications concerning any seeding, overseeding or sprigging situation.

Note: Not all bermudagrass seed have similar germination characteristics. Check germination vigor of your seeded hybrid bermudagrass prior to use of **Drive 75** before, at and 7 days after seeding.

III. Addition of Adjuvants

Additives in Spray Mix to Achieve Control:

Methylated seed oil is the preferred adjuvant for postmergence applications. **DO NOT** include additives when tank mixing with EC (Emulsifiable Concentrate) products as this may cause phytotoxicity. Consult your local BASF representative or distributor for recommendations for your area. *Other additives or adjuvants may be used, such as **DyneAmic® Lesco Spreader Sticker®, LI-700®, Surf King® Spreader, Target Pro® Spreader or Thoroughbred®**. However, under some environmental conditions, some of these products may cause phytotoxicity or less than optimal efficacy.*

Methylated Seed Oil:

The methylated seed oil used as the adjuvant with **Drive 75 DF** must meet all the following criteria:

- be nonphytotoxic,
- contain only EPA-exempt ingredients,
- provide good mixing quality in the jar test, and
- be successful in local experience.

The exact composition of suitable products will vary; however, any methylated seed oil used should contain emulsifiers to provide good mixing quality.

IV. Mixing Instructions for Drive® 75 DF herbicide

Respective sections of this leaflet define **Drive 75 DF** tank mix treatments. **If tank mixing with products in PVA bags, always allow water-soluble packets to dissolve before agitating.**

Ground Driven Sprayer:

1. Begin with a clean spray tank. Fill tank one-half to three-quarters full with clean water.
2. **Drive 75 DF** alone:
When using **Drive 75 DF** alone, either premix (1 part product + 2 parts water) or slowly add **Drive 75 DF** to the partially filled tank while agitating until completely dispersed. Fill the remainder of the tank with water.
3. **Drive 75 DF** tank mixes:
 - a. Wettable Powder Formulations:
 - 1) When using **Drive 75 DF** with a wettable powder formulation, add **Drive 75 DF** to the partially filled tank while agitating.
 - 2) When **Drive 75 DF** is properly mixed, add a slurry of the wettable powder (1 part WP + 2 parts water) slowly into the tank. Fill the remainder of the tank while agitating.
 - b. Flowable Formulations:
 - 1) When using **Drive 75 DF** with a flowable formulation, add **Drive 75 DF** to the partially filled tank while agitating.
 - 2) When **Drive 75 DF** is properly mixed, premix one part of the flowable with two parts water and add this diluted mixture slowly into the tank. Fill the remainder of the tank while agitating.
 - c. Emulsifiable Concentrate Formulations:
 - 1) When using **Drive 75 DF** with an emulsifiable concentrate formulation, add **Drive 75 DF** to the partially filled tank while agitating.
 - 2) When **Drive 75 DF** is properly mixed, premix one part of the emulsifiable concentrate with two parts water and add this diluted mixture slowly into the tank. Fill the remainder of the tank while agitating.
 - d. Dry Flowable (Water Dispersible Granule) Formulations:
 - 1) When using **Drive 75 DF** with a dry flowable formulation, add **Drive 75 DF** to the partially filled tank while agitating.
 - 2) When **Drive 75 DF** is properly mixed, premix one part of the dry flowable with two parts water and add this diluted mixture slowly into the tank. Fill the remainder of the tank while agitating.

4. MAINTAIN CONTINUOUS AGITATION WHILE ADDING HERBICIDES AND UNTIL SPRAYING IS COMPLETED. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.

Backpack Sprayer: Begin with a clean spray tank. Fill the spray tank ½ full with clean water and add the required amount of **Drive 75 DF** to the sprayer. Cap sprayer and agitate to ensure mixing. Uncap sprayer and finish filling tank to desired level. Cap sprayer and agitate once again. During application it is desirable to agitate the mixture on occasion to ensure mixing. If the mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed.

Liquid Fertilizers: Prior to mixing, small quantities should always be tested using a simple jar test. Add the required amount of **Drive 75 DF** to half-filled tank while agitating, then add the fertilizer product. Complete filling spray tank to desired level.

V. Spraying Instructions for Drive 75 DF herbicide

Apply with properly calibrated ground equipment in sufficient water per acre to provide uniform spray distribution (at least 20 gallons of water per acre or at least 0.5 gallons per 1000 sq ft). Use low pressure sprayers at 20-40 psi.

Maintain continuous agitation during spraying with good mechanical or bypass agitation. Nozzle screens must be no finer than 50 mesh (100 mesh is finer than 50 mesh). Check sprayer routinely to determine proper calibration. Flat fan, flood or cone nozzles may be used. Nozzles should be arranged to obtain uniform coverage for turf and weeds to be controlled. Boom height, nozzle selection, and pressure should be adjusted to provide uniform coverage and minimize spray drift.

Avoid overlaps that will increase rates above those labeled for use.

Avoid application when winds may cause drift.

Procedure For Cleaning Spray Equipment:

Clean application equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturers directions before and after applying this product.

VI. Compatibility Test for Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of labeled rate per acre.

1) **Water:**

For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.

2) **Products in PVA bags:**

Cut an opening in the water-soluble PVA bag just large enough to use a teaspoon for measuring purposes. Use the opened water-soluble PVA bag first when preparing spray solution. Boron-containing fertilizers can be incompatible with PVA material. Include PVA material if a boron fertilizer is intended to be used. Cap the jar and invert 10 cycles.

3) **Water-dispersible products (such as Drive® 75 DF herbicide):**

-dry flowables, wettable powders, suspension concentrates, or suspo-emulsions
-Cap the jar and invert 10 cycles.

4) **Water-soluble products (such as Basagran® T/O herbicide):**

-Cap the jar and invert 10 cycles.

5) **Emulsifiable concentrates:**

-(methylated seed oil)
-Cap the jar and invert 10 cycles.

6) **Water-soluble additives:**

-Cap the jar and invert 10 cycles.

7) Let the solution stand for 15 minutes.

8) **Evaluate** the solution for uniformity and stability.

The spray solution should not have free oil on the surface nor thick (clabbered) texture. For WG or WP products, a fine precipitate that is easily resuspended is normal; large non-dispersible particles (>300 microns) that precipitate on standing are a sign of tank mix incompatibility. **DO NOT** use any spray solution that could clog spray nozzles.

VII. General Restrictions and Limitations

- **DO NOT** apply more than 2 pounds of **Drive 75 DF** per acre (or 0.73 oz per 1000 sq ft) in one year.
- **DO NOT** apply to golf course collars or greens.
- **DO NOT** make applications of **Drive 75 DF** to drought-stressed turfgrass.
- **DO NOT** apply to fine fescue unless it is part of a seed blend.
- **DO NOT** apply to Bahiagrass, carpetgrass, St. Augustinegrass, centipedegrass, dichondra, or lawns or turf where desirable clovers are present.
- **DO NOT** apply within 4 weeks after seeding emergence of Kentucky bluegrass, creeping bentgrass, fine fescue blends and perennial ryegrass.

- **DO NOT** apply to exposed feeder roots of trees or ornamentals. Be particularly careful within the drip line of trees and other ornamental species.
- **DO NOT** apply into any ornamental bed.
- **DO NOT use clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.**
- **DO NOT** plant eggplants or tobacco within 12 months on fields treated with **Drive 75 DF**.
- **DO NOT** plant tomatoes or carrots within 24 months on fields treated with **Drive 75 DF**.
- Apply when wind speed is less than 10 mph as drift may cause damage or death of nontarget area vegetation. **DO NOT** apply when conditions favor drift from target area.
- Use a lawn-type sprayer with coarse spray as wind drift is less likely.
- Avoid mist and spray onto vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants, **especially plants belonging to the Solanaceae family, such as tomatoes, eggplants, and bell peppers.**
- **DO NOT** pour spray solutions near these plants.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** use to formulate or reformulate any other pesticide product which is not registered by EPA.
- For best results, weeds should not be under stress from lack of water, excessive water, low fertility, mowing shock, excessive hot or cold temperatures, or injury from other herbicide applications.

VIII. Turfgrass Tank Mixes

Read and follow the applicable **Restrictions and Limitations** and **Directions For Use** on all products involved in tank mixing. The most restrictive labeling applies to any tank mix. To increase spectrum of control of broadleaf weed species, a tank mix with **2,4-D**, triclopyr, **MCPA**, **MCP**, **Three Way™**, **Trimec®**, or other broadleaf herbicide can be used. For extended residual control, apply **Drive 75 DF** with **Pendulum®** or **PRE-M® herbicides**. For sedge control, applications of **Drive 75 DF** with **Basagran® SG**, **Basagran T/O**, **Lescogran® herbicides**, or **Image® 70 DG herbicide**, or MSMA can be made.

Combinations with MSMA will aid in control of certain grassy weeds, such as Bahiagrass or kikuyugrass. Consult tolerant turfgrass species on all labels. Separate applications should be made if all target weeds are not at the correct growth stage for treatment at the same time.

Physical incompatibility, reduced weed control, or turf injury may result from mixing **Drive 75 DF** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers.

Before tank mixing, a simple jar test is required to insure compatibility of herbicides or other pesticide and/or additives. Refer to manufacturers labels for specific use directions, precautions, and limitations before tank mixing with **Drive 75 DF**, and follow those that are most restrictive.

9/12

Table 1. Turf Tolerance (Established)		
Highly Tolerant	Moderately Tolerant	Susceptible
Bermudagrass, Common ¹ Bluegrass, Annual Bluegrass, Kentucky Buffalograss Fescue, Tall Ryegrass, Annual Ryegrass, Perennial Zoysiagrass	Bentgrass, Creeping ¹ Bermudagrass, Hybrid ¹ Bluegrass, Rough (<i>Poa trivialis</i>) Fescue, Chewing's Fescue, Fine ² Fescue, Hard Fescue, Red Paspalum, Seashore	Bahiagrass Bentgrass, Colonial Bentgrass, Seaside Centipedegrass Dichondra St. Augustinegrass
<p>¹ Yellowing that occurs on these species can be reduced by the addition of chelated iron or sprayable soluble nitrogen fertilizer (see II. Application Information, III. Addition of Adjuvants).</p> <p>² Apply Drive® 75 DF herbicide to fine fescue only when it is part of a blend.</p> <p>DO NOT use on golf course greens and collars.</p> <p>Consult Table 4. Seeding/Overseeding/Sprigging Timing Chart for this situation.</p>		

Table 2. Drive 75 DF Application to Established Creeping Bentgrass		
Turfgrass Species	Application Rate/Timing	Additive Rate
Bentgrass, Creeping ¹	Drive 75 DF must be applied in 2 - 3 split applications at 0.123 - 0.245 oz per 1000 sq ft (not to exceed 2 lbs of product per acre per season). Sequential application(s) should be timed 21 days apart.	Use methylated seed oil at 0.55 oz per 1000 sq ft (1.5 pints per acre).
<p>¹ Yellowing that occurs on these species can be reduced by the addition of chelated iron or sprayable soluble nitrogen fertilizer (see II. Application Information, III. Addition of Adjuvants).</p> <p>DO NOT use on golf course greens and collars.</p> <p>Consult Table 4. Seeding/Overseeding/Sprigging Timing Chart for this situation.</p>		

Table 3. Application Rates and Timing for Postemergence Weed Control in Turf			
Weed Species		Drive® 75 DF herbicide Rate	Additive Rate
Grasses Controlled		Broadcast Applications: 1 pound of product per acre or 0.367 oz per 1000 sq ft	Apply: 1.5 pints per acre (0.55 oz per 1000 sq ft) methylated seed oil
Common Name	Scientific Name		
Barnyardgrass	<i>Echinochloa crusgalli</i>	Spot Applications: Apply 0.367 oz of Drive 75 DF per 1000 sq ft of treated area. Refer to footnotes in Tables 2 and 3 for specific turfgrass or weed instructions.	
Crabgrass, Large ^{1,4}	<i>Digitaria sanguinalis</i>		
Crabgrass, Smooth ^{1,4}	<i>Digitaria ischaemum</i>		
Foxtail, Giant ¹	<i>Setari faberi</i>		
Foxtail, Green ¹	<i>Setari viridis</i>		
Foxtail, Yellow ¹	<i>Setari glauca</i>		
Kikuyugrass ^{2,3}	<i>Pennisetum clandestinum</i>		
Signalgrass, Broadleaf ¹	<i>Brachiaria platyphylla</i>		
Torpedograss ³	<i>Panicum repens</i>		
Broadleaf Weeds Controlled			
Common Name	Scientific Name		
Bindweed, Field	<i>Convolvulus arvensis</i>		
Clover, Hop	<i>Trifolium aureum</i> Pollich		
Clover, Red	<i>Trifolium pratense</i>		
Clover, White	<i>Trifolium repens</i>		
Daisy, English ^{2,5}	<i>Bellis perenne</i>		
Dandelion, Common ²	<i>Taraxacum officinale</i>		
Dollarweed	<i>Hydrocotyle umbellata</i>		
Geranium, Carolina	<i>Geranium carolinium</i>		
Medic, Black	<i>Medicago lupulina</i>		
Morningglory spp.	<i>Ipomea</i> sp.		
Speedwell, Common	<i>Veronica officinalis</i>		
Speedwell, Slender	<i>Veronica filiformis</i>		
Speedwell, Thymeleaf	<i>Veronica serpyllifolia</i>		
Violet, Wild	<i>Viola</i> sp.		

¹ Under certain conditions, application of **Drive 75 DF** made to annual grasses at 2 - 4 tiller may not provide complete control. A sequential application will be required for grass control in these situations. Optimum control is achieved when applications of **Drive 75 DF** + methylated seed oil are applied either before second tiller or as weed grasses mature.

² Tank mix partner or sequential application required.

³ Make 3 sequential applications of 0.245 oz of **Drive 75 DF** per 1000 sq ft at 14 - 21 day intervals.

⁴ Biotypes of large and smooth crabgrass in California have shown varied response to **Drive 75 DF**. If control failure occurs following a full or split application, **DO NOT** reapply **Drive 75 DF**. Change to a herbicide with a different mode of action.

⁵ **NOT FOR USE** to control this weed in California.

11/12

Table 4. Seeding/Overseeding/Sprigging Timing Chart¹

Variety	Before seeding ²	At seeding	7 days after emergence	14 days after emergence	28 days after emergence
Annual Bluegrass	OK	OK	OK	OK	OK
Annual Ryegrass	OK	OK	OK	OK	OK
Buffalograss	OK	OK	OK	OK	OK
Common Bermudagrass ³ (For sprigging see footnote 3)	OK	OK	OK	OK	OK
Creeping Bentgrass	OK	NO	NO	NO	OK
Fine Fescue (in blend)	OK	NO	NO	NO	OK
Hybrid Bermudagrass ³ (For sprigging see footnote 3)	OK	OK	OK	OK	OK
Kentucky Bluegrass	OK	NO	NO	NO	OK
Perennial Ryegrass	OK	OK	NO	NO	OK
Seashore Paspalum ^{3,4} (For sprigging see footnote 3)	NO	NO	NO	OK	OK
Tall Fescue	OK	OK	OK	OK	OK
Zoysiagrass ³ (For sprigging see footnote 3)	OK	OK	OK	OK	OK

¹Note: No adjuvant or additive should be used when **Drive® 75 DF herbicide** applications are made on newly emerged turf seedlings until 28 days after emergence; with the exception of seashore paspalum, a **Drive 75 DF** application rate of 0.367 oz/1000 sq ft can be made to all other turfgrass species in above Table 4.

²**Drive 75 DF** can be applied 7 days or greater prior to seeding.

³**Drive 75 DF** can be used anytime prior to, at or after sprigging as indicated by turfgrass species in above **Table 4**.

⁴0.18 oz to 0.367 oz/1000 sq ft application can be made at times indicated in above **Table 4**.

Application of **Drive 75 DF** should be timed around the seeding operations using the above chart as a reference point.

Table 5. Spot Spraying with Drive 75 DF

• Spray Mix Volume (gallons)	Drive Product in Mix (tablespoons)	MSO Adjuvant in Mix (tablespoons)
1	1	2
2	2	4
3	3	6

• Apply at the rate of 1 gallon per 1000 sq ft.
1 tablespoon = 0.367 oz of **Drive 75 DF** product.

NOTE: For consistent results make **Drive 75 DF** application to newly germinated 1 - 2 leaf crabgrass, to 1-tiller crabgrass and when crabgrass has matured to 5 tillers or greater.*

* Under certain conditions, applications of **Drive 75 DF** made to annual grasses 2 - 4 tiller may not provide complete control. A sequential application will be required for grass control in these situations.

12/12

Conditions of Sale and Warranty

The **Directions For Use** of this product reflects the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Basagran, Drive, Image and Pendulum are registered trademarks of BASF.

DyneAmic is a registered trademark of Helena Chemical Company.

LI 700 is a registered trademark of UHS (United Horticultural Supply).

PRE-M and Lescogran, Lescro Spreader Sticker and Three Way are registered trademarks of Lesco, LLC.

Surf King and Thoroughbred are registered trademarks of Estes.

Target Pro is a registered trademark of Target.

Trimec is a registered trademark of PBI-Gordon Corporation.

© 2006 BASF Corporation
All rights reserved.

007969-00130.20060802b.NVA 2006-04-073-0199
Supersedes: NVA 2004-04-073-0248

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

