| | 7969-1 | 46 3 | 124/98 | · · · · · · · · | 1 Page 1710 |
|---|---|--|---|-------------------------------|---|
| | United States Environmental Protection Washington, DC 204 | | Form Approve | Registratio Amendme | e e state |
| | Applicatio | on for Pesti | icide - Section | 1 ···· | the course which the mail series and |
| L. Company/Product Number 4. Company/Product (Name) | 7969-146 | 1 | | 1 | 3. Proposed Classification |
| Guardsman h 5. Name and Address of App BASF Corporati P.O. Box 13528 Research Trian | on | (b)(i) to: | | | with FIFRA Section 3(c)(3) in composition and labeling |
| Check if this | is a new address | Pro | duct Name | | |
| | | Section | - 11 | | |
| Amendment - Explain Resubmission in respo X Notification - Explain | onse to Agency letter dated | [| Final printed lab Agency letter da "Me Too" Applie Other - Explain b | ation. | MAR 2 4 1998 |
| EPA regulation | tion is consistent s at 40 CFR 152.46 | , and no | other chan f formula o | ges have l | |
| 1. Material This Product Will | 1 | | | | |
| Child-Resistant Packaging Yes X No <i>certification must</i> | Unit Packaging Yes X No If "Yes" No. per Unit Packaging wgt. container | Water Schubl Yes X No If "Yes" Package wgt | No. per | | Itainer letel astic lass aper ther (Specify) |
| 3. Location of Net Contents I | | tail Container , 30 bulk | | ocation.of Label I On labe | |
| 6. Manner in Which Label is . | Affixed to Product | graph glued illed | Other | | |
| <u></u> | | Section | | | |
| ······································ | items directly below for identificatio | (' | o be contacted, if ne | <u> </u> | |
| Jonathan B | ryant | Title Manager, | Regulatory | y Affairs | ephone No. (Include Area Code) 919-547-2978 |
| | Certificat ments I have made on this form and y knowlingity false or misteading sta law. | i all attachments | | | |
| 2. Signature | · · · · | 3. Title Manager, | Regulatory | y Affairs | |
| 4. Typed Name | | 5. Date | | | |
| Jonathan Br | yant | Mar | ch 18, 1998 | 3 | |
| PA Form 8570-1 (Rev. 3-94 | Previous editions are obsolete. | <u></u> | White - | EPA File Copy (or | iginal) Yellow - Applicant Copy |

RESTRICTED USE PESTICIDE (GROUND AND SURFACE WATER CONCERNS)

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

THIS PRODUCT IS A RESTRICTED USE HERBICIDE DUE TO GROUND AND SURFACE WATER CONCERNS.



RT 12-3-97 3-19-98

NOTIFICATION

MAR 2 4 1998

FOR WEED CONTROL IN FIELD CORN, SEED CORN, SWEET CORN, POPCORN, AND GRAIN SORGHUM



ACTIVE INGREDIENTS:

ĺ

-_

| Dimethenamid: 2-chloro-N- | |
|---|---------|
| [(1-methyl-2-methoxy)ethyl]-N- | |
| (2,4-dimethyl-thien-3-yl)-acetamide | 24.8P÷ |
| Atrazine: 2-chloro-4-ethylamino-6-isopropyl- | |
| amino-s-triazine. | 2874 |
| INERT INGREDIENTS: | 46.8% |
| Total | 100.0%+ |
| Cuardeman [®] contains 2.32 nounds of dimethossend and 1 | |

Guardsman[®] contains 2.33 pounds of dimethenamid and 2.47 pounds of atrazine per gallon.

* Contains petroleum distillates

EPA Reg No. 7969-146

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a ciguien para que se la explique a usted en detalle. (It you do not understand the label, find somecne to explain it to you in detail.) See next page for additional Precautionary Statements and Statement of Practical Treatment.

SHAKE BEFORE USING

Users must read and follow all precautionary statements and instructions for use in order to minimize potential for atrazine to reach ground and surface water.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING AVISO

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapor or spray. Avoid contact with skin. Prolonged or frequent repeated skin contact may cause allergic reactions in some individuals.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Hold eyelids open and flush with steady, gentle stream of water for 15 minutes. Get medical attention.

If swallowed: Call a doctor or get medical attention. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Do not give anything by mouth to an unconscious person. Avoid alcohol. If inhaled: Remove victim to fresh air. If not

If inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth to mouth. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage! May bose an aspiration pneumonia hazard....

.....

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category section chart.

Applicators and other handlers (other than mixers and loaders) must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton
- Chemical-resistant footwear plus socks
- Protective eyewear
- Mixers and Loaders must wear:
- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton
- Chemical resistant footwear plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) [4-6]], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Atrazine is toxic to aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters or rinsate. Do not apply when weather conditions (gusty winds, high temperatures, low humidity, and when a temperature inversion exists) favor drift from treated areas.

Dimethenamid has properties that may result in groundwater contamination. Application in areas where soils are permeable or coarse and groundwater is near the surface could result in groundwater contamination. Following application and during rainfall events that cause run-off, this chemical may reach surface water bodies including streams, rivers, and reservoirs. **Guardsman[®]** herbicide (**Guardsman**) contains the active ingredient atrazine. Atrazine can leach through soil and has been found to result in contamination of water supplies by way of groundwater. Therefore, growers are advised to avoid use of **Guardsman** in well-drained loamy-sand to sand soils, particularly in areas having high groundwater tables.

39

Groundwater contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material. This product may not be mixed, loaded, or used within 50 feet of all wells including abandoned wells, drainage wells, and sinkholes. This product may not be mixed or loaded within 50 feet of intermittent streams or rivers, natural or impounded lakes and reservoirs. This product may not be applied aerially or by ground within 66 feet of the points where field surface water run-off enters perennial or intermittent streams and rivers or within 200 feet around natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 foot buffer or set-back from run-off points must be planted to crop or seeded with grass or other suitable crop.

Care must be taken when using this product to prevent 1) back siphoning into wells, 2) spills, or 3) improper disposal of excess pesticide, spray mixtures or rinsates.

Check valves or antisiphoning devices must be used on all mixing equipment.

TILE-TERRACED FIELDS CONTAINING STANDPIPES

To ensure protection of surface water from run-off through standpipes and tile outlets in terraced fields, one of the following options may be used:

- Do not apply this product within 66 feet of standpipes in tile-outletted terraced fields.
- Apply this product to the entire tile-outletted field and immediately incorporate it to a depth of 2-3 inches in the entire tile-outletted terraced field.
- Apply this product to the entire tile-outletted field under a no-til practice only when high crop residue management practices are used. High crop residue management practice is described as a crop management practice where little or no crop residue is removed from the field during or after crop harvest.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE ****

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a vay that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.



AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the product is soil injected or soil

incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

Coveralls

- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton
- Chemical resistant footwear plus socks
- Protective eyewear

Where there are state/local requirements regarding atrazine use (including lower maximum rates and/or higher set-backs) which are different from this label, the more restrictive/protective requirements apply.

Do not apply this product through any type of irrigation system. **IMPORTANT:** Read the Limited Warranty and Liability and all applicable Directions for Use before using this product.

GENERAL INFORMATION

Guardsman[®] herbicide is a selective preemergence herbicide for control of most annual grasses, certain annual broadleaf weeds and sedges in field corn, seed corn, sweet corn, or popcorn and grain sorghum. This product will provide most effective weed control when applied (by ground or aerial equipment), and subsequently incorporated into soil by rainfall, sprinkler irrigation, or mechanical tillage prior to weed seedling emergence from soil.

Guardsman is recommended for preplant incorporated, preplant surface, preemergence or early postemergence treatment. **Guardsman** application may be made using either water or fluid fertilizer as the spray carrier or it may be impregnated on and applied with dry bulk fertilizer. Sprayable fluid fertilizer as a carrier is not recommended for use after crop emergence. Observe all precautionary statements and limitations on labeling of all products used in tank mixtures. Tank mix recommendations are for use only in states and on application sites where the tank mix product is registered. Do not apply this product through any type of irrigation system.

4910

Single or split applications of **Guardsman** may not exceed a total of 5 pints per acre in one crop year.

Recommended use rates of this product vary by soil type. The most accurate indicator of appropriate use rate for **Guardsman** is the Cation Exchange Capacity (CEC) of the soil to be treated. CEC values are available in standard soil testing procedures. If CEC values are not available, the recommended use rate of **Guardsman** may be determined using the soil texture and organic matter. Soil texture groupings used in this label are coarse, medium and fine. Soil textures included in these groupings are as follows:

| Coarse | Medium | Fine |
|--------------------------|-----------------|--------------------------|
| Sand | Silt | Sandy Clay |
| Loamy Sand | Silt Loam | Sandy Clay Silty Clay |
| Loamy Sand Sandy Loam | Loam | Silty Clay Loam |
| • | Sandy Clay Loam | Ćlay Loam |
| | | Ćlav |

Guardsman use rates, by either soil CEC values or by soil texture and organic matter, are given in specific crop use sections of this label. When use rates are expressed in ranges, use the lower end of the rate range for lower CEC values and use the higher end of the rate range for higher CEC values. If texture and organic matter are used to determine use rates, use the lower end of the rate range for more coarsely textured soils low in organic matter and the higher end of the rate range for more finely textured soils that are high in organic matter.

WEED SPECIES CONTROLLED BY GUARDSMAN

Guardsman applied at recommended rates and application timings will control many annual grass and broadleaf weeds.*

ANNUAL GRASSES

Barnyardgrass Broadleaf, Signalgrass Crabgrass, Smooth, Large Fall Panicum Foxtail, Giant, Green, Yellow

SEDGE Rice Flatsedge

ANNUAL BROADLEAVES

Carpetweed Cocklebur** Common Purslane Florida Pusley Jimsonweed Kochia Lambsquarters Momingglory, Annual Mustards Nightshade, Black, Eastern Black, Hairy Gocsegrass Johnsongrass (seedling)** Red Rice Southwestern Cupgrass Wild Oats Witchgrass

Yellow Nutsedge**

Palmer Amaranth Pigweed, Prostrate, Redroot, Smooth, Tumble Ragweed, Common, Giant** Smortweed Spurge, Nodding, Spotted Velvetleaf** Warthump Wild Burkwheat

- Some weed species may have tractine resistant/biotypes that will not be controlled adequately by this product. If resistant biotypes are suspected use an ctarnate program or use nontriazine products such as BANVEL® or CLARITY® Herbicide in combination or sequentially with this product ,
- ** For best control of these species use the highest recommended rate by soil type. If dry conditions exist near application or excessive rainfall occurs early in season, a postemergence herbicide and/or cultivation may be required to aid in control of these weeds.

3

ADDITIONAL WEED SPECIES PARTIALLY CONTROLLED

Guardsman will provide partial control or suppression of eclipta, woolly cupgrass, wild proso millet, shattercane, sandbur, and Texas panicum. Control of these species will generally not be commercially acceptable. To complement control **Guardsman** should be used in tank mixtures or sequential applications with other herbicides that provide additional control of these weed species.

GUARDSMAN USE METHODS AND TIMINGS

Guardsman may be applied preplant surface, preplant incorporated, or preemergence to corn or grain sorghum; or may be applied early postemergence to corn. Best weed control will result when applied (by ground or aerial equipment), and subsequently incorporated into soil by rainfall, sprinkler irrigation, or mechanical tilloge prior to weed seedling emergence from soil.

PREPLANT SURFACE APPLICATIONS

For use in minimum tillage or no-tilage production systems, application of Guardsman alone or in tank mixtures may be made up to 45 days before planting. When making early preplant applications (15 to 45 days prior to planting) use the highest rate recommended for the specific soil type. Early preplant applications are not recommended for use on coarse textured soils or in areas where average annual rainfall (or rainfall + irrigation) typically exceeds 40 inches. Early preplant applications may be applied as part of a split application program where the second application is made after planting (use $^{2}/_{3}$ of Guardsman rate early followed by 1/3 of rate after planting). A split application is recommended when initial application is made more than 30 days prior to planting. Tank mixtures with postemergence herbicides such as Roundup® or Gramoxone® Extra must be used when weeds are present at the time of application.

PREPLANT INCORPORATED APPLICATIONS

Apply **Guardsman** and incorporate into the upper (1 to 2 inch) soil surface up to two weeks before planting. Use a harrow, rolling cubivator, finishing disk, or other implement capable of giving uniform shallow incorporation. Avoid deeper incorporation or reduced weed control may result.

PREEMERGENCE SURFACE APPLICATIONS

Broadcast treatment uniformly to the soil surface after planting and before crop emergence. Rainfall, sprinkler irrigation, or shallow mechanical incorporation after application is required to move this product into the upper soil surface where weed seeds germinate. If adequate rainfall or irrigation does not occur and weed seedling emergence begins, a stallow cultivation or rotary hoeing will improve performance.

EARLY POSTEMERGENCE APPLICATIONS

Guardsman may be used in early postemergence applications up to 8 inch tall corn. Application must be made before weeds are greater than 1.5 inches tall or in a tank mixture with products that control the emerged weeds. Note restrictions and use directions on tank mix product labels. Do not apply to grain sorghum after crop emergence.

GUARDSMAN FIELD CORN, SEED CORN, SWEET CORN, OR POPCORN USE PROGRAMS

5710

Guardsman may be used as a preplant surface applied, preplant incorporated, preemergence, or early postemergence (up to 8 inches tall) treatment to corn. Corn, as used in this label, refers to field corn (grown for grain, silage or seed), sweet corn (does not include sweet corn grown for seed), and popcorn. Before application to seed corn, sweet corn, or popcorn, first verify with your local seed company (supplier) the **Guardsman** selectivity on your inbred line or variety. This precaution will help avoid potential injury to sensitive varieties.

Com may be grazed or fed to livestock at 40 or more days after application of **Guardsman**. Sweet corn ears may be harvested at 50 or more days after application of **Guardsman**.

GUARDSMAN USE RATES IN FIELD CORN, SEED CORN, SWEET CORN, AND POPCORN

Recommended broadcast use rates for Guardsman when used alone or in tank mix combinations and/or sequential applications with other herbicides in corn are given in the following tables^{*}. Cation exchange capacity (CEC) of soil is the preferred method for determining use rate. If CEC is not known, select use rate based on soil texture and organic matter content. If Guardsman is used in early preplant applications, and/or used on soils with heavy surface plant residue add 0.5 pints per acre to the rates given in the following tables. Do not exceed 4.75 pints per acre on highly erodible soils with less than 30% plant residue cover prior to crop emergence. Do not exceed 5 pints per acre per crop year on any soil.

USE RATE DETERMINED BY CATION EXCHANGE CAPACITY (CEC) OF SOIL

| Cation Exchange Capacity (CEC) of Soil | < 5.0 | 5.0-9.0 | 10-14 | 15-20 | > 20.0** |
|---|-------|---------|---------|---------|----------|
| Guardsman Use Rate (Pints/acre) | 2.5 | 3.0-3.5 | 3.5-4.0 | 4.0-4,5 | 5.0 |

USE RATE DETERMINED BY SOIL TEXTURE AND ORGANIC MATTER CONTENT

| | Guardsman Use Rate (Pints per Acre) | | | |
|--------------|-------------------------------------|--------------|--|--|
| • | Organic Matter | | | |
| Soil Texture | Less than 3% | 3% or more** | | |
| Caarse | 2.5 to 3.0 | 3.0 - 3.5 | | |
| Medium | 3.0 to 4.0 | 4.0 - 4.5 | | |
| Fine | 4.0 to 4.5 | 4.5.5.0 | | |

* See tank mix descriptions below for the recommended use rate ranges of other herbicides used in combination with **Guardsman**. Refer to tank mix section of companion product labels for actual use rate by soil type, weed species, and weed or crop prowth stage.

type, weed species, and weed or crop growth stage. * On all soils with 8% to 20% organic motter, use 5 pints per acre of **Guardsman**. **Guardsman** is not recommended for use on soils with more than 20% organic matter.



GUARDSMAN TANK MIXTURES AND SEQUENTIAL USE PROGRAMS IN CORN

Guardsman may be applied prior to, in tank mix with, or following the use of one or more of the following herbicides: Accent[®], atrazine, BANVEL[®], Basagran[®], Beacon[®], Bladex[®], CLARITY[®], FRONTIER[®], Gramoxoner[®] Extra, Laddok[®] S-12, MARKSMAN[®], Princep[®], Prowl[®], Pursuit[®], Roundup[®], or 2,4-D. Guardsman use rates for herbicide combination programs and specific tank mixtures are presented in the above label section, <u>Guardsman Use</u> Rates in Field Corn. Seed Corn. Sweet Corn. and Popcorn.

Guardsman[•] herbicide contains 2.67 pounds active of atrazine per gallon (0.334 lb ai per pint). When tank mixing or making sequential applications with atrazine or **Marksman** do not exceed the following total combined amounts of atrazine:

Prior to corn emergence

1.6 lb ai/A - Highly erodible soils with less than 30% plant residue cover
2.0 lb ai/A - Other soils

After corn emergence

(

2.0 lb ai/A - Any soil

Prior to and after corn emergence (sequential applications)

2.5 lb ai/A - Any soil

When using tank mix or sequential applications with **Guardsman**, always read the companion.product label(s) to determine specific use rates by soil types, weed species, and weed or crop growth stage. Tank mixing must be in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing prohibition against such mixing.

Guardsman may be tank mixed with one or more of the following herbicide combination programs.

| Guardsman plus BANVEL (0.5 to 1.0 pt/A) or CLARITY (8.0 to 16.0 fl oz/A) or MARKSMAN (2.0 to 3.5 pt/A) | Application may be made preplant surface or preemergence in minimum or no tillage corn in all soil types. Preemergence applications may be used in all tillage systems where corn is grown on soils with CEC greater than 15, or medium and fine textured soils with 2.5% or more organic matter (For best per- formance make application just prior to corn emergence). Early postemergence (up to 8 inch tall corn) applications may be made on all soil types. This treatment must be combined with a herbicide that provides post- emergence control of grass weeds if they are greater than 1.5 inches tall at the time of application. MARKSMAN con- tains atrazine. Do not exceed the maximum amount of atrazine allowed for the specific condition. This tank mixture is not labeled for use on sweet corn |
|---|---|

| Guardsman plus FRONTIER {4.0 to 12.0 fl oz/A} | Application may be made pre- plant surface, preplant incorpor- ated, preemergence, or early postemergence (up to 8 inch tall corn). FRONTIER may be added for improved annual grass control and/or for reducing atrazine use rate. Do not exceed 1.5 lb ai/A of dimethenamid (dimethenamid active is 0.29 lb ai/pint in Guardsman and 0.06 lb ai/fl oz in FRONTIER). |
|---|--|
| Guardsman plus ATRAZINE {0.5 to 1.0 lb ai/A} | Application may be made pre- plant surface, preplant incorpor- ated, preemergence, or early postemergence (up to 8 inch tall corn). This treatment must be combined with a herbicide that provides postemergence control of weeds if they are greater than 1.5 inches tall at the time of application. Crop oil concentrate may be added to this mixture to aid in control of emerged weeds. Do not add crop oil concentrate if this mixture is combined with other herbicides for early post- emergence applications. Do not exceed the amount of atrazine allowed for the specific condition. |
| Guardsman plus BLADEX® {1.25 to 3.0 lb ai/A} | Application may be made pre- plant surface, preplant incorpor- ated, or preemergence to corn. This mixture is not recommended for use on soils with CEC less than 5 or on sand or loamy sand soils with organic matter less than 1%. |
| Guardsman plus PROWL® 3.3 EC [1.8 to 3.6 pt/A] | Application may be made pre- emergence to early postemer- gence (up to 3 inch tall corn) but before weeds are greater than 1.5 inches tall. |
| Guardanaa alaa | |
| Guardsman plus PRINCEP [®] {1.0 lb ai/A} | Application may be made preplant surface, preplant incor- porated or preemergence to corn. |
| PRINCEP® | preplant surface, preplant incor- |
| PRINCEP [®] (1.0 lb ai/A) Guardsman plus PURSUIT® 2.0 SL (4 fl oz/A) or | preplant surface, preplant incor- porated or preemergence to corn. Use only on Pursuit resistant (IR) or tolerant (IT) corn varieties. Appli- cation may be made preplant surface, preplant incorporated, preemergence, or early postemer- |
| PRINCEP [®] {1.0 lb ai/A} Guardsman plus PURSUIT [®] 2.0 SL [4 fl oz/A] or PURSUIT [®] 70DG (1.4 fl oz/A) Guardsman plus GRAMOXONE EXTRA [®] {1.5 to 3.0 pt/A] or ROUNDUP [GLYPHOSATE] - | preplant surface, preplant incor- porated or preemergence to corn. <u>Use only on Pursuit resistant (IR) or</u> tolerant (III) com varieties. Appli- cation may be made preplant surface, preplant incorporated, preemergence, or early postemer- gence (up to 3 inch tall weeds). Application may be made preplant surface or preemergence to control emerged weeds in minimum or no tillage corn. Applications must be made prior to corn emergence. To improve control of emerged broadleaf weeds, herbicides such as Banvel and/or 2,4-D should be added to |
| PRINCEP [®] {1.0 lb ai/A} Guardsman plus PURSUIT [®] 2.0 SL [4 fl oz/A] or PURSUIT [®] 70DG (1.4 fl oz/A) Guardsman plus GRAMOXONE EXTRA [®] {1.5 to 3.0 pt/A] or ROUNDUP [GLYPHOSATE] - | preplant surface, preplant incor- porated or preemergence to corn. <u>Use only on Pursuit resistant (IR) or</u> tolerant (III) com varieties. Appli- cation may be made preplant surface, preplant incorporated, preemergence, or early postemer- gence (up to 3 inch tall weeds). Application may be made preplant surface or preemergence to control emerged weeds in minimum or no tillage corn. Applications must be made prior to corn emergence. To improve control of emerged broadleaf weeds, herbicides such as Banvel and/or 2,4-D should be added to |
| PRINCEP [®] {1.0 lb ai/A} Guardsman plus PURSUIT [®] 2.0 SL [4 fl oz/A] or PURSUIT [®] 70DG (1.4 fl oz/A) Guardsman plus GRAMOXONE EXTRA [®] {1.5 to 3.0 pt/A] or ROUNDUP [GLYPHOSATE] - | preplant surface, preplant incor- porated or preemergence to corn. <u>Use only on Pursuit resistant (IR) or</u> tolerant (III) com varieties. Appli- cation may be made preplant surface, preplant incorporated, preemergence, or early postemer- gence (up to 3 inch tall weeds). Application may be made preplant surface or preemergence to control emerged weeds in minimum or no tillage corn. Applications must be made prior to corn emergence. To improve control of emerged broadleaf weeds, herbicides such as Banvel and/or 2,4-D should be added to |
| PRINCEP [®] {1.0 lb ai/A} Guardsman plus PURSUIT [®] 2.0 SL [4 fl oz/A] or PURSUIT [®] 70DG (1.4 fl oz/A) Guardsman plus GRAMOXONE EXTRA [®] {1.5 to 3.0 pt/A] or ROUNDUP [GLYPHOSATE] - | preplant surface, preplant incor- porated or preemergence to corn. <u>Use only on Pursuit resistant (IR) or</u> tolerant (III) com varieties. Appli- cation may be made preplant surface, preplant incorporated, preemergence, or early postemer- gence (up to 3 inch tall weeds). Application may be made preplant surface or preemergence to control emerged weeds in minimum or no tillage corn. Applications must be made prior to corn emergence. To improve control of emerged broadleaf weeds, herbicides such as Banvel and/or 2,4-D should be added to |
| PRINCEP [®] {1.0 lb ai/A} Guardsman plus PURSUIT [®] 2.0 SL [4 fl oz/A] or PURSUIT [®] 70DG (1.4 fl oz/A) Guardsman plus GRAMOXONE EXTRA [®] {1.5 to 3.0 pt/A] or ROUNDUP [GLYPHOSATE] - | preplant surface, preplant incor- porated or preemergence to corn. <u>Use only on Pursuit resistant (IR) or</u> tolerant (III) com varieties. Appli- cation may be made preplant surface, preplant incorporated, preemergence, or early postemer- gence (up to 3 inch tall weeds). Application may be made preplant surface or preemergence to control emerged weeds in minimum or no tillage corn. Applications must be made prior to corn emergence. To improve control of emerged broadleaf weeds, herbicides such as Banvel and/or 2,4-D should be added to |
| PRINCEP [®] {1.0 lb ai/A} Guardsman plus PURSUIT [®] 2.0 SL [4 fl oz/A] or PURSUIT [®] 70DG (1.4 fl oz/A) Guardsman plus GRAMOXONE EXTRA [®] {1.5 to 3.0 pt/A] or ROUNDUP [GLYPHOSATE] - | preplant surface, preplant incor- porated or preemergence to corn. <u>Use only on Pursuit resistant (IR) or</u> tolerant (III) com varieties. Appli- cation may be made preplant surface, preplant incorporated, preemergence, or early postemer- gence (up to 3 inch tall weeds). Application may be made preplant surface or preemergence to control emerged weeds in minimum or no tillage corn. Applications must be made prior to corn emergence. To improve control of emerged broadleaf weeds, herbicides such as Banvel and/or 2,4-D should be added to |

5

Tank mixture for postemergence use in corn.

Guardsman plus ACCENT 75 WDG {1/3 to 2/3 oz/A}

Application may be made from emergence up to 8 inch tall corn when weeds are emerged and actively growing. This tank mixture is not labeled for use on sweet corn. Use lower rates $(^{1}/_{3} \text{ oz})$ of ACCENT for control of foxtail, barnyardgrass, fall panicum, or seedling Johnsongrass when all are less than 3 inches tall. **Guardsman** at reduced use rates: 2 to 2.5 pints/acre - on soils with CEC less than 10 (coarse soils or medium and fine textured soils with less than 3% organic matter), or 2.5 to 3.5 pints/acre - on soils with CEC of 10 or more (medium or fine textured soils with 3% or more organic matter) may be used in this mixture. Addition of BANVEL, CLARITY, or BAS 653 00H Herbicide to this mixture will provide burndown and residual control of broadleaf weeds. Add non-ionic surfactant at .25% v/v plus 4% v/v liquid fertilizer (28, 30, or 32% UAN). Other herbicides are not recommended for use with this mixture.

GUARDSMAN GRAIN SORGHUM USE PROGRAMS

Guardsman® herbicide may be used as a preplant surface, preplant incorporated, or preemergence application in sorghum. A single or split application program may be used. All **Guardsman** applications must only be made to sorghum seed that has been properly treated by the seed company with an approved chloroacetamide herbicide safener or severe injury may occur.

Under high soil moisture and/or cool conditions, Guardsman application may cause temporary stunting or leaf wrapping of sorghum. Sorghum will normally outgrow these symptoms in 10 to 14 days.

For best performance make preemergence surface applications within 5 days of the last preplant tillage. If weeds are emerged apply **Guardsman** with contact herbicides to control emerged vegetation.

If grasses are emerged **Guardsman** must be applied before they reach the 2 leaf stage or must be used in combination with an effective postemergence herbicide.

Sorghum forage may be grazed or fed to livestock at 60 or more days after application. Grain and fodder may be harvested and fed at 80 or more days after application of **Guardsman**.

Guardsman is not registered for use on sweet or forage sorghum.

GUARDSMAN USE RATES IN SORGHUM

For **Guardsman[®] herbicide** rates in sorghum use rates given for corn in this Directions for Use Booklet. Additional rate recommendations specific for sorghum: for best weed control for sorghum produced under irrigation, use a minimum of 3.5 pt/A. **Guardsman** is not recommended for use on coarse textured soils.

GUARDSMAN TANK MIXTURES AND SEQUENTIAL USE PROGRAMS IN SORGHUM

Guardsman may be applied prior to, in tank mix with, or following use of one or more of the following herbicides in sorghum: atrazine, BANVEL®, Cyclone® CF, Fallowmaster®, Gramoxone® Extra, glyphosate (Roundup®), Landmaster® BW and Mito-Pro. In addition this product can be used prior to or used in sequential applications with other herbicides labeled for use in grain sorghum such as: Basagran®, Buctril®, Laddok® S-12, MARKSMAN, WEEDMASTER® or 2,4-D. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

.....

When using tank mix or sequential applications with **Guardsman**, always read the companion product label(s) to determine specific use rates by soil types, weed species, and weed or crop growth stage. In addition, follow precautions and restrictions including state and local use restrictions that may apply to specific products.

| Guardsman plus ATRAZINE [0.5 to 1.0 lb ai/A] | Applications may be made preplant, preplant incorporated, or preemergence. Do not apply on coarse textured soils or medium textured soils with less than 1.0% organic matter. |
|---|---|
| Guardsman plus MILOPRO [0.60 to 2.3 lb ai/A] | Applications may be made preplant or preemergence in states where Milo-Pro is approved for use. Do not apply on coarse textured soils or medium textured soils with less than 1.5% organic matter. |
| Guerdsman plus CYCLONE CF or GRAMOXONE EXTRA {1.5 to 3.0 pt/A} or ROUNDUP (GLYPHOSATE] {0.5 to 4.0 lb ai/A} or LANDMASTER BW or FALLOWMASTER | Apply for burndown of emerged weeds in minimum or no-tillage production system. Application with Cyclone, Gramoxone, or Roundup must be made before crop emergence. For improved burndown of broadleaf weeds, BANVEL may be added to mixtures with Cyclone, Gramoxone, and Roundup (glyphosate) if appli- cation is made 15 or more days before planting. Landmaster or Fallow Master applications must be made 14 or more days before sorghum seeding. |

MIXING AND APPLICATION DIRECTIONS MIXING OF SPRAY SOLUTIONS

Guardsman is a concentrated liquid herbicide that can be mixed for application in water, sprayable fluid fertilizer, or it may be impregnated on dry bulk fertilizer. When herbicide tank, mixtures or mixtures with sprayable fluid fertilizer are to be used, a compatibility test should be conducted prior to tank mixing (SEE COMPATIBILITY TEST section in this label). When loading a spray tank for application of **Guardsman** (applied alone or in tank mixtures with other products) use the following order for mixing: 1) Thoroughly clean spray tank prior to use.

2) Fill sprayer tank one-half full of either water or sprayable fluid fertilizer.

- 3) Start sprayer agitation system and continue agitating throughout mixing and application.
- 4) If compatibility test indicated the need for addition of a compatibility agent, add the amount recommended on the label of the selected compatibility agent.
- 5) If tank mixing with wettable powder or dry flowable products, make a slurry of these products with water and add slowly to spray tank.
- 6) If tank mixing with a flowable formulation, add this product slowly to spray tank.
- 7) Add Guardsman to spray tank.
- 8) If tank mixing with emulsifiable concentrates or soluble liquid products add these products to spray tank.
- 9) If tank mixing with Gramoxone Extra or Roundup add these products as remainder of spray tank is filled with carrier.
- 10)If adding spray adjuvants, add them after all other products have been mixed.

GROUND OR AERIAL APPLICATIONS

Apply the recommended rates of **Guardsman** in 2 or more gallons of spray solution per acre. Actual minimum spray volume per acre is determined by spray equipment used for application. Application must be made in adequate volume to provide accurate and uniform distribution of spray particles over treated area and to avoid drift of spray particles to non-target areas. In addition, when **Guardsman** is used in tank mixtures, spray volume used must be no less than the minimum volume required by the tank mix product. Use nozzle screens no finer than 50 mesh when spraying tank mixtures with wettable powder or flowable formulations.

Guardsman[®] herbicide may be applied as a band treatment. Use rates within the treated area of the band must **not** exceed the use rates given for broadcast application. Use the formula below to determine the appropriate amount of product and volume of carrier needed per acre of field.

| Band width <u>in inches</u> Row width in inches | ٠X | Broadcast RATE per acre | = | Band RATE per acre |
|--|----|---------------------------------|---|-------------------------|
| Band width in inches Row width in inches | x | Broadcast VOLUME per àcre | = | Band VOLUME per acre |

USE OF SPRAY ADJUVANTS

Spray adjuvants have little or no influence on performance of **Guardsman** when applications are made prior to weed emergence: To improve burndown of emerged weeds; surfactants and/or low rate fertilizer (28%, 30%, or 32% UAN or ammonium sulfate), or crop oil concentrate may be used with **Guardsman** or **Guardsman** tank mixes applied preplant, preemergence, or early postemergence to the crop. Crop oil concentrates are allowed after crop emergence only when **Guardsman** is used alone or in tank mixtures with atrazine.

COMPATIBILITY TESTING

A compatibility test is recommended for applications with liquid fertilizer. Before mixing products in the spray tank, small amounts of all products can be mixed in proportionate quantities to determine compatibilities. The amounts of carrier and products in this compatibility test are based on a spray volume of 25 gallons per acre. Make appropriate changes in amounts for your specific spray volume.

To conduct this test begin by adding one pint of the liquid fertilizer solution to a one quart container. Then add, in sequence as given above, $1^{1}/_{2}$ level teaspoons per lb use rate for dry formulated products and $1/_{2}$ teaspoon per pint use rate of liquid formulated products. Agitate materials by covering and gently shaking container for 5 to 10 seconds between each product addition.

If herbicide(s) do not ball-up or form flakes, gels, sludge, oily film or layers, or other precipitates, then the tested spray mix is compatible. Incompatibility in any of the above described forms will normally occur within 5 minutes after mixing. If components are incompatible, the use of a compatibility agent is recommended. Rerun the above compatibility test with a suitable compatibility agent (1/4 teaspoon in test is equivalent to a use rate of 2 pints per 100 gallons of spray volume). If the mixture is then compatible, use the compatibility agent as directed on its label. If the mixture is still incompatible, the liquid fertilizer and herbicide(s) should not be mixed for use in the same spray tank.

DRY BULK FERTILIZER APPLICATIONS

Guardsman may be impregnated or coated onto dry bulk granular fertilizer carriers for preplant surface, preplant incorporated or preemergence applications. Impregnation or coating may be conducted by either the in-plant bulk system or the onboard system. Guardsman may also be applied in herbicide tank mixtures where the tank mix companion product is also registered for these application systems. Individuals or agents selling Guardsman in these application systems are responsible for following all state and local regulations regarding fertilizer/herbicide blending.

Addition of a drying agent may be necessary if the fertilizer/herbicide blend is too wet for uniform application due to high humidity, high urea concentration, and/or low fertilizer use rate. Slowly add the drying agent to the blend until a flowable mixture is obtained. Drying agents are not recommended for use with on-board impregnation systems.

Under some conditions fertilizer impregnated with Guardsman may clog air tubes or deflector plates on pneumatic application systems. Mineral oil may be added to Guardsman before blending with fertilizer to reduce plugging. Do not use drying agents when mineral oil is used. To avoid separation of Guardsman and mineral cil mixes in cold temperatures either keep mixture heated or agitated prior to blending with fertilizer. Mineral cil may be used at in-plant blending stations or on-board injection systems.

7

~ . 뷁는 2월 화 파트 반의 보니 같~

Apply 200 to 750 pounds of fertilizer/herbicide blend per acre. Application must be made uniformly to the soil to prevent possible crop injury and offer satisfactory weed control. Impregnated fertilizer spread at half rate and overlapped to obtain a full rate will offer a more uniform distribution. A shallow (1 to 2 inch) incorporation is desirable for improved weed control. Deeper incorporation may result in unsatisfactory weed control.

Formula to Determine Herbicide Rate(s):

Pints or Pints or <u>Pounds Herbicide/Acre</u> X 2000 = Pounds of Herbicide per Pounds of Fertilizer/Acre Ton of Fertilizer

INCOMPATIBLE MIXTURES - DO NOT impregnate Guardsman or Guardsman mixes on ammonium nitrate, potassium nitrate, or sodium nitrate fertilizers or fertilizer blends. Single superphosphate (0-20-0) and triple superphosphate (0-46-0), may be impregnated only with Guardsman alone.

ROTATIONAL CROPS

If the corn or grain sorghum crop treated with Guardsman[®] herbicide is lost to adverse weather or for other reasons, the area treated may be replanted to either of these crops immediately. If the original Guardsman treatment was broadcast do not make a second application of **Guardsman**. If the original application was banded and the second crop is planted in the row middles a second band application may be applied. Corn, sorghum, soybeans, cotton or peanuts may be planted the year following treatment. Injury may occur to soybeans planted on soils having a calcareous surface layer. Do not graze or feed forage or fodder from cotton to livestock. Do not plant sugarbeets, tobacco, vegetables (including dry beans), spring-seeded small grains, or small seeded legumes and grasses the year following application, or injury may occur.

Guardsman used in tank mixtures: Refer to the labels of tank mix products for additional rotational crop information and restrictions.

STORAGE AND CONTAINER DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container away from fertilizer, feed, or foodstuffs and separated from other pesticides.

PESTICIDE DISPOSAL: Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures under Subtitle C of the Resource Conservation and Recovery Act.

PLASTIC CONTAINERS

Triple rinse (or equivalent) and add rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If containers are burned, do not inhale smoke. **Metal:** Triple rinse (or equivalent). Then offer for recycling or containers reconditioning, or puncture and dispose of in a sanitary landfill, (non-aerosol) or by other procedures approved by state and local authorities.

Five Steps to Repacking BASF Bulk Containers

Step 1. Inspect all mini-bulk tanks before filling with bulk products.

- Is the tank empty and clean (according to the current approved mini-bulk tank cleanout procedure), or does it contain only residue of the same BASF product with which it is about to be filled?
- Does the tank have a capacity greater than 55 gallons?
- Has the tank been inspected to be sure it is free of any punctures or structural defects?
- Step 2. Inspect all tank valves, hoses, pumps, meter, and seals before filling with bulk product.
- Are all fittings free of visual signs of leaking or heavy wear?
- Are all fittings and plumbing clean (according to the current approved mini-bulk tank cleanout procedure)?

Step 3. Label all tanks properly before filling with bulk product.

- Is there an up-to-date legible product label including Directions For Use displayed on the tank.
- Did you write the net contents (in gallons) on the product label after every refill.
- Did you write your EPA Establishment Number on the product label?
- Step 4. Properly secure the tank to any vehicle prior to transportation.
- Step 5. Maintain a file of the following items at the location where repackaging occurs.
 - A copy of the manufacturer's cleanout procedure.
 - A signed copy of the manufacturer's repackaging label authorization for this retail location.
 - While it is not required by the EPA currently, BASF has developed mini-bulk checklist to aid in your inspection (Steps 1-3 above). We recommend that you complete these checklists every time a mini-bulk container is refilled and keep them on file.

Please post these instructions in an areavisible by all employees and be sure to follow them prior to filling any mini-bulk tank.

>>>>>>

33333

,,,,

Conditions of Sale and Warranty

10 %

10

÷

The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop. injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Selier. All such risks shall be assumed by the Buyer. BASF warrants that this product conforms to the

chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use, subject to the inherent risks, referred to above. BASF MÁKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

TRADEMARKS

Accent and Bladex are registered tracemarks of E.I. duPont de Nemours & Company.

Basagran and Laddok are registered trademarks of BASF AG. Beacon and Princep are registered trademarks of Novartis, Buctril is a registered trademark of Prone-Poulenc Ag Company.

Cyclone and Gramoxone are registered trademarks of Zeneca, Inc

Landmaster and Roundup are trademarks of Monsanto Company.

Prowl and Pursuit are registered tracemarks c* American

Cyanamid Company, BANVEL, CLARITY, FRONTIER, GUARDSMAN, FALLOWMASTER, and WEEDMASTER are registered trademarks of BASF Corporation.

Laddok is a restricted use pesticide.

© 1997 BASF Corporation All rights reserved. NVA 97-4-67-0118 bulk

BASF Corporation P.O. Box 13528 Research Triangle Park, NC 27709

