

PM 25.

7969-168

4/27/98

BASF

RT Date: 2-22-98
Copy 2f

1/8

ACCEPTED

APR 27 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 7969-168

BM OK

TJ OK

MK OK

KB OK

Conclude® Ultra

herbicide

For use on soybeans

Active Ingredients*:

Sodium salt of bentazon: sodium (3-isopropyl-1H-2,1,3-benzothiadiazin-4(3H)-one-2,2-dioxide).....	20.0%
Sodium salt of acifluorfen: sodium 5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoate	9.1%
Sethoxydim: 2-[1-(ethoxyimino)butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one*	14.0%
Inert Ingredients	56.9%
Total	100.0%

* Equivalent to 1.69 pounds per gallon bentazon: 3-isopropyl-1H-2,1,3-benzothiadiazin-4(3H)-one 2,2-dioxide; and 0.84 pounds per gallon of sodium acifluorfen, sodium 5-[2-chloro-4-(trifluoromethyl) phenoxy]-2-nitrobenzoate.

* Equivalent to 1.29 pounds of sethoxydim per gallon

EPA Reg. No. 7969- RAI

EPA Establishment Number

KEEP OUT OF REACH OF CHILDREN.

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See the attached booklet for complete **Precautionary Statements, Statements of Practical Treatment, Directions For Use, and Conditions of Sale and Warranty.**

Net contents:

Precautionary Statements

Hazards to Humans and Domestic Animals

DANGER. Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Avoid contact with skin. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Statement of Practical Treatment

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

If swallowed: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol.

If inhaled: Move to fresh air. Aid in breathing if necessary, and get immediate medical attention.

Note to physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
 - Chemical-resistant gloves such as barrier laminate, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, or viton \geq 14 mils
 - Chemical-resistant footwear plus socks
 - Protective eyewear
 - Chemical-resistant headgear for overhead exposure
- Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to aquatic organisms. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. Do not apply when weather conditions favor drift from target area.

Groundwater Advisory

Bentazon and acifluorfen are present in this product. These chemicals are known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this product in areas where soils are permeable, such as sand and soils with loamy sand textures, particularly where water tables are shallow could result in contamination of groundwater. The utilization of irrigated water in these areas will increase the likelihood of contamination.

Endangered Species Concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of federal law.

Tank Mix of Conclude® Ultra B and Conclude® Ultra G herbicides

(Hereafter referred to as Conclude® Ultra)

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty are to be followed. This labeling must be in the user's possession during application.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, or viton \geq 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Do not store below 32° F or above 100° F. Store in a dry place away from heat or open flame. Avoid contamination of feed or foodstuffs.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal:

• **Plastic Containers for Duplex® II system:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

• **Bulk/Mini-bulk Containers:**

Reusable containers should be returned to the point of purchase for cleaning and refilling. Reusable containers can only be refilled with **Conclude® Ultra herbicide**. Do not reuse this container with any other product.

In Case of Emergency

In case of large-scale spillage regarding this product, call:

CHEMTREC 800-424-9300
BASF Corporation 800-832-HELP

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment.
- Your local poison control center (hospital).
- BASF Corporation (800-832-HELP).

Returnable Container Operating Instructions.

Prodigy® System Operating Procedure

Conclude Ultra is supplied in the **Prodigy System**, a unique, 120-gallon mini-bulk closed delivery system that contains enough product to treat 320 acres. It consists of a self-discharging tank that does not require any pumping mechanism, and has a dry lock connector which protects the user from exposure to tank contents.

Do not refill **Prodigy System**. Return **Prodigy System** to BASF for cleaning and refilling.

Conclude Ultra in a dedicated, returnable **Prodigy System** can only be used with the closed **Prodigy System** in which it comes packaged.

The **Prodigy System**, when operated according to directions, will discharge **Conclude Ultra G** and **Conclude Ultra B** in a 1:1.7 ratio.

Attention! The **Prodigy System** is a pressurized delivery system. Do not attempt to open the container. Transfer product only by following these steps:

1. Install a male dry lock connector to the spray tank.
2. Uncoil the hose from the rack and connect the female dry lock connector (at the end of the hose attached to the tank) with the male dry lock connector installed on the spray tank.
3. Turn on the nitrogen gas supply.

4. Push down on the activation handle in the front near the meter until the handle is locked in the lower position allowing the manifold to fill with product and become pressurized. Some tanks do not have a handle; move on to the next step.
5. Turn the meter on by pressing the "ON/TOTAL" button.
6. Press "RESET" button to set current total to "0.00" if desired.
7. Turn the yellow product delivery valve counterclockwise (to horizontal) until the desired amount of product, as indicated on the measuring meter, has been discharged into the spray tank.
8. Turn the yellow product delivery valve clockwise (to vertical) to stop the discharge of product into your spray tank.
9. Lift the activation handle to the unlocked position (in front near the meter) to stop liquid and pressurization from flowing into the manifold. Some tanks do not have a handle; move on to the next step.
10. Turn off the nitrogen gas valve when the **Prodigy System** is not in use.
11. **Hose draining:** Starting at the yellow handle on the **Prodigy Tank**, grasp the hose and walk toward the receiving tank holding the hose level or higher than the dry lock connection allowing all of the product to drain out of the hose.
12. Disconnect the female dry lock connector on the tank hose from the male dry lock connector on the spray tank.
13. Recoil the hose onto the hose rack.
14. Be sure to turn off the nitrogen gas valve on the nitrogen cylinder when the **Prodigy System** operation is completed, or when the tank is empty, or when the tank is ready to be returned to the point of purchase.


Leave all product and bar code labels in place. Product labels must remain in place to comply with Department of Transportation regulations.

Return Container Promptly to Distributor

The **Prodigy System** containers are tracked with bar codes and serial numbers. Distributors are responsible for the containers assigned to them. Return this container to the distributor from which it was purchased. Notify the distributor if the container cannot be returned by a specific time. The distributor is responsible for returning the container to BASF. The distributor will be charged for any container not returned within 30 days.

Duplex™ II System

Conclude Ultra B and **Conclude Ultra G** are provided in a molded jug pack that contains enough **Conclude Ultra** to treat 5 acres.



I. General Information

Conclude® Ultra herbicide is intended for the early postemergence control of a wide spectrum of broadleaf weeds and annual grasses in soybeans (see **Table 1**). **Conclude Ultra B** must be used in combination with **Conclude Ultra G**.

Mode of Action

Conclude Ultra rapidly enters the target weed through its foliage and translocates throughout the plant. The effects range from slowing or stopping growth (generally within 2 days), to foliage reddening and leaf tip burn. Subsequently, foliage burnback may occur. These symptoms will generally be observed within 3 weeks depending on environmental conditions.

Crop Tolerance

All soybean varieties are tolerant to **Conclude Ultra** at all stages of growth. Leaf speckling may occur, but plants generally outgrow this condition within 10 days. New growth is normal and crop vigor is not reduced.

Herbicide Resistance

Repeated use of **Conclude Ultra** (or similar postemergence grass herbicides with the same mode of action) may lead to the selection of naturally occurring biotypes with resistance to these products. If poor performance cannot be attributed to adverse weather conditions or improper application methods, a resistant biotype may be present. Consult your local representative or agricultural advisor for assistance.

Coverage

Apply **Conclude Ultra** to the foliage of grasses on a spray-to-wet basis uniformly and completely because large leaf canopies shelter smaller weeds and can prevent adequate spray coverage. Do not spray to the point of runoff.

Irrigation

In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth.

Cultivation

Do not cultivate within 5 days before or 7 days after applying **Conclude Ultra**. Cultivating 7-14 days after treatment may help provide season-long control.

Cleaning Spray Equipment

Clean spray equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product, particularly if a herbicide with the potential to injure crops was used.

II. Application Instructions

For optimum results with **Conclude Ultra** in a total postemergence, one-pass weed control system for soybeans, the following recommendations should be followed:

- plant rows 15" wide or less
- eliminate all vegetative weed growth prior to soybean planting
- apply **Conclude Ultra** according to weed sizes stated on this label (about 21 days after soybean planting).

4/8
Apply 38 ounces of **Conclude Ultra** (14 ounces of **Conclude Ultra G** and 24 ounces of **Conclude Ultra B**) per acre early postemergence (generally when soybeans are in the 2nd to 3rd trifoliate leaf stage of growth). The most effective control will result from making postemergent applications of **Conclude Ultra** early, when weeds are small. Delaying application permits weeds to exceed the maximum size stated and will prevent adequate control.

Do not apply when conditions favor drift from target area or when windspeed is greater than 10 mph.

All **Conclude Ultra** applications to control volunteer cereals (barley, corn, oats, rye, and wheat) should be made before tillering. Volunteer cereals that emerged the previous fall may not be adequately controlled with **Conclude Ultra** for spring control.

Air Application

Water Volume: Use a minimum of 5 gallons of water per acre.

Spray Pressure: Use up to 40 psi.

Application Equipment: Use only diaphragm-type nozzles that produce fan spray patterns.

Nozzle Height: 6-10 feet above crop.

Nozzle Orientation: Nozzles must be oriented so as to discharge straight back with the air stream (opposite the direction of travel of the aircraft) and not more than 20° downwind. Nozzles must be located no farther out than 3/4 the distance from the center of the aircraft to the end of the wing or rotor.

Special Directions for Aerial Application

To obtain uniform coverage and to avoid drift hazards, follow these guidelines:

- Do not apply **Conclude Ultra** by aircraft when wind is blowing more than 10 mph. Use coarse sprays (larger droplets) as they are less likely to drift.
- Do not apply **Conclude Ultra** by air if ornamental or sensitive nontarget crops such as cotton, sugar beets, sunflowers, or okra are within 200 feet downwind.

The applicator must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

Ground Application (Broadcast)

Water Volume: Use a minimum of 10 gallons of water per broadcast acre.

Spray Pressure: Use a minimum of 40 psi (measured at the boom, not at the pump or in the line).

Application Equipment: Use standard high-pressure pesticide flat fan or hollow cone nozzles spaced up to 20 inches apart. Do not use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles as erratic coverage can cause inconsistent weed control. Do not use selective application equipment such as recirculating sprayers or wiper applicators. When tall weeds such as volunteer corn are to be controlled, the boom should be high enough to cover the entire plant. Refer to the nozzle manufacturer's directions for recommended height.

Table 1. Maximum weed heights controlled by 33 ounces of Conclude Ultra per acre (24 ounces of Conclude Ultra B and 14 ounces of Conclude Ultra G per acre) with crop oil concentrate (1 pint per acre)

Broadleaves*	Leaf Stage	Maximum Weed Height	Grasses*	Maximum Weed Height
Anoda, Spurred*	4	2"	Barnyardgrass	4"
Carpetweed	3" diam.	2"	Crabgrass, Large	4"
Cocklebur	6	6"	Smooth	4"
Copperleaf, Hophornbeam	4	4"	Cupgrass, Woolly	4"
Crotalaria	6	6"	Foxtail, Giant	4"
Croton, Tropic	2	2"	Green	4"
Woolly	2	2"	Yellow	4"
Eclipta	6	6"	Goosegrass	4"
Jimsonweed	6	6"	Johnsongrass (seedling)	4"
Ladysthumb	6	6"	Junglegrass	4"
Lambsquarters*	6	2"	Millet, Wild Proso	12"
Mallow, Venice	6	2"	Oats, Wild	3"
Morningglories	4	2"	Panicum, Browntop	4"
Mustard, Wild	6	4"	Fall	4"
Nightshade, Black	6	2"	Texas	4"
Pigweed, Redroot	6	3"	Red Rice	1"
Smooth	6	3"	Shattercane	6"
Ragweed, Common	6	3"	Signalgrass, Broadleaf	4"
Giant	4	6"	Sprangletop, Red	4"
Redweed	4	3"	Volunteer Corn	12"
Sesbania, Hemp	4	6"	Witchgrass	4"
Sida, Prickly or Teaweed*	4	2"		
Smartweed, Pennsylvania	6	6"	Perennials:	Maximum Weed Height
Starbur, Bristly	6	3"	(top growth suppression)	
Texasweed	3	3"	Johnsongrass (Rhizome)	Based on application timing of annual grasses
Velvetleaf*	4	2"	Nutsedge, Yellow*	
Waterhemp, Common	6	3"	Thistle, Canada*	
Tail	6	3"		

* For new germination or perennial regrowth, follow up with Basagran*, Poast Plus*, Storm*, or Blazer* herbicide. Refer to the respective labels for Directions For Use.

† Control may be inconsistent. A later application of Basagran may be necessary. (See Basagran label.)

III. Additives

When using the Prodigy System, add oil concentrate after Conclude* Ultra herbicide has been added to the spray tank. Under excessively dry, wet, or cold conditions which may reduce herbicidal activity, use the maximum rate of oil concentrate (refer to Table 2.)

Oil Concentrate

A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:

- be nonphytotoxic,
- contain only EPA-exempt ingredients,
- provide good mixing quality in the jar test, and
- be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. Refer to Compatibility Test for Mix Components for more information.

Table 2. Additive Rates Per Acre

Additive	Ground Application	Aerial Application
Oil Concentrate	1-2 pints	1-2 pints

Compatibility Test for Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

- 1) Water. For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.

- 2) Correct amounts of Conclude Ultra G and B. Cap the jar and invert 10 cycles.
- 3) Emulsifiable concentrates (oil concentrate). Cap the jar and invert 10 cycles.
- 4) Let the solution stand for 15 minutes.
- 5) Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. Do not use any spray solution that could clog spray nozzles.

IV. Mixing Order

The following steps apply when using the Duplex* II container system.

- 1) Water: Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
- 2) Agitation: Maintain constant agitation throughout mixing and application.
- 3) Conclude Ultra B
- 4) Emulsifiable concentrates (oil concentrate)
- 5) Conclude Ultra G
- 6) Remaining quantity water

Maintain constant agitation during application. Do not attempt to pour the contents of the Duplex II container system into the tank simultaneously or poor mixing will result.

V. Tank Mixing Application

No tank mixes are specifically recommended with this product. Consult your local BASF representative or agricultural advisor for assistance.

6/8

VI. General Restrictions and Limitations — All Crops

- **Maximum seasonal use rate:** Do not apply more than **38 ounces** of **Conclude® Ultra herbicide** (14 ounces of **Conclude Ultra G** and 24 ounces of **Conclude Ultra B**) per acre, per season.
- **Make only one** application of **Conclude Ultra** per acre, per season.
- After an application of **Conclude Ultra**, do not apply more than 1.5 pounds of a.i. of bentazon per acre, per season; 0.25 pounds of a.i. of sodium acifluorfen per acre, per season; or 0.55 pounds of a.i. of sethoxydim per acre, per season.
- Do not apply sequential applications of **Blazer**, **Galaxy™ herbicide**, or **Storm** within 15 days following the application of **Conclude Ultra**.
- **Preharvest Interval (PHI):** Do not apply within **75 days** of soybean harvest.
- **Restricted Entry Interval (REI): 48 hours.**
- In case of **crop failure**, only peanuts or soybeans may be immediately replanted.
- **Crop Rotation Restriction:** Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in fields treated with **Conclude Ultra** for **18 months** following treatment.
- Do not use treated plants for feed or forage.
- Do not apply to grasses or crops under **stress** such as stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control will probably result.
- Do not apply to crops that show **injury** (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced or prolonged.
- Avoid drift to all other crops and nontarget areas.
- Do not use **selective application equipment** such as recirculating sprayers, wiper applicators, or shielded applicators.
- **Rainfast Period:** Do not apply if rainfall or overhead irrigation is expected soon after application.
- Do not apply through any type of **irrigation** equipment.
- This product cannot be used to **formulate** or reformulate any other pesticide product.

Table 3. Crop-Specific Restrictions and Limitations for Conclude Ultra herbicides

Crop	Minimum Time From Application to Harvest (PHI)	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application
Soybeans	75 days	38 ounces	38 ounces	No	Yes

VII. Crop-Specific Information

Soybeans

Special Directions for Problem Weeds

Rhizome Johnsongrass:

Rhizome Johnsongrass is best controlled when **Conclude® Ultra** herbicide is followed by **Poast Plus®** herbicide when Johnsongrass is 6-8" tall. The timing of **Conclude Ultra** should follow label directions for control of the annual grasses and broadleaf weeds. This timing will usually not be optimum for controlling rhizome Johnsongrass, however, the **Conclude Ultra** will provide effective control of the Johnsongrass vegetation and some rhizomes.

The sequential application of **Poast Plus** (1.5 pints per acre) will control the newly emerging vegetation as well as deplete the rhizome reserves when the following rates are used:

- **Conclude Ultra:** 38 fluid ounces per acre based on the annual grass labeled directions.
- **Poast Plus:** 1.5 pints per acre on 6-8" rhizome Johnsongrass

Use 2 pints of oil concentrate per acre in each spray mix.

Yellow Nutsedge:

At the optimum application time of **Conclude Ultra** for most pests, yellow nutsedge may not be at the correct growth stage for optimum control. The best nutsedge control will be achieved by applying **Conclude Ultra** and a sequential application of **Basagran®** herbicide at 1.5-2.0 pints per acre.

Crops:

This product can be used on the following crops:

Soybeans

Look inside for complete **Restrictions and Limitations and Application Instructions.**

Weeds listed in this label:

Common Name	Scientific Name
Anoda, Spurred	<i>Anoda cristata</i>
Barnyardgrass (Watergrass)	<i>Echinochloa crus-galli</i>
Carpetweed	<i>Mollugo verticillata</i>
Cocklebur	<i>Xanthium strumarium</i>
Copperleaf, Hophornbeam	<i>Acalypha ostryifolia</i>
Crabgrass, Large	<i>Digitaria sanguinalis</i>
Smooth	<i>Digitaria ischaemum</i>
Crotalaria	<i>Crotalaria spectabilis</i>
Croton, Tropic	<i>Croton glandulosus</i>
Woolly	<i>Croton capitatus</i>
Cupgrass, Woolly	<i>Eriochloa villosa</i>
Eclipta	<i>Eclipta alba</i>
Foxtail, Giant	<i>Setaria faberi</i>
Green	<i>Setaria viridis</i>
Yellow	<i>Setaria glauca</i>
Goosegrass	<i>Eleusine indica</i>
Jimsonweed	<i>Datura stramonium</i>
Johnsongrass	<i>Sorghum halepense</i>
Junglegrass	<i>Cenchrus incertus</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, Common	<i>Chenopodium album</i>
Mallow, Venice	<i>Hibiscus trionum</i>
Millet, Wild Proso	<i>Panicum miliaceum</i>
Morningglory, Cypressvine	<i>Ipomoea quamoclit</i>
Entireleaf	<i>Ipomoea hederacea, var. integruscula</i>
Ivyleaf	<i>Ipomoea hederacea</i>
Palmleaf	<i>Ipomoea wrightii</i>
Pitted	<i>Ipomoea lacunosa</i>
Purple	<i>Ipomoea turbinate</i>
Mustard, Wild	<i>Sinapis arvensis</i>
Nightshade, Black	<i>Solanum nigrum</i>
Nutsedge, Yellow	<i>Cyperus esculentus</i>
Oats, Wild	<i>Avena fatua</i>
Panicum, Browntop	<i>Panicum fasciculatum</i>
Fall	<i>Panicum dichotomiflorum</i>
Texas	<i>Panicum texanum</i>
Pigweed, Redroot	<i>Amaranthus retroflexus</i>
Smooth	<i>Amaranthus hybridus</i>
Ragweed, Common	<i>Ambrosia artemisiifolia</i>
Giant	<i>Ambrosia trifida</i>
Red Rice	<i>Oryza sativa</i>
Redweed	<i>Melochia corchorifolia</i>
Sesbania, Hemp	<i>Sesbania exaltata</i>
Shattercane/Wildcane	<i>Sorghum bicolor</i>
Sida, Prickly, (Teaweed)	<i>Sida spinosa</i>
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Sprangletop, Red	<i>Leptochloa filiformis</i>
Starbur, Bristly	<i>Acanthospermum hispidum</i>
Texasweed	<i>Cyperonia palustris</i>
Thistle, Canada	<i>Cirsium arvense</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Volunteer Corn	<i>Zea mays</i>
Waterhemp, Common	<i>Amaranthus rudis</i>
Tall	<i>Amaranthus tuberculatus</i>
Witchgrass	<i>Panicum capillare</i>

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF Corporation ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. BASF **MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.** BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Basagran, Galaxy, and Poast are registered trademarks of BASF AG.

Blazer, CommServ, Conclude, Duplex, Poast Plus, Prodigy, and Storm are registered trademarks of BASF Corporation.

© 1998 BASF Corporation
All rights reserved
NVA 98-4-11-0035

Additional Information

For additional information, call BASF's **COMM SERV**® at 1-800-874-0081.

BASF Corporation
P.O. Box 13528
Research Triangle Park, NC 27709

BASF