


7969-179

3-29-2000

1/9

	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460	EPA Reg. Number: 7969-179	Date of Issuance: MAR 29 2000	
	NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration	Term of Issuance: Conditional		Name of Pesticide Product: Conclude Xact
	(under FIFRA, as amended)	Name and Address of Registrant (include ZIP Code): BASF Corporation Agricultural Products P.O. Box 13528 Research Triangle Park, NC 27709-3528		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) or 4(a) when the Agency requires all registrants of similar products to submit such data. 2. Add the phrase "EPA Registration No. 7969-179" before you release the product for shipment. 3. Submit three (3) copies of your final printed labeling before you release the product for shipment. <p>If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of this product constitutes acceptance of these conditions.</p> <p>A stamped copy of the label is enclosed for your records. Vkw</p>				
Signature of Approving Official: 		Date: 3-29-00		

ACCEPTED

MAR 29 2000

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 7964-179

Conclude® Xact

herbicide

For use on soybeans

Active Ingredients:

Sodium salt of bentazon: sodium (3-isopropyl-1H-2, 1,3-benzothiadiazin-4(3H)-one-2,2-dioxide)	16.28%
Sodium salt of acifluorfen: sodium 5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoate	7.47%
Sethoxydim: 2-[1-(ethoxyimino)butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one*	11.15%
Inert Ingredients	65.10%
Total	100.00%

* Equivalent to 2.67 pounds per gallon bentazon: 3-isopropyl-1H-2,1,3-benzothiadiazin-4(3H)-one 2,2-dioxide; and 1.33 pounds per gallon of sodium acifluorfen, sodium 5-[2-chloro-4-(trifluoromethyl) phenoxy]-2-nitrobenzoate.
* Equivalent to 2.0 pounds of sethoxydim per gallon

EPA Reg. No. 7969-

EPA Establishment Number

**KEEP OUT OF REACH OF CHILDREN.
DANGER/PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See the attached booklet for complete **Precautionary Statements, Statements of Practical Treatment, Directions For Use, and Conditions of Sale and Warranty.**

Net contents:

Precautionary Statements

Hazards to Humans and Domestic Animals

DANGER. Corrosive. Causes irreversible eye damage. Do not get in eyes or on clothing. Avoid contact with skin. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Statement of Practical Treatment

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Note to physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. For more options, refer to category **G** on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or viton ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, and loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow manufacturer's instructions for cleaning/maintaining. E. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is toxic to aquatic organisms. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. Do not apply when weather conditions favor drift from target area.

Groundwater Advisory

Bentazon and acifluorfen are present in this product. These chemicals are known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this product in areas where soils are permeable, such as sand and soils with loamy sand textures, particularly where water tables are shallow could result in contamination of groundwater. The utilization of irrigated water in these areas will increase the likelihood of contamination.

Endangered Species Concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of federal law.

Tank Mix of Conclude® Xact B and Conclude® Xact G herbicides

(Hereafter referred to as **Conclude® Xact**)

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** are to be followed. This labeling must be in the user's possession during application.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **48 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or viton ≥ 14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Do not store below 32° F or above 100° F. Store in a dry place away from heat or open flame. Avoid contamination of feed or foodstuffs.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal:

• **Plastic Containers for Duplex® II system:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

• Bulk/Mini-bulk Containers:

Reusable containers should be returned to the point of purchase for cleaning and refilling. Reusable containers can only be refilled with **Conclude® Xact herbicide**. Do not reuse this container with any other product.

In Case of Emergency

In case of large-scale spillage regarding this product, call:

CHEMTREC 800-424-9300
BASF Corporation 800-832-HELP

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment.
- Your local poison control center (hospital).
- BASF Corporation (800-832-HELP).

Duplex® II System

Conclude Xact B and **Conclude Xact G** are provided in a molded jug pack that contains enough **Conclude Xact** to treat 5 acres.

I. General Information

Conclude® Xact herbicide is intended for the early postemergence control of a wide spectrum of broadleaf weeds and annual grasses in soybeans (see **Table 1**). **Conclude Xact B** must be used in combination with **Conclude Xact G**.

Mode of Action

Conclude Xact rapidly enters the target weed through its foliage and translocates throughout the plant. The effects range from slowing or stopping growth (generally within 2 days), to foliage reddening and leaf tip burn. Subsequently, foliage burnback may occur. These symptoms will generally be observed within 3 weeks depending on environmental conditions.

Crop Tolerance

All soybean varieties are tolerant to **Conclude Xact** at all stages of growth. Leaf speckling may occur, but plants generally outgrow this condition within 10 days. New growth is normal and crop vigor is not reduced. 4/9

Herbicide Resistance

Repeated use of **Conclude Xact** (or similar postemergence grass herbicides with the same mode of action) may lead to the selection of naturally occurring biotypes with resistance to these products. If poor performance cannot be attributed to adverse weather conditions or improper application methods, a resistant biotype may be present. Consult your local representative or agricultural advisor for assistance.

Coverage

Apply **Conclude Xact** to the foliage of grasses on a spray-to-wet basis uniformly and completely because large leaf canopies shelter smaller weeds and can prevent adequate spray coverage. Do not spray to the point of runoff.

Irrigation

In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth.

Cultivation

Do not cultivate within 5 days before or 7 days after applying **Conclude Xact**. Cultivating 7-14 days after treatment may help provide season-long control.

Cleaning Spray Equipment

Clean spray equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product, particularly if a herbicide with the potential to injure crops was used.

II. Application Instructions

For optimum results with **Conclude Xact** in a total postemergence, one-pass weed control system for soybeans, the following recommendations should be followed:

- plant rows 15" wide or less
- eliminate all vegetative weed growth prior to soybean planting
- apply **Conclude Xact** according to weed sizes stated on this label (about 21 days after soybean planting).

Apply 3 pints of **Conclude Xact** (1.5 pints of **Conclude Xact G** and 1.5 pints of **Conclude Xact B**) per acre early postemergence (generally when soybeans are in the 2nd to 3rd trifoliate leaf stage of growth). The most effective control will result from making postemergent applications of **Conclude Xact** early, when weeds are small. Delaying application permits weeds to exceed the maximum size stated and will prevent adequate control.

Do not apply when conditions favor drift from target area or when windspeed is greater than 10 mph.

All **Conclude Xact** applications to control volunteer cereals (barley, corn, oats, rye, and wheat) should be made before tillering. Volunteer cereals that emerged the previous fall may not be adequately controlled with **Conclude Xact** for spring control.

Air Application

Water Volume: Use a minimum of 5 gallons of water per acre.

Spray Pressure: Use up to 40 psi.

Application Equipment: Use only diaphragm-type nozzles that produce fan spray patterns.

Nozzle Height: 6-10 feet above crop.

Nozzle Orientation: Nozzles must be oriented so as to discharge straight back with the air stream (opposite the direction of travel of the aircraft) and not more than 20° downwind. Nozzles must be located no farther out than $\frac{3}{4}$ the distance from the center of the aircraft to the end of the wing or rotor.

Special Directions for Aerial Application

To obtain uniform coverage and to avoid drift hazards, follow these guidelines:

- Do not apply **Conclude Xact** by aircraft when wind is blowing more than 10 mph. Use coarse sprays (larger droplets) as they are less likely to drift.
- Do not apply **Conclude Xact** by air if ornamental or sensitive nontarget crops such as cotton, sugar beets, sunflowers, or okra are within 200 feet downwind.

The applicator must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

Ground Application (Broadcast)

Water Volume: Use a minimum of 10 gallons of water per broadcast acre.

Spray Pressure: Use a minimum of 40 psi (measured at the boom, not at the pump or in the line).

Application Equipment: Use standard high-pressure pesticide flat fan or hollow cone nozzles spaced up to 20 inches apart. Do not use flood, whirl chamber, or

controlled droplet applicator (CDA) nozzles as erratic coverage can cause inconsistent weed control. Do not use selective application equipment such as recirculating sprayers or wiper applicators. When tall weeds such as volunteer corn are to be controlled, the boom should be high enough to cover the entire plant. Refer to the nozzle manufacturer's directions for recommended height.

5/9

Table 1. Maximum weed heights controlled by 3 pints of Conclude Xact per acre (1.5 pints of Conclude Xact B and 1.5 pints of Conclude Xact G per acre) with crop oil concentrate (1 pint per acre)

6/9

Broadleaves*	Leaf Stage	Maximum Weed Height	Grasses*	Maximum Weed Height
Anoda, Spurred*	4	2"	Barnyardgrass	8"
Carpetweed	3" diam.	2"	Crabgrass, Large	6"
Cocklebur	6	6"	, Smooth	6"
Copperleaf, Hophornbeam	4	4"	Cupgrass, Woolly	8"
Crotalaria	6	6"	Foxtail, Giant	8"
Croton, Tropic	2	<2"	, Green	8"
, Woolly	2	<2"	, Yellow	8"
Eclipta	6	<2"	Goosegrass	6"
Jimsonweed	6	6"	Johnsongrass (seedling)	8"
Ladysthumb	6	6"	Junglegrass	8"
Lambsquarters*	6	2"	Millet, Wild Proso	10"
Mallow, Venice	6	2"	Oats, Wild	4"
Morningglories	4	2"	Panicum, Browntop	8"
Mustard, Wild	6	4"	, Fall	8"
Nightshade, Black	6	2"	, Texas	8"
Pigweed, Redroot	6	3"	Red Rice	4"
, Smooth	6	3"	Shattercane	18"
Ragweed, Common	6	3"	Signalgrass, Broadleaf	8"
, Giant	4	6"	Sprangletop, Red	8"
Redweed	4	3"	Volunteer Corn	20"
Sesbania, Hemp	4	6"	Witchgrass	8"
Sida, Prickly or Teaweed*	4	2"		
Smartweed, Pennsylvania	6	6"	Perennials:	Maximum Weed Height
Starbur, Bristly	6	3"	(top growth suppression)	
Texasweed	3	3"	Johnsongrass (Rhizome)	Based on application timing of annual grasses
Velvetleaf*	4	2"	Nutsedge, Yellow*	
Waterhemp, Common	6	3"	Thistle, Canada*	
, Tall	6	3"		

* For new germination or perennial regrowth, follow up with Basagran*, Poast Plus*, Storm*, or Blazer* herbicide. Refer to the respective labels for Directions For Use.
 † Control may be inconsistent. A later application of Basagran may be necessary. (See Basagran label.)

III. Additives

Add oil concentrate after **Conclude* Xact herbicide** has been added to the spray tank. Under excessively dry, wet, or cold conditions which may reduce herbicidal activity, use the maximum rate of oil concentrate (refer to **Table 2**.)

Oil Concentrate

A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:

- be nonphytotoxic,
- contain only EPA-exempt ingredients,
- provide good mixing quality in the jar test, and
- be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. Refer to **Compatibility Test for Mix Components** for more information.

Table 2. Additive Rates Per Acre

Additive	Ground Application	Aerial Application
Oil Concentrate	1-2 pints	1-2 pints

Compatibility Test for Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

- 1) **Water.** For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.

- 2) Correct amounts of **Conclude Xact G** and **B**. Cap the jar and invert 10 cycles.
- 3) **Emulsifiable concentrates** (oil concentrate). Cap the jar and invert 10 cycles.
- 4) Let the solution stand for 15 minutes.
- 5) **Evaluate** the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. Do not use any spray solution that could clog spray nozzles.

IV. Mixing Order

The following steps apply when using the **Duplex* II container system**.

- 1) **Water:** Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
- 2) **Agitation:** Maintain constant agitation throughout mixing and application.
- 3) **Conclude Xact B**
- 4) **Emulsifiable concentrates** (oil concentrate)
- 5) **Conclude Xact G**
- 6) Remaining quantity water

Maintain constant agitation during application. Do not attempt to pour the contents of the **Duplex II** container system into the tank simultaneously or poor mixing will result.

V. Tank Mixing Application

No tank mixes are specifically recommended with this product. Consult your local BASF representative or agricultural advisor for assistance.

VI. General Restrictions and Limitations — All Crops

7/9

- **Maximum seasonal use rate:** Do not apply more than **3 pints of Conclude® Xact herbicide (1.5 pints of Conclude Xact G and 1.5 pints of Conclude Xact B)** per acre, per season.
- Make only one application of **Conclude Xact** per acre, per season.
- After an application of **Conclude Xact**, do not apply more than 1.5 pounds of a.i. of bentazon per acre, per season; 0.25 pounds of a.i. of sodium acifluorfen per acre, per season; or 0.55 pounds of a.i. of sethoxydim per acre, per season.
- Do not apply sequential applications of **Blazer, Galaxy® herbicide, or Storm** within **15** days following the application of **Conclude Xact**.
- **Preharvest Interval (PHI):** Do not apply within **75 days** of soybean harvest.
- **Restricted Entry Interval (REI): 48 hours.**
- In case of **crop failure**, only peanuts or soybeans may be immediately replanted.
- **Crop Rotation Restriction:** Root crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in fields treated with **Conclude Xact** for **18 months** following treatment.
- Do not use treated plants for feed or forage.
- Do not apply to grasses or crops under **stress** such as stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control will probably result.
- Do not apply to crops that show **injury** (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced or prolonged.
- Avoid drift to all other crops and nontarget areas.
- Do not use **selective application equipment** such as recirculating sprayers, wiper applicators, or **shielded applicators**.
- **Rainfast Period:** Do not apply if rainfall or overhead irrigation is expected soon after application.
- Do not apply through any type of **irrigation** equipment.
- This product cannot be used to **formulate** or reformulate any other pesticide product.

Table 3. Crop-Specific Restrictions and Limitations for Conclude Xact herbicides

Crop	Minimum Time From Application - to Harvest (PHI)	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application
Soybeans	75 days	3 pints	3 pints	No	Yes

VII. Crop-Specific Information

Soybeans

8/9

Special Directions for Problem Weeds

Rhizome Johnsongrass:

Rhizome johnsongrass is best controlled when **Conclude® Xact herbicide** is followed by **Poast Plus® herbicide** when johnsongrass is 6-8" tall. The timing of **Conclude Xact** should follow label directions for control of the annual grasses and broadleaf weeds. This timing will usually not be optimum for controlling rhizome johnsongrass, however, the **Conclude Xact** will provide effective control of the johnsongrass vegetation and some rhizomes.

The sequential application of **Poast Plus** (1.5 pints per acre) will control the newly emerging vegetation as well as deplete the rhizome reserves when the following rates are used:

- **Conclude Xact:** 3 pints per acre based on the annual grass labelled directions.
- **Poast Plus:** 1.5 pints per acre on 6-8" rhizome johnsongrass

Use 2 pints of oil concentrate per acre in each spray mix.

Yellow Nutsedge:

At the optimum application time of **Conclude Xact** for most pests, yellow nutsedge may not be at the correct growth stage for optimum control. The best nutsedge control will be achieved by applying **Conclude Xact** and a sequential application of **Basagran® herbicide** at 1.5-2.0 pints per acre.

Crops:

This product can be used on the following crops:

Soybeans

Look inside for complete **Restrictions and Limitations** and **Application Instructions**.

Weeds listed in this label:

Common Name	Scientific Name
Anoda, Spurred	<i>Anoda cristata</i>
Barnyardgrass (Watergrass)	<i>Echinochloa crus-galli</i>
Carpetweed	<i>Mollugo verticillata</i>
Cocklebur	<i>Xanthium strumarium</i>
Copperleaf, Hophornbeam	<i>Acalypha ostryifolia</i>
Crabgrass, Large	<i>Digitaria sanguinalis</i>
, Smooth	<i>Digitaria ischaemum</i>
Crotalaria	<i>Crotalaria spectabilis</i>
Croton, Tropic	<i>Croton glandulosus</i>
, Woolly	<i>Croton capitatus</i>
Cupgrass, Woolly	<i>Eriochloa villosa</i>
Eclipta	<i>Eclipta alba</i>
Foxtail, Giant	<i>Setaria faberi</i>
, Green	<i>Setaria viridis</i>
, Yellow	<i>Setaria glauca</i>
Goosegrass	<i>Eleusine indica</i>
Jimsonweed	<i>Datura stramonium</i>
Johnsongrass	<i>Sorghum halepense</i>
Junglegrass	<i>Cenchrus incertus</i>
Ladythumb	<i>Polygonum persicaria</i>
Lambsquarters, Common	<i>Chenopodium album</i>
Mallow, Venice	<i>Hibiscus trionum</i>
Millet, Wild Proso	<i>Panicum miliaceum</i>
Morningglory, Cypressvine	<i>Ipomoea quamoclit</i>
, Entireleaf	<i>Ipomoea hederacea, var. integruscula</i>
, Ivyleaf	<i>Ipomoea hederacea</i>
, Palmleaf	<i>Ipomoea wrightii</i>
, Pitted	<i>Ipomoea lacunosa</i>
, Purple	<i>Ipomoea turbinata</i>
Mustard, Wild	<i>Sinapis arvensis</i>
Nightshade, Black	<i>Solanum nigrum</i>
Nutsedge, Yellow	<i>Cyperus esculentus</i>
Oats, Wild	<i>Avena fatua</i>
Panicum, Browntop	<i>Panicum fasciculatum</i>
, Fall	<i>Panicum dichotomiflorum</i>
, Texas	<i>Panicum texanum</i>
Pigweed, Redroot	<i>Amaranthus retroflexus</i>
, Smooth	<i>Amaranthus hybridus</i>
Ragweed, Common	<i>Ambrosia artemisiifolia</i>
, Giant	<i>Ambrosia trifida</i>
Red Rice	<i>Oryza sativa</i>
Redweed	<i>Melochia corchorifolia</i>
Sesbania, Hemp	<i>Sesbania exaltata</i>
Shattercane/Wildcane	<i>Sorghum bicolor</i>
Sida, Prickly, (Teaweed)	<i>Sida spinosa</i>
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Sprangletop, Red	<i>Leptochloa filiformis</i>
Starbur, Bristly	<i>Acanthospermum hispidum</i>
Texasweed	<i>Caperonia palustris</i>
Thistle, Canada	<i>Cirsium arvense</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Volunteer Corn	<i>Zea mays</i>
Waterhemp, Common	<i>Amaranthus rudis</i>
, Tall	<i>Amaranthus tuberculatus</i>
Witchgrass	<i>Panicum capillare</i>

Conditions of Sale and Warranty

9/9

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF Corporation ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL BASF OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Basagran, Galaxy, and Poast are registered trademarks of BASF AG.

Blazer, Conclude, Duplex, Poast Plus and Storm are registered trademarks of BASF Corporation.

© 2000 BASF Corporation
All rights reserved
NVA 99-4-11-0146

Additional Information

For additional information, call BASF at 1-800-551-2767.

BASF Corporation
P.O. Box 13528
Research Triangle Park, NC 27709

BASF