

## **PLEASE NOTE**

**This image contains more than one label  
approved for this product on this date.**

# Endura™

fungicide

## Supplemental Labeling

### For use in soybean

EPA Reg. No. 7969-197

#### Active ingredient

Boscalid, 3-pyridinecarboxamide, 2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl) ..... 70.0%

Inert ingredients ..... 30.0%

Total ..... 100.0%

#### Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation. Refer to the **Endura™ fungicide** main label for precautionary statements, first aid and personal protective equipment requirements. This supplemental label must be in the user's possession during application.

#### General Information

**Endura** provides optimum disease control when applied in a regularly scheduled protective fungicide program and is used in a spray program that rotates fungicides with different modes of action. Refer to the **Endura** main label for general resistance management information and to the crop-specific use recommendations and restrictions found in this label.

#### Application Information

Apply **Endura** according to the rate, timing, resistance management and adjuvant use recommendations in the **Crop Specific Use Directions (Table A)** in this label.

Observe the additional general recommendations on application instructions, additive use and mixing order on the **Endura** main label.

**Endura** may be applied by ground sprayer, aerial equipment or through sprinkler irrigation systems. Refer to the **Endura** main label for specific instructions on these methods.

#### Restrictions and Limitations

**Endura** is not for use in greenhouse or transplant production systems.

Follow the restrictions and limitations outlined in the Crop-Specific **Use Restrictions and Limitations** table (**Table B**) in this label for:

- Minimum pre-harvest interval (PHI)
- Maximum rate per acre
- Maximum number of applications
- Maximum rate per season
- Livestock grazing or feeding restrictions
- Aerial application restrictions




Table A – Endura™ Fungicide Crop-Specific Use Directions

Crop	Target Diseases	Use Rate per Application	Maximum Number of Applications per Season	Maximum Rate per Season	Minimum Time from Application to Harvest (PHI)
Soybean	Alternaria leaf spot ( <i>Alternaria</i> spp.)	3.5 - 5.5 oz. per acre	2	22 oz. per acre	21 days
	<b>Suppression of:</b> Brown spot ( <i>Septoria glycines</i> ) Cercospora blight and leaf spot ( <i>Cercospora kikuchii</i> ) Frogeye leaf spot ( <i>Cercospora sojina</i> ) Rhizoctonia aerial blight ( <i>Rhizoctonia solani</i> )	5.5 oz. per acre			
	<b>Suppression of:</b> White mold ( <i>Sclerotinia sclerotiorum</i> )	5.5 - 11 oz. per acre			
<b>Application Directions:</b> Begin applications of <b>Endura</b> prior to disease development or when conditions are conducive for disease development, and continue on a 7- to 14-day interval. Use the shorter interval when disease pressure is high. <b>Resistance Management:</b> To limit the potential for development of resistance, do not make more than two (2) applications of <b>Endura</b> per season. Do not make more than two (2) sequential applications of <b>Endura</b> before alternating to a labeled fungicide with a different mode of action for at least one (1) application.					

Table B - Endura Use Restrictions and Limitations

Crop	Minimum Time from Application to Harvest (PHI days)	Maximum Rate per Acre per Application (oz.)	Maximum Number of Applications per Season	Maximum Rate per Acre per Season (oz.)	Livestock Grazing or Feeding	Aircraft Application
Soybean	21	11	2	22	No (Forage and Hay)	Yes

Endura is a trademark of BASF.  
© 2003 BASF Corporation  
All rights reserved.

007969-00197.20040402.NVA 2003-04-201-0082.pdf

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709

**BASF**

**NEXT**

**LABEL**

# Endura™

fungicide

## Supplemental Labeling

### For use on pomefruits

#### Active ingredient

Boscalid, 3-pyridinecarboxamide, 2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl) ..... 70.0%

Inert ingredients ..... 30.0%

Total ..... 100.0%

EPA Reg. No. 7969-197

#### Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Refer to the **Endura™ fungicide** main label for precautionary statements, first aid and personal protective equipment requirements. This supplemental label must be in the user's possession during application.

#### General Information

**Endura** provides optimum disease control when applied in a regularly scheduled protective fungicide program and is used in a spray program that rotates fungicides with different modes of action. Refer to the **Endura** main label for general resistance management information and to the crop-specific use recommendations and restrictions found in this label.

#### Application Information

Apply **Endura** according to the rate, timing, resistance management and adjuvant use recommendations in the **Crop Specific Use Directions (Table 1)** in this label.

Observe the additional general recommendations on application instructions, additive use and mixing order on the **Endura** main label.

**Endura** may be applied by ground sprayer, aerial equipment or through sprinkler irrigation systems. Refer to the **Endura** main label for specific instructions on these methods.

#### Restrictions and Limitations

**Endura** is not for use in greenhouse or transplant production systems.

Follow the restrictions and limitations outlined in the crop-specific **Use Restrictions and Limitations table (Table 2)** in this label for:

- Minimum pre-harvest interval (PHI)
- Maximum rate per acre
- Maximum number of applications
- Maximum rate per season
- Livestock grazing or feeding restrictions
- Aerial application restrictions




Table 1 – Endura™ Fungicide Crop-Specific Use Directions

Crop	Target Diseases	Use Rate per Application	Maximum Number of Applications per Season	Maximum Rate per Season	Minimum Time from Application to Harvest (PHI)
<b>Pomefruits Group</b> Apple Pear Oriental pear Quince Crabapple Loquat	Apple scab ( <i>venturia inaequalis</i> ) Pear scab ( <i>Venturia pirina</i> ) Powdery mildew ( <i>Podosphaera leucotricha</i> ) Fly speck ( <i>Zygophiala jamaicensis</i> ) Alternaria blotch ( <i>Alternaria mali</i> )	6.5 oz. per acre	4	26 oz. per acre	0 days
<p><b>Application Directions for scab and powdery mildew:</b> Begin applications of <b>Endura</b> prior to disease development and continue on a 7- to 10-day interval. Use the shorter interval when disease pressure is high.</p> <p><b>Application Directions for fly speck and Alternaria blotch:</b> Begin applications of <b>Endura</b> prior to disease development and continue on a 7- to 14-day interval. Use the shorter interval when disease pressure is high.</p> <p><b>Resistance Management:</b> To limit the potential for development of resistance, do not make more than four (4) applications of <b>Endura</b> per season. Do not make more than two (2) sequential applications of <b>Endura</b> before alternating to a labeled fungicide with a different mode of action.</p>					

Table 2 - Endura Use Restrictions and Limitations

Crop	Minimum Time from Application to Harvest (PHI days)	Maximum Rate per Acre per Application (oz.)	Maximum Number of Applications per Season	Maximum Rate per Acre per Season (oz.)	Livestock Grazing or Feeding	Aircraft Application
<b>Pome fruits<sup>1</sup></b> Apple Pear, etc.	0	6.5	4	26	Yes	Yes
<sup>1</sup> For a complete list of crops within a crop group, see <b>Table 1- Endura® fungicide Crop-Specific Use Directions</b> .						

Endura is a trademark of BASF.  
© 2003 BASF Corporation  
All rights reserved.

007969-00197.20040402.NVA 2003-04-201-0141.pdf

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709