

7969-197

02/14/2006

1/21

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

Application for Pesticide - Section I

1. Company/Product Number EPA Reg. No. 7969-197	2. EPA Product Manager Tony Kish	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Endura fungicide	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) BASF Corporation, Agricultural Products P.O. Box 13528, 26 Davis Drive Research Triangle Park, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated <u>July 23, 2003</u>
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
FEB 24 2006

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Submission of Final Printed Labeling for Endura fungicide (EPA Reg. No. 7969-197). Minor changes made to the label include change of the BASF Logo, Change of the trademark from Endura™ fungicide to the registered trademark Endura® fungicide, change of the Establishment Number, adding of emergency information on the front panel, minor change to the "Conditions of Sale and Warranty.

PRIA Fee Category: None; Response to EPA request in letter dated July 23, 2003.
 E-Mail: akkarik@basf-corp.com or akkarik@basf.com

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	* Certification must be submitted	
	If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container			
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 6.5 Pounds	5. Location of Label Directions <input checked="" type="checkbox"/> on labeling adhered to jug		
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Khalid H. Akkari	Title Product Registration Manager	Telephone No. (Include Area Code) (919) 547-2976
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment, or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Product Registration Manager	
4. Typed Name Khalid H. Akkari	5. Date February 14, 2006	

2/21

Paperwork Reduction Act Notice and Instructions

Paperwork Reduction Act Notice: Public reporting burden for this collection of information is estimated to average 0.85 hour per response, including time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding the burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Chief, Information Policy Branch, (2136), U.S. Environmental Protection Agency, 401 M Street, SW, Washington, D.C. 20460.

Instructions: This form is to be used for all applications for new registration, end use reregistration, amendment, resubmission, to applications for notifications, final printed labeling, reregistration, etc. In order to process an application for a new registration submitted on this form, the following material must accompany the application:

1. Certification with Respect to Citation of Data (EPA Form 8570-29). [If not exempted by 40 CFR 152.81 (b) (4)];
2. Confidential Statement of Formula (EPA Form 8570-4);
3. Formulator's Exemption Statement (EPA Form 8570-27);
4. Five copies of draft labeling;
5. Three copies of any data submitted;
6. Authorization letter where applicable;
7. Matrices where applicable.

Submission of Labeling - Labeling should first be submitted in the form of draft labels with all applications for new registration. Such draft labels may be in the form of typed label text on 8.5 x 11 inch file. Mockup labels significantly smaller than 8.5 x 11 inches should be mounted on 8.5 x 11 inch paper for submission.

Submission of Data - Data submitted in support of this application must be submitted in accordance with PR Notice 86-5.

Specific Instructions: Please read the instructions listed below before completing this application. First determine the type of registration action, listed in Block A, for which you are submitting this application. For applications submitted in connection with New Registration actions, Sections I, III, and IV must be completed by the applicant. For applications submitted in connection with amended reregistration actions, resubmissions, notifications, reregistrations, etc., Sections I, II, and IV must be completed by the applicant.

Block A - Check the appropriate action for which you are submitting this form.

Section I - This section must be completed, as applicable, for all registration actions.

1. **Company/Product Number** - Insert your Company Number, if one has been assigned by EPA. This number may have been assigned to you as a basic registrant, a distributor, or as an establishment. If your product is registered, insert the Product Number.
2. **EPA Product Manager** - If known, fill in the name and PM number of the EPA Product Manager.
3. **Proposed Classification** - Specify the proposed classification of this product.
4. **Product Name** - Enter the complete product name of this pesticide as it will appear on the label. The name must be specific to this product only. Duplication of names is not permitted among products of the same company. Do not include any brand name or company line designations.
5. **Name and Address of Applicant** - The name of the firm or person and address shown in your application is the person or firm to whom the registration will be issued. If you are acting in behalf of another party, you must submit authorization from the party to act for them in registration matters. An applicant not residing in the United States must have an authorized agent residing in the United States to act for them in all registration matters. The name and complete mailing address of such an agent must accompany this application.
6. **Expedited Review** - FIFRA section 3(c) 3(b) provides for expedited review of applications for registration, or amendments to existing registrations, that are similar, or identical to other pesticide products that are currently registered with the EPA. In order for you application to be eligible for expedited review, you must provide us with the EPA Registration Number and product name of the product you believe is similar to or identical to your product. The product must be similar or identical in both formulation and labeled uses.

Section II - This section must be completed for all applications submitted to amend the registration only of a currently registered product (Amendment), for a resubmission in response to an Agency letter, for notifications to the Agency, for the submission of final printed labeling, for reregistration and for any other action that pertains to a specific EPA-registered product. This section is not to be used for a new application for registration.

1. **Subject of submission** - Check the applicable block and provide the Agency letter date if appropriate. Provide a brief explanation of the purpose(s) for the submission, such as "the addition of a site, pest or crop (specify)"; "amend the Confidential Statement of Formula by . . ."; "reregistration submission"; "general label revision of use directions." Attach a separate page if additional space is needed.

Section III (Packaging and Container Information) - This Section must be completed for all applications submitted in connection with new registration or applicable amendments.

1. **Type of Packaging** - Check the appropriate block if your product will be packaged in the indicated packaging types. Indicate the size of the individual packets and number per retail container.
2. **Type of Retail Container** - Indicate type of container in which product will be marketed.
3. **Location of Net Contents** - Indicate the location of the net contents information for your product.
4. **Size(s) of Retail Container** - Specify the net contents of all retail containers for your product.
5. **Location of the Use Directions** - Indicate the location of the use directions for your product.
6. **Manner in which label is affixed to product** - Indicated the method product label is attached to retail container.

Section IV (Contact Point) - This Section must be completed for all applications for Registration actions, i.e., new products registration, resubmission, "me-too," reregistration, etc.

- 1-5. Self-explanatory
6. EPA Use only

BASF
The Chemical Company

GROUP 7 FUNGICIDE

• **Endura**®

fungicide

For use in dry and succulent beans, bulb vegetables, canola, carrots, fruiting vegetables, grapes, lettuce, peanuts and potatoes

Active Ingredient

Boscalid, 3-pyridinecarboxamide,2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl)..... 70.0%

Inert Ingredients 30.0%

Total 100.0%

[0.7 oz. a.i. (0.044 lb a.i.) in 1 oz. of **Endura**® Fungicide]

EPA Reg. No. 7969-197

EPA Est. No. 67545-AZ-01

KEEP OUT OF REACH OF CHILDREN. WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)


See the attached booklet for complete **First Aid, Precautionary Statements, Directions For Use, and Conditions of Sale and Warranty.**

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net contents: 6.5 pounds

Product of Brazil
BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

NVA 2005-05-201-0317
2064922



FIRST AID

If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

Precautionary Statements

Hazards to Humans and Domestic Animals

Warning. Causes substantial but temporary eye injury. Harmful if absorbed through skin. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Protective eyewear (goggles, face shield or safety glasses)
- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Do not apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow watertables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The require-

ments in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Protective eyewear (goggles, face shield or safety glasses)
- Coveralls
- Chemical-resistant gloves
- Shoes plus socks

Storage and Disposal

Storage:

Do not contaminate water, food, or feed by storage or disposal.

• **Pesticide disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

• **Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

In Case of Spill

In case of large-scale spillage regarding this product, call:

CHEMTREC	800-424-9300
BASF Corporation	800-832-HELP

Steps to be taken in case material is released or spilled:

Dike and contain spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.

Remove contaminated clothing and wash affected skin areas with water.

Wash clothing before re-use.

Keep spill out of all sewers and open bodies of water.

I. General Information

This package contains **Endura® fungicide**, a water dispersable granule (WG). Boscalid, active ingredient in **Endura** belongs to the group of respiration inhibitors classified by the U.S. EPA and Canada PMRA as carboxamides, or Target Site of Action Group 7 fungicides. **Endura** has a new mode of action and is effective against pathogens resistant to other fungicides.

Endura has a protective effect because it inhibits spore germination and a curative-eradivative effect because it inhibits mycelial growth and sporulation of the fungus on the leaf surface. **Endura** can therefore be applied in either pre- or postinfection situations. However, optimum disease control is achieved when **Endura** is applied in a regularly scheduled protective spray program and is used in a rotation program with other fungicides.

Because of its high specific activity and rainfastness, **Endura** has good residual activity against target fungi.

Resistance Management

Endura contains boscalid, a Group 7 fungicide, and is effective against pathogens resistant to fungicides with modes of action different from those of carboxamide (anilide) fungicides (Target site Group 7), such as for example, dicarboximides, sterol inhibitors, benzimidazoles, QoI fungicides, and phenylamides. Fungal isolates resistant to Group 7 fungicides may eventually dominate the fungal population if Group 7 fungicides are

used predominantly and repeatedly in the same field in successive years as the primary method of control for the targeted pathogen species. This may result in reduction of disease control of **Endura** or other Group 7 fungicides.

To maintain the performance of **Endura** in the field, do not exceed the total number of sequential applications of **Endura** and the total number of applications of **Endura** per season stated in Sections V. and VI. Adhere to the label instructions regarding the consecutive use of **Endura** or other target site of action Group 7 fungicides that have a similar site of action on the same pathogens.

The following recommendations may be considered to delay the development of fungicide resistance:

1. **Tank mixtures:** Use tank mixtures with fungicides from different target site of action groups that are registered/permitted for the same use and that are effective against the pathogens of concern. BASF recommends using at least the minimum labeled rates of each fungicide in the tank mix.
2. **IPM: Endura** should be integrated into an overall disease and pest management program. Cultural practices known to reduce disease development should be followed. Consult your local extension specialist, certified crop advisor and/or BASF representative for additional IPM strategies established for your area. **Endura** may be used in Agricultural Extension advisory (disease forecasting) programs, which recommend application timing based on environmental factors favorable for disease development.
3. **Monitoring:** Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development.
4. **Reporting:** If a group 7 target site fungicide, such as **Endura**, appears to be less or no longer effective against a pathogen that it pre-

7/21

viously controlled or suppressed, contact a BASF representative, local extension specialist, or certified crop advisor to assist in determining the cause of reduced performance.

Cleaning Spray Equipment

Spraying equipment must be cleaned thoroughly before and after applying this product, particularly if a product with the potential to injure crops was used prior to **Endura® fungicide**.

II. Application Instructions

Apply recommended rates of **Endura** as instructed by the **Crop-Specific Recommendations**. Apply **Endura** with ground sprayer, aerial equipment or through sprinkler irrigation equipment. Equipment should be checked frequently for calibration. Under low-level disease conditions, the minimum application rates can be used while maximum application rates and shortened spray schedules are recommended for severe or threatening disease conditions. Do not apply when conditions favor drift from target area.

Ground Application: Apply **Endura** in sufficient water to ensure thorough coverage of foliage, bloom, and fruit. Thorough coverage is required for optimum disease control.

Aerial Application: BASF recommends ground applications for thorough coverage. For those crops or in conditions where applications cannot be done by ground equipment, aerial application can be made. Avoid applications under conditions when uniform coverage cannot be obtained or when spray drift may occur. Use no less than 5 gallons of spray solution per acre. For aerial applications to tree and vine crops, use no less than 10 gallons of spray solution per acre. For all other crops, thorough coverage is required for optimum disease control.

Directions for Use Through Sprinkler Irrigation Systems

Sprayer Preparation: Chemical tank and injector system should be thoroughly cleaned. Flush system with clean water.

Application Instructions: Apply **Endura** at rates and timings as described in this label.

Use Precautions for Sprinkler Irrigation Applications:

- Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.
- Add this product to the pesticide supply tank containing sufficient water to maintain a continuous flow by the injection equipment. In continuous moving systems, inject this product-water mixture continuously, applying the labeled rate per acre for that crop. Do not exceed 1/2 inch (13,577 gallons) per acre. In stationary or non-continuous moving systems, inject the product-water mixture in the last 15-30 minutes of each set allowing sufficient time for all of the required pesticide to be applied by all the sprinkler heads and applying the labeled rate per acre for that crop. Do not apply when wind speed favors drift beyond the area intended for treatment. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. Thorough coverage of foliage is required for good control. Good agitation should be maintained during the entire application period.
- If you have questions about calibration you should contact State Extension Service specialist, equipment manufacturers or other experts.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also con-

tain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Specific Instructions for Public Water Systems:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

III. Additives and General Tank Mixing Information

Endura® fungicide can be tank mixed with most recommended fungicides and insecticides herbicides. liquid fertilizers, biological control products, adjuvants, and additives as specified in **Section VI. Crop-Specific Recommendations.**

Under some conditions, the use of additives or adjuvants may improve the performance of **Endura.**

However, all varieties and cultivars have not been tested with possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing **Endura** with other products. Therefore, before using tank mix (fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

Consult a BASF representative or local agricultural authorities for more information concerning additives.

IV. Compatibility Test and Mixing Order

If tank mixtures are used, adhere to restrictions due to rates, label recommendations and precautions on all labels.

Compatibility Test for Tank Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

- 1) **Water:** — For 100 gallons per acre spray volume, use 16 cups (1 gallon) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
- 2) **Water-dispersible products:** — (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions) Cap the jar and

invert 10 cycles.

- 3) **Water-soluble products:** — Cap the jar and invert 10 cycles.
- 4) **Emulsifiable concentrates:** — (oil concentrate or methylated seed oil when applicable) Cap the jar and invert 10 cycles.
- 5) **Water-soluble additives:** — Cap the jar and invert 10 cycles.
- 6) Let the solution stand for 15 minutes.
- 7) **Evaluate** the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. Do not use any spray solution that could clog spray nozzles.

Mixing Order

- 1) **Water:** Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
- 2) **Agitation.** Maintain constant agitation throughout mixing and application.
- 3) **Inductor.** If an inductor is used, rinse it thoroughly after each component has been added.
- 4) **Products in PVA bags:** Place any product contained in water-soluble PVA bags into the mixing tank. Wait until water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
- 5) **Water-dispersible products:** (such as **Endura**, dry flowables, wettable powders, suspension concentrates, or suspo-emulsions).
- 6) **Water-soluble products.**
- 7) **Emulsifiable concentrates:** (such as oil concentrates when applicable).
- 8) **Water-soluble additives:** (such as AMS or UAN when applicable).
- 9) **Remaining quantity water:** Make sure that each component is thoroughly mixed and suspended before adding tank mix partners. Make constant agitation during application. See **Section VI. Crop-Specific Restrictions and Limitations** for more details.

V. General Restrictions and Limitations - All Crops

- **Maximum seasonal use rate:** Do not apply more than the maximum rate per acre as listed in **Table 1. Crop-Specific Restrictions and Limitations** and **Table 2. Crop-Specific Recommendations**.
- **Maximum rate per application:** Do not apply more than the maximum rate per acre per application as listed in **Table 1. Restrictions and Limitations** and **Table 2. Crop-Specific Recommendations**.
- Do not make more than the total number of applications of **Endura® fungicide** per season, as listed in **Table 1. Crop-Specific Restrictions and Limitations** and **Table 2. Crop-Specific Recommendations**.
- **Preharvest Interval (PHI):** See **Table 1. Crop-Specific Restrictions and Limitations** and **Table 2. Crop-Specific Recommendations**.
- **Livestock Feeding Restrictions:** Do not feed peanut hay to livestock.
- Do not apply more than the maximum season use rate of a.i./A for each specific crop from any combination of products (e.g. **Pristine®**, **Endura**, **Emerald®**).
- **Plantback Restrictions:** Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be planted after 14 days.
- Do not use on soybean, cowpea, field pea and grain lupin.
- Do not use on garden beet, radish and turnip.

Table 1. Crop-Specific Restrictions and Limitations

Crop	Minimum Time from Application to Harvest (PHI) (days)	Maximum Rate per Acre per Application (oz. product)	Maximum Number of Applications per Season ²	Maximum Rate per Acre per Season (oz. product)	Livestock Grazing or Feeding
Beans, dry	21	11	2	22	NA ³
Beans, succulent	7	11	2	22	NA
Bulb Vegetables Group ¹ : Onion, Garlic, Leeks	7	6.8	6	41	NA
Canola	21	6	2	12	Yes
Carrots	0	4.5	5	22.5	Yes, for carrot culls
Fruiting Vegetables Group ¹ : Tomato Bell Pepper Chili Pepper Eggplant	0	12.5	2	25	NA

(continued)

Table 1. Crop-Specific Restrictions and Limitations (continued)

Crop	Minimum Time from Application to Harvest (PHI) (days)	Maximum Rate per Acre per Application (oz. product)	Maximum Number of Applications per Season ²	Maximum Rate per Acre per Season (oz. product)	Livestock Grazing or Feeding
Grapes	14	8	3	25	NA
Lettuce	14	11	2	22	NA
Peanut	14	10	3	30	Restricted
Potato	30	10	2	20.5	Yes

¹ For a complete list of crops within a crop group, see Section **VI. Crop-Specific Recommendations**.

² At the maximum use rate only

³ NA = not applicable Aerial Application is permitted for all labeled crop uses.

VI. Crop-Specific Recommendations

Table 2. Crop Specific Recommendation

Crop	Target Diseases	Use Rate per Application	Maximum Number of Applications per Season	Maximum Rate per Season	Minimum Time from Application to Harvest (PHI)
Beans, dried and succulent Broad bean Guar Lablab bean <i>Lupinus spp.</i> Sweet lupin White lupin White sweet Lupin	Ascochyta blight (<i>Phoma exigua</i> , <i>Ascochyta spp.</i>) Botrytis gray mold (<i>Botrytis cinerea</i>) Rust (<i>Uromyces appendiculatus</i>) White mold (<i>Sclerotinia sclerotiorum</i>)	8 to 11 oz. per acre	2	22 oz. per acre	Dried Beans 21 days Succulent Beans 7 days

(continued)

Table 2. Crop-Specific Recommendations (CONT):

Crop	Target Diseases	Use Rate per Application	Maximum Number of Applications per Season	Maximum Rate per Season	Minimum Time from Application to Harvest (PHI)
<p><u>Phaseolus spp.</u> Field bean Kidney bean Pink bean Lima bean Navy bean Pinto bean Runner bean Snap bean Tepary bean Wax bean</p> <p><u>Vigna spp.</u> Adzuki bean Asparagus bean Blackeyed pea Catjang (Chinese) long bean Crowder pea Jack bean Yard long bean Moth bean Mung bean Rice bean Southern pea Urd bean</p>	<p>Ascochyta blight (<i>Phoma exigua</i>, <i>Ascochyta spp.</i>)</p> <p>Botrytis gray mold (<i>Botrytis cinerea</i>)</p> <p>Rust (<i>Uromyces appen- diculatus</i>)</p> <p>White mold (<i>Sclerotinia sclerotiorum</i>)</p>	<p>8 to 11 oz. per acre</p>	<p>2</p>	<p>22 oz. per acre</p>	<p>Dried Beans 21 days</p> <p>Succulent Beans 7 days</p>
<p>Chickpea (garbanzo bean) Lentils</p>	<p>Ascochyta blight (<i>Phoma exigua</i>, <i>Ascochyta spp.</i>)</p> <p>Botrytis gray mold (<i>Botrytis cinerea</i>)</p> <p>Rust (<i>Uromyces appen- diculatus</i>)</p> <p>White mold (<i>Sclerotinia sclerotiorum</i>)</p>	<p>6 oz. per acre</p>			

See next page for instructions.

(continued)

Application Directions: Apply **Endura® fungicide** at the beginning of flowering or prior to onset of disease.

Use the higher rate for extended protection and maximum yield benefit. Apply a second time at full bloom if conditions are favorable for disease development or if heavy disease has already set in.

Ascochyta blight in chickpeas and lentils develops quickly once established so early detection and application is essential to reduce losses.

Apply at the beginning of flowering. Make a second application 7 to 10 days later if disease persists or weather conditions are favorable for disease development.

Resistance management: To limit the potential for development of resistance, do not make more than two (2) applications of **Endura** per season.

Table 2. Crop-Specific Recommendations (CONT):

Crop	Target Diseases	Use Rate per Application	Maximum Number of Applications per Season	Maximum Rate per Season	Minimum Time from Application to Harvest (PHI)
Bulb Vegetables Group Onions (all varieties) Garlic Leek Shallot	Purple blotch (<i>Alternaria porri</i>) Botrytis leaf blight (<i>Botrytis</i> spp.)	6.8 oz. per acre	6	41 oz. per acre	7 days
<p>Application Directions: For control of purple blotch and botrytis leaf blight begin applications of Endura prior to disease development and continue on a 7 to 14-day interval. Use the higher rate and shorter interval when disease pressure is high.</p> <p>Resistance Management: To limit the potential for development of resistance, do not make more than six (6) applications of Endura per season. Do not make more than two (2) applications of Endura before alternating to a labeled fungicide with a different mode of action.</p>					
Canola	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	5 to 6 oz. per acre	2	12 oz. per acre	21 days
<p>Application Directions: Begin applications of Endura at 20 to 50% flowering or prior to the onset of disease. Use the higher rate for extended protection. Apply a second time if conditions continue to be favorable for disease development.</p> <p>Resistance management: To limit the potential for development of resistance, do not make more than two (2) applications of Endura per season.</p>					

(continued)

Table 2. Crop-Specific Recommendations (CONT):

Crop	Target Diseases	Use Rate per Application	Maximum Number of Applications per Season	Maximum Rate per Season	Minimum Time from Application to Harvest (PHI)
Carrots	Alternaria leaf spot (<i>Alternaria</i> spp.)	4.5 oz. per acre	5	22.5 oz. per acre	0 days
<p>Application Directions: Begin applications of Endura® fungicide prior to disease development and continue on a 7 to 14 day interval. Use the shorter interval when disease pressure is high.</p> <p>Resistance Management: To limit the potential for development of resistance, do not make more than five (5) applications of Endura per season. Do not make more than two (2) sequential applications of Endura before alternating to a labeled fungicide with a different mode of action.</p>					
Fruiting Vegetable Group Eggplant Ground cherry Pepino Pepper (all varieties) Tomatillo Tomato	Early blight (<i>Alternaria solani</i>)	2.5 to 3.5 oz. per acre or 2.5 to 3.5 oz. per 100 gallons of spray volume (dilute)*	6	21 oz. per acre	0 days
Tomato	Botrytis gray mold (<i>Botrytis cinerea</i>)	9 to 12.5 oz. per acre	2	25 oz. per acre	<p>Application Directions: Begin applications of Endura prior to disease development and continue on a 7 to 14 day interval for early blight and Botrytis gray mold. Use the higher rate and the shorter interval when disease pressure is high.</p> <p>*For applications based on dilute volume, plants should be sprayed to runoff. Apply a minimum of 20 gallons of spray volume per acre, and increase the spray volume as the plants grow during the season. Spray volume should be proportional to the amount of plant tissue to be covered such that 100 gallons of spray per acre is used on mature plants.</p> <p>Resistance Management: To limit the potential for development of resistance, do not make more than six (6) applications of Endura per season. Do not make more than two (2) sequential applications of Endura before alternating to a labeled fungicide with a different mode of action.</p>

(continued)

Table 2. Crop-Specific Recommendations (CONT):

Crop	Target Diseases	Use Rate per Application	Maximum Number of Applications per Season	Maximum Rate per Season	Minimum Time from Application to Harvest (PHI)
Grapes	Powdery mildew (<i>Uncinula necator</i>)	4.5 oz. per acre	5	25 oz. per acre	14 days
	Botrytis gray mold (<i>Botrytis cinerea</i>)	8 oz. per acre	3		
<p>Application Directions: For powdery mildew control, begin applications of Endura® fungicide at budbreak or prior to the onset of disease and continue on a 10 to 14 day interval. For the control of Botrytis gray mold begin applications of Endura prior to disease development and when conditions favor disease development during early bloom, bunch pre-closure or veraison.</p> <p>Use the higher rate and the shorter interval when disease pressure is high.</p> <p>Resistance Management: To limit the potential for development of resistance, do not make more than five (5) applications of Endura per season.</p> <p>Do not make more than three (3) sequential applications of Endura before alternating to a labeled fungicide with a different mode of action.</p>					
Lettuce, head and leaf	Lettuce drop (<i>Sclerotinia minor</i> , <i>S. Sclerotiorum</i>) Botrytis rot (<i>Botrytis cinerea</i>) Suppression Only Rhizoctonia bottom rot (<i>Rhizoctonia solani</i>)	8 to 11 oz. per acre	2	22 oz. per acre	14 days
<p>Application Directions: On direct-seeded lettuce, make the first application immediately after emergence or prior to disease development.</p> <p>On transplanted lettuce, make the first application immediately after transplanting or prior to the onset of disease. If conditions continue to favor disease development, or if soil surface is disturbed by cultivation, make a second application.</p> <p>Use the higher rate when disease pressure is high.</p> <p>Resistance management: To limit the potential for development of resistance, do not make more than two (2) applications of Endura per season.</p>					

(continued)

Table 2. Crop-Specific Recommendations (CONT):

Crop	Target Diseases	Use Rate per Application	Maximum Number of Applications per Season	Maximum Rate per Season	Minimum Time from Application to Harvest (PHI)
Peanut	Early leaf spot (<i>Cercospora arachidicola</i>) Late leaf spot (<i>Cercosporidium personatum</i>)	6.5 to 10 oz. per acre	3	30 oz. per acre	14 days
	Sclerotinia blight (<i>Sclerotinia minor</i>) Web blotch (<i>Phoma arachidicola</i>) Suppression Only Sclerotium stem rot - Southern stem rot (<i>Sclerotium rolfsii</i>)	8 to 10 oz. per acre			
<p>Application Directions: For control of early and late leaf spot, and web blotch, begin applications of Endura® fungicide prior to the onset of disease and continue on a 14 day interval.</p> <p>For control of Sclerotinia blight, begin applications of Endura prior to the onset of disease or at 45 to 60 days after planting. Make a second application 14 to 21 days later.</p> <p>For suppression of Southern stem rot (<i>Sclerotium rolfsii</i>), apply Endura prior to disease development or at 45 to 60 days after planting. Two additional applications may be made at 14 day intervals. For improved control of Southern stem rot, Endura may be mixed with other labeled, effective fungicides.</p> <p>Use the higher rate and/or shorter spray interval when disease pressure is high or in fields with a history of disease.</p> <p>Do not feed peanut hay to livestock.</p> <p>Resistance Management: To limit the potential for development or resistance, do not make more than three (3) applications of Endura per season.</p> <p>Do not make more than two (2) sequential applications of Endura before alternating to a labeled fungicide with a different mode of action.</p>					

(continued)

17/21

Table 2. Crop-Specific Recommendations (CONT):

Crop	Target Diseases	Use Rate per Application	Maximum Number of Applications per Season	Maximum Rate per Season	Minimum Time from Application to Harvest (PHI)
Potato	Early blight (<i>Alternaria</i> spp.)	2.5 to 4.5 oz. per acre	4	20.5 oz. per acre	30 days
	White mold (<i>Sclerotinia sclerotiorum</i>)	5.5 to 10 oz. per acre	2		
<p>Application Directions: For control of <i>Sclerotinia</i> white mold, begin applications of Endura® fungicide prior to row closure or at the onset of disease. Make a second application 14 days later if conditions continue to be favorable for disease development. For control of <i>Alternaria</i> early blight, begin applications of Endura prior to the onset of disease and continue applications at 7 to 14 day intervals if conditions continue to be favorable for disease development.</p> <p>Use the higher rates once disease has been confirmed in your area or weather conditions are conducive to disease development.</p> <p>Resistance Management: To limit the potential for development of resistance, do not make more than four (4) applications of Endura per season. Do not make more than two (2) sequential applications of Endura before alternating to a labeled fungicide with a different mode of action.</p>					

18/21

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Emerald, Endura, and Pristine are registered trademarks of BASF Corporation.

© 2004 BASF Corporation
All rights reserved

007969-00197.20040916.NVA 2004-04-201-0250.pdf
Supersedes NVA 2003-04-201-0217

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

 **BASF**

The Chemical Company

19/21

• **Endura**[®]

fungicide

GROUP 7 FUNGICIDE

For use in dry and succulent beans, bulb vegetables, canola, carrots, fruiting vegetables, grapes, lettuce, peanuts and potatoes

Active Ingredient

Boscalid, 3-pyridinecarboxamide, 2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl) 70.0%

Inert Ingredients 30.0%

Total 100.0%

[0.7 oz. a.i. (0.044 lb a.i.) in 1 oz. of **Endura**[®] Fungicide]

EPA Reg. No. 7969-197 EPA Est. No. 67545-AZ-01

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements: Hazards to Humans and Domestic Animals: Warning. Causes substantial but temporary eye injury. Harmful if absorbed through skin. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **HOT LINE NUMBER:** Have the product container or label with you when calling a poison con-

trol center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

Environmental Hazards: Do not apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

Storage and Disposal: Storage: Do not contaminate water, food, or feed by storage or disposal. **Pesticide disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

See the attached booklet for complete **First Aid, Precautionary Statements, Directions For Use, and Conditions of Sale and Warranty.**

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

NVA 2005-05-201-0317
2064922

Product of Brazil

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709



The Chemical Company

Net contents: 6.5 pounds

2006



The Chemical Company

Helping Make
Products Better™

February 14, 2006

U.S. Environmental Protection Agency
Office of Pesticide Programs (7505C)
Crystal Mall 2, Room 266A
1801 South Bell Street
Arlington, VA 22202

ATTN: Tony Kish – (PM 22)

Dear Mr. Kish:

SUBJECT: Endura® Fungicide: EPA Reg. No. 7969-197

Reference: Final Printed Labeling Notification

BASF is submitting 5 copies of the final printed labeling for Endura® fungicide (EPA Reg. No. 7969-197). This version of the labeling has the following minor changes:

1. Change of the BASF Logo from "BASF" to the new logo "BASF, The Chemical Company".
2. Change of the trademark from Endura™ fungicide to the registered trademark Endura® fungicide throughout the label
3. Change of the Establishment Number
4. Adding of emergency information on the front panel
4. Including the Front Panel as page 1 in the pagination
5. Change to the "Conditions of Sale and Warranty: Remove the statement "IN NO CASE SHALL BASF OR THE SELLER BE LIABLE..." and replace with "TO THE EXTENT PERMITTED BY LAW, BASF AND THE SELLER DISCLAIM..."
6. Minor editorial changes such as formatting, font size, headers, label numbers, etc.

The following are enclosed:

- (1) EPA Form 8570-1 "Application for Pesticide"
- (2) Five (5) copies of the final printed labeling for Endura® fungicide
- (3) Certification with Respect to Citation of Data
- (4) Copy of EPA Letter / Notice of Pesticide Registration dated July 23, 2003

2/21



The Chemical Company

If you have any questions or need further information, I can be reached by phone at 919-547-2976, or via e-mail at akkarik@basf.com.

Sincerely,
BASF Corporation
The Chemical Company

A handwritten signature in black ink that reads "K. H. Akkari".

Khalid H. Akkari, Ph.D.
NAFTA Registration Manager
Regulatory Affairs.

