

7969-197

12/17/2013

1154



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Ms. Christi Keating
BASF Corporation, Agricultural Products
26 Davis Drive
P.O. Box 13528
Research Triangle Park, NC 27709

DEC 17 2013

Subject: Endura Fungicide
EPA Reg. No. 7969-197
EPA Decision Number: 468290
Your labels submitted on July 27, 2012 and resubmitted on December 5, 2013

Dear Ms. Keating:

The master and supplemental labeling referred to above, submitted under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, to expand the use of: Vegetable, bulb group 3-07; Vegetable, fruiting, group 8-10; Fruit, pome, group 11-10; Caneberry subgroup 13-07A; Bushberry subgroup 13-07B; Fruit, small, vine climbing, except fuzzy kiwifruit, subgroup 13-07F; Berry, low growing, subgroup 13-07G; Oilseed, group 20; and Turnip greens is unconditionally accepted under section 3(c)(5).

Submit and/or cite all data required for reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data. Boscalid is scheduled to begin Registration Review in 2015. Please refer to the following link for more information: http://www.epa.gov/oppsrrd1/registration_review/fy12-fy15-schedule-summary.pdf

Stamped copies of the master and supplemental labels "Accepted" are enclosed for your records. These labels supersede all other previously accepted labels. Please submit one copy of the final printed labels before the product is released for shipment. If you have any questions please contact Heather Garvie by phone at: 703-308-0034 or via email at: garvie.heather@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Shaja B. Joyner".

Shaja B. Joyner
Product Manager (20)
Fungicide Branch (7504P)
Registration Division

Enclosure: Stamped labels "Accepted"

2154



The Chemical Company

Group 7 Fungicide

ACCEPTED
DEC 17 2013
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg No. 7969-197

Endwura®

fungicide

For use in the following crops: alfalfa; dry and succulent beans; berries; head and stem *Brassica*; leafy *Brassica* greens; bulb vegetables; celery; cucurbit vegetables; fruiting vegetables; grapes; leafy greens; leafy petioles; lettuce; low growing berry; oilseed crops; peanut; succulent and dried shelled peas; pome fruits; potato; rapeseed; root and tuber vegetables; small fruit, vine climbing; soybean; spinach; stone fruits; strawberries; sunflower; tree nuts; and turnip greens

Active Ingredient:

boscalid*: 3-pyridinecarboxamide, 2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl)-... 70.0%

Other Ingredients: 30.0%

Total: 100.0%

* 0.7 oz [0.044 lb ai] of boscalid in 1 oz of product

EPA Reg. No. 7969-197

EPA Est. No.

**KEEP OUT OF REACH OF CHILDREN
WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net Contents:

| FIRST AID | |
|--|---|
| If in eyes | <ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes. • Call a poison control center or doctor for treatment advice. |
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice. |
| If swallowed | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything to an unconscious person. |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice. |
| HOTLINE NUMBER | |
| <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).</p> | |

Precautionary Statements

Hazards to Humans and Domestic Animals

Warning. Causes substantial but temporary eye injury. Harmful if absorbed through skin. Harmful if swallowed. **DO NOT** get in eyes or on clothing. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Protective eyewear (goggles, face shield or safety glasses)
- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

DO NOT apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

Groundwater Advisory

Boscalid is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of boscalid from runoff water and sediment. Runoff of this product will be reduced

by avoiding applications when rainfall irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, precautions, and **Conditions of Sale and Warranty** are to be followed. This labeling must be in the user's possession during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Protective eyewear (goggles, face shield or safety glasses)
- Coveralls
- Chemical-resistant gloves
- Shoes plus socks

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 50 pounds) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity > 50 pounds) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

In Case of Emergency

In case of large-scale spillage regarding this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

Steps to be taken in case material is released or spilled:

- Dike and contain spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with water.
- Wash clothing before reuse.
- Keep spill out of all sewers and open bodies of water.

Product Information

This package contains **Endura® fungicide**, a water dispersible granule (WG). Boscalid, the active ingredient in **Endura**, belongs to the group of respiration inhibitors classified by the US EPA and Canada PMRA as carboxamides, or target site of action **Group 7** fungicides. **Endura** has a new mode of action and is effective against pathogens resistant to other fungicides.

Endura has a protective effect because it inhibits spore germination and a curative effect because it inhibits mycelial growth and sporulation of the fungus on the leaf surface. However, optimum disease control is achieved when **Endura** is applied in a regularly scheduled protective spray program and used in a rotation program with other fungicides.

Because of its high specific activity and rainfastness, **Endura** has good residual activity against target fungi.

Endura is not for use in greenhouse or transplant production systems.

Resistance Management

Endura contains boscalid, a **Group 7** fungicide, and is effective against pathogens resistant to fungicides with modes of action different from those of carboxamide (anilide) fungicides (target site **Group 7**), such as dicarboximides, sterol inhibitors, benzimidazoles, QoI fungicides, and phenylamides. Fungal isolates resistant to **Group 7** fungicides may eventually dominate the fungal population if **Group 7** fungicides are used predominantly and repeatedly in the same field in successive years as the primary method of control for the targeted pathogen species. This may result in reduction of disease control of **Endura** or other **Group 7** fungicides.

To maintain the performance of **Endura** in the field, **DO NOT** exceed the total number of sequential applications of **Endura** and the total number of applications of **Endura** per season stated in **Table 1. Endura® fungicide Crop-specific Restrictions and Limitations Overview** and **Table 2. Endura® fungicide Crop-specific Requirements**. Adhere to the label instructions regarding the consecutive use of **Endura** or other target site of action **Group 7** fungicides that have a similar site of action on the same pathogens.

The following instructions may delay the development of fungicide resistance:

1. **Tank mixtures** - Use tank mixtures with fungicides from different target site of action groups that are registered/permitted for the same use and that are effective against the pathogens of concern. Use at least the minimum labeled rates of each fungicide in the tank mix.
2. **IPM** - Integrate **Endura** into an overall disease and pest management program. Follow cultural practices known to reduce disease development. Consult your local extension specialist, certified crop advisor and/or BASF representative for additional IPM strategies established for your area. **Endura** may be used in Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.
3. **Monitoring** - Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development.
4. **Reporting** - If a **Group 7** target site fungicide, such as **Endura**, appears to be less or no longer effective against a pathogen that it previously controlled or suppressed, contact a BASF representative, local extension specialist, or certified crop advisor to assist in determining the cause of reduced performance.

Cleaning Spray Equipment

Clean spray equipment thoroughly before and after applying this product, particularly if a product with the potential to injure crops was used prior to **Endura**.

Application Instructions

Apply **Endura** rates as instructed by **Table 2. Endura® fungicide Crop-specific Requirements**. Apply **Endura** with ground sprayer, aerial equipment or through sprinkler irrigation equipment. Check equipment frequently for calibration. Under low-level disease conditions, use the minimum application rates; use maximum application rates and shortened spray schedules for severe or threatening disease conditions. **DO NOT** apply when conditions favor drift from target area.

Ground Application

Apply **Endura® fungicide** in sufficient water to ensure thorough coverage of foliage, bloom, and fruit. Thorough coverage is required for optimum disease control.

Directed or Banded Sprays

The rates on the **Endura** label reflect the amount of product that should be applied uniformly over an acre of ground on a broadcast basis.

In some crops, **Endura** may be applied as a directed or banded spray over the rows or plant beds, with the alleys or row middles left unsprayed. For such uses, the labeled **Endura** rates should be reduced in proportion to the area actually sprayed. This adjustment is necessary to avoid applying the product at use rates higher than permitted according to label directions.

Use the following formula to determine the broadcast equivalent rate for directed or banded sprays:

$$\begin{array}{r} \text{sprayed bed} \\ \text{width} \end{array} + \begin{array}{r} \text{unsprayed} \\ \text{row middles width} \end{array} = \begin{array}{r} \text{total row} \\ \text{width} \end{array}$$

$$\frac{\text{sprayed bed width in inches}}{\text{total row width in inches}} \times \frac{\text{broadcast rate}}{\text{treated acre}} = \frac{\text{band rate}}{\text{field acre}}$$

EXAMPLE: Directed spray application to 45-inch plant beds that are separated by 15-inches of unsprayed row middles.

$$\begin{array}{r} 45 \text{ inches} \\ \text{sprayed bed width} \end{array} + \begin{array}{r} 15 \text{ inches} \\ \text{unsprayed row middles} \end{array} = \begin{array}{r} 60 \text{ inches} \\ \text{total row width} \end{array}$$

The calculation to determine the appropriate equivalent rate of product to use for this situation based on a label broadcast rate of 6 ozs/A follows:

$$\frac{45 \text{ inches sprayed bed width}}{60 \text{ inches total row width}} \times \frac{6 \text{ ozs Endura}}{\text{treated acre}} = \frac{4.5 \text{ ozs Endura}}{\text{field acre}}$$

Aerial Application

DO NOT apply by air to the berries group. For all other crops listed on this label, aerial application can be made and thorough coverage is required to obtain optimum disease control. Avoid applications under conditions when uniform coverage cannot be obtained or when spray drift may occur. Use no less than 5 gallons of spray solution per acre. For aerial applications to tree and vine crops, use no less than 10 gallons of spray solution per acre. For all crops, thorough coverage is required for optimum disease control.

Directions For Use Through Sprinkler Irrigation Systems

Sprayer Preparation

Thoroughly clean chemical tank and injector system. Flush system with clean water.

Application Instructions

Apply **Endura** at rates and timings as described in this label.

Sprinkler Irrigation Application Use Precautions

- Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side [wheel] roll, traveler, big gun, solid set, or hand move irrigation systems. **DO NOT** apply this product through any other type of irrigation system.
- Add this product to the pesticide supply tank containing sufficient water to maintain a continuous flow by the injection equipment. In continuous moving systems, inject this product-water mixture continuously, applying the labeled rate per acre for that crop. **DO NOT** exceed 1/2 inch (13,577 gallons) per acre. In stationary or noncontinuous moving systems, inject the product-water mixture in the last 15 to 30 minutes of each set allowing sufficient time for all of the required pesticide to be applied by all the sprinkler heads and applying the labeled rate per acre for that crop. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. Thorough coverage of foliage is required for good control. Maintain good agitation during the entire application.
- If you have questions about calibration, contact State Extension Service specialist, equipment manufacturers or other experts.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

Additives and General Tank Mixing Information

Endura® fungicide can be tank mixed with most recommended fungicides and insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives as specified in **Table 2. Endura® fungicide Crop-specific Requirements.**

Under some conditions, the use of additives or adjuvants may improve the performance of **Endura**. However, all varieties and cultivars have not been tested with possible tank mix combinations. Local conditions can also influence crop tolerance and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing **Endura** with other products. Therefore, before using any tank mix (fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

When an adjuvant is to be used with this product, BASF recommends the use of a Chemical Producers and Distributors Association certified adjuvant.

Consult a BASF representative or local agricultural authorities for more information concerning additives.

Compatibility Test and Mixing Order

If tank mixtures are used, adhere to restrictions due to rates, label directions and precautions on all labels.

Compatibility Test for Tank Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of label rate per acre.

1. **Water** - For 100 gallons per acre spray volume, use 16 cups (1 gallon) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
2. **Water-dispersible products** (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions). Cap the jar and invert 10 cycles.
3. **Water-soluble products** - Cap the jar and invert 10 cycles.
4. **Emulsifiable concentrates** (oil concentrate or methylated seed oil when applicable). Cap the jar and invert 10 cycles.
5. **Water-soluble additives** - Cap the jar and invert 10 cycles.
6. Let the solution stand for 15 minutes.
7. **Evaluate** the solution for uniformity and stability. The spray solution should not have free oil on the surface, or fine particles that precipitate to the bottom, or thick (clabbered) texture. **DO NOT** use any spray solution that could clog spray nozzles.

Mixing Order

1. **Water** - Begin by agitating a thoroughly clean sprayer tank 3/4 full of clean water.
2. **Agitation** - Maintain constant agitation throughout mixing and application.
3. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
4. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. **Water-dispersible products** (such as **Endura[®] fungicide**, dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
6. **Water-soluble products**
7. **Emulsifiable concentrates** (such as oil concentrates when applicable)
8. **Water-soluble additives** (such as AMS or UAN when applicable)
9. **Remaining quantity water** - Make sure that each component is thoroughly mixed and suspended before adding tank mix partners. Make constant agitation during application.

Restrictions and Limitations - Alfalfa

- **DO NOT** use harvested seed for sprouting.
- Processed seed must be labeled "Not for human or animal consumption" at the processing plant. All alfalfa seed screenings must be disposed of in such a way that they cannot be distributed or used for food or feed.
- No portion of treated plant used for seed production may be grazed, used, or distributed for food or feed purposes.
- Livestock grazing or feeding is permitted for alfalfa for forage and hay only.

Restrictions and Limitations

- **DO NOT** exceed the maximum product rate (ozs/A) per year (season), the maximum rate per application, or the total number of applications of **Endura** per year (season) as stated in **Table 1. Endura[®] fungicide Crop-specific Restrictions and Limitations Overview** and **Table 2. Endura[®] fungicide Crop-specific Requirements**. Preharvest interval (PHI) restrictions are also included in these tables.
- **Livestock Feeding Restrictions - DO NOT** feed peanut hay to livestock.
- **DO NOT** apply more than the maximum annual use rate of ai/A or ozs of product/A for each specific crop from any combination of products containing boscalid (e.g. **Pristine[®] fungicide, Endura**).
- **Plantback Restrictions** - Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be planted after **14 days**.
- **DO NOT use on cowpea, field pea, grain lupin, sugar beet, garden beet, radish and turnip roots.**
- **Endura** is not for use in greenhouse or transplant production systems.

Table 1. Endura® fungicide Crop-specific Restrictions and Limitations Overview*

| Crop Group** | Minimum Time from Application to Harvest (PHI) (days) | Maximum Rate per Acre per Application (ozs product) | Maximum Number of Applications per Season**** | Maximum Rate per Acre per Season (ozs product) |
|---|--|--|--|---|
| Alfalfa for forage and hay | 14 | 6.5 | 3 | 19.5 |
| Alfalfa grown for seed | 14 | 11 | 2 | 22 |
| Beans, dried and succulent | 21 (dried) 7 (succulent) | 11 | 2 | 22 |
| Berry subgroups | 0 | 8 | 4 | 32 |
| <i>Brassica</i> , head and stem | 0 | 9 | 2 | 18 |
| <i>Brassica</i> , leafy greens | 14 | 9 | 2 | 18 |
| Bulb vegetable group | 7 | 6.8 | 6 | 41 |
| Celery | 0 | 9 | 2 | 18 |
| Cucurbit vegetables group | 0 | 6.5 | 4 | 26 |
| Fruiting vegetable group | 0 | 3.5 | 6 | 21 |
| Tomato*** | | 12.5 | 2 | 25 |
| Grapes*** | 14 | 4.5 | 5 | 24 |
| | | 8 | 3 | |
| Leafy greens (except <i>Brassica</i> , head lettuce, and leaf lettuce) | 14 | 9 | 2 | 18 |
| Leafy petioles (except <i>Brassica</i>) | 0 | 9 | 2 | 18 |
| Lettuce | 14 | 11 | 2 | 22 |
| Low growing berry subgroup (except cranberry and strawberry) | 0 | 8 | 5 | 40 |
| Oilseed crops (except rapeseed and sunflower) | 21 | 6 | 2 | 12 |
| Peanut | 14 | 10 | 3 | 30 |
| Peas, dried shelled | 21 | 11 | 2 | 22 |
| Peas, succulent | 7 | 11 | 2 | 22 |
| Pome fruits group | 0 | 6.5 | 4 | 26 |
| Potato*** | 10 | 4.5 | 4 | 20 |
| | | 10 | 2 | |
| Rapeseed | 21 | 6 | 2 | 12 |
| Root and tuber vegetables: root vegetables subgroup 1A*** | 0 | 4.5 | 5 | 23.4 |
| | | 7.8 | 3 | |
| Root and tuber vegetables: tuberous and corm vegetables subgroup 1C*** | 10 | 4.5 | 4 | 20 |
| | | 10 | 2 | |
| Small fruit, vine climbing subgroup (except fuzzy kiwifruit and grapes) | 14 | 8 | 5 | 24 |
| Soybean | 21 | 11 | 2 | 22 |
| Spinach | 0 | 9 | 2 | 18 |
| Stone fruits group | 0 | 5.3 | 5 | 26.5 |
| Strawberries | 0 | 8 | 5 | 40 |
| Sunflower | 21 | 9 | 2 | 18 |

(continued)

Table 1. Endura® fungicide Crop-specific Restrictions and Limitations Overview (continued)

| Crop Group** | Minimum Time from Application to Harvest (PHI) (days) | Maximum Rate per Acre per Application (ozs product) | Maximum Number of Applications per Season**** | Maximum Rate per Acre per Season (ozs product) |
|-----------------|---|---|---|--|
| Tree nuts group | 14 (for almond - 25) | 5.3 | 4 | 21.2 |
| Turnip greens | 14 | 9 | 2 | 18 |

* See **Table 2. Endura® fungicide Crop-specific Requirements** for complete directions and exceptions.
 ** For a complete list of crops within a crop group, see **Table 2. Endura® fungicide Crop-specific Requirements**.
 *** Maximum rate per acre, number of applications per season, and maximum rate per acre per season vary for tomato, grapes, potato, and root and tuber vegetables depending on the target disease. Refer to **Crop-specific Use Directions** for fruiting vegetables, grapes, potato, and root and tuber vegetables for maximum rates and number of applications by target disease.
 **** At the maximum use rate only, except for tomato, grapes, potato, and root and tuber vegetables.

Crop-specific Use Directions

Table 2. Endura® fungicide Crop-specific Requirements

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|-----------------------------|---|---|---|---|---|
| Alfalfa for forage and hay* | Common leaf spot (<i>Pseudopezizza medicaginis</i>) Leaf spot (<i>Leptosphaerulina briosiani</i>) Spring black stem and leaf spot (<i>Phoma medicaginis</i>) Suppression Only: Powdery mildew (<i>Erysiphe pisi</i>) Southern blight (<i>Sclerotium rolfsii</i>) White mold (<i>Sclerotinia sclerotiorum</i> , <i>S. trifoliorum</i>) | 6.5 | 2 per cutting and 3 total per growing season | 19.5 | 14 |
| | Alfalfa grown for seed* | Common leaf spot (<i>Pseudopezizza medicaginis</i>) Leaf spot (<i>Leptosphaerulina briosiani</i>) Spring black stem and leaf spot (<i>Phoma medicaginis</i>) Suppression Only: Powdery mildew (<i>Erysiphe pisi</i>) Southern blight (<i>Sclerotium rolfsii</i>) White mold (<i>Sclerotinia sclerotiorum</i> , <i>S. trifoliorum</i>) | 6 to 11 | 2 | 22 |
| | | 8 to 11 | | | |

(continued)

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

Alfalfa, continued

Application Directions. Begin **Endura** applications when conditions favorable for disease are expected, but prior to onset of disease development. For stand establishment of fall seeded alfalfa, begin applications in fall through early winter prior to first snowfall or extended cool, wet conditions. For seed pod protection, begin applications at 10% to 30% bloom.

Disease control can be improved when application equipment and spray volumes are adjusted to achieve thorough canopy penetration and coverage.

Alfalfa for forage and hay. Repeat application on a 14- to 21-day interval if conditions are conducive for disease development. Use the shorter interval when disease pressure is high.

No restriction for livestock grazing or feeding for alfalfa for forage and hay.

Alfalfa grown for seed. Repeat application on a 7- to 14-day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

Under some conditions, the use of additives or adjuvants may improve the performance of **Endura**.

DO NOT use harvested seed for sprouting.

Processed seed must be labeled "Not for human or animal consumption" at the processing plant. All alfalfa seed screenings must be disposed of in such a way that they cannot be distributed or used for food or feed.

No portion of treated plant used for seed production may be grazed, used, or distributed for food or feed purposes.

Resistance Management. To limit the potential for development of resistance in **alfalfa for forage and hay**, **DO NOT** make more than three (3) applications of **Endura** per season. To limit the potential for development of resistance in **alfalfa grown for seed**, **DO NOT** make more than two (2) applications of **Endura** per season.

* Not registered for use in California.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|---|--|---|---|---|
| <p>Beans, dried and succulent</p> <p>Broad bean Chickpea (garbanzo bean) Guar Jack bean Lablab bean</p> <p>Lupinus spp. Sweet lupin White lupin White sweet lupin</p> <p>Phaseolus spp. Field bean Kidney bean Lima bean Navy bean Pink bean Pinto bean Runner bean Snap bean Tepary bean Wax bean</p> <p>Vigna spp. Adzuki bean Asparagus bean Blackeyed pea Catjang Chinese longbean Crowder pea Moth bean Mung bean Rice bean Southern pea Urd bean Yardlong bean</p> | <p>Ascochyta blight (<i>Phoma exigua</i>, <i>Ascochyta</i> spp.)</p> | <p>6</p> | <p>2</p> | <p>22</p> | <p>21 (Dried Beans)</p> |
| | <p>Botrytis gray mold (<i>Botrytis cinerea</i>)</p> <p>White mold (<i>Sclerotinia sclerotiorum</i>)</p> | <p>8 to 11</p> | | | <p>7 (Succulent Beans)</p> |

Application Directions. Apply **Endura** at the beginning of flowering or prior to onset of disease. Use the higher rate for extended protection and maximum yield benefit. Apply a second time at full bloom if conditions are favorable for disease development or if heavy disease has already set in.

Ascochyta blight in chickpeas develops quickly once established, so early detection and application is essential to reduce losses.

Apply at the beginning of flowering. Make a second application 7 to 10 days later if disease persists or weather conditions are favorable for disease development.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

Restrictions. DO NOT use on cowpea, field pea, and grain lupin. DO NOT feed treated pea commodities to livestock.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|---|--|---|---|---|
| <p>Berry subgroups</p> <p>Bushberry subgroup</p> <p>Aronia berry</p> <p>Black currant</p> <p>Blueberry (highbush and lowbush)</p> <p>Buffalo currant</p> <p>Chilean guava</p> <p>Currant</p> <p>Elderberry</p> <p>European barberry</p> <p>Gooseberry</p> <p>Highbush cranberry</p> <p>Honeysuckle, edible</p> <p>Huckleberry</p> <p>Jostaberry</p> <p>Juneberry</p> <p>Lingonberry</p> <p>Native currant</p> <p>Red currant</p> <p>Salal</p> <p>Sea buckthorn</p> <p>Caneberry subgroup</p> <p>Blackberry (all varieties)</p> <p>Loganberry</p> <p>Raspberry (black and red)</p> <p>Wild raspberry</p> | <p>Botrytis gray mold (<i>Botrytis cinerea</i>)</p> | <p>8</p> | <p>4</p> | <p>32</p> | <p>0</p> |

Application Directions. Begin applications of **Endura** prior to disease development and continue on a 7- to 14-day interval.

DO NOT apply by air.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Endura** per season.

DO NOT make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|--|--|---|---|---|
| Brassica leafy vegetables Head and stem Brassica Broccoli Brussels sprouts Chinese broccoli (gai lan) Cabbage Chinese cabbage (napa) Chinese mustard cabbage (gai choy) Cauliflower Cavalo broccolo Kohlrabi | Alternaria blight (Black spot) (<i>Alternaria</i> spp.) Gray mold (<i>Botrytis cinerea</i>) Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i> , <i>S. minor</i>) Suppression Only: Powdery mildew (<i>Erysiphe polygoni</i>) Rhizoctonia bottom rot (<i>Rhizoctonia solani</i>) | 6 to 9 | 2 | 18 | 0 |
| <p>Application Directions. Begin applications of Endura prior to disease development and continue on a 7- to 14-day interval.</p> <p>Use the higher rate and the shorter interval when disease pressure is high.</p> <p>Resistance Management. To limit the potential for development of resistance, DO NOT make more than two (2) applications of Endura per season.</p> | | | | | |

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|--|--|---|---|---|
| Brassica leafy vegetables Leafy Brassica greens Broccoli raab (rapini) Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens | Alternaria blight (Black spot) (<i>Alternaria</i> spp.) Gray mold (<i>Botrytis cinerea</i>) Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i> , <i>S. minor</i>) Suppression Only: Powdery mildew (<i>Erysiphe polygoni</i>) Rhizoctonia bottom rot (<i>Rhizoctonia solani</i>) | 6 to 9 | 2 | 18 | 14 |
| <p>Application Directions. Begin applications of Endura prior to disease development and continue on a 7- to 14-day interval.</p> <p>Use the higher rate and the shorter interval when disease pressure is high.</p> <p>Resistance Management. To limit the potential for development of resistance, DO NOT make more than two (2) applications of Endura per season.</p> | | | | | |

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|---|--|---|---|---|
| Bulb vegetable group Chive, fresh leaves Chive, Chinese, fresh leaves Daylily, bulb Elegans hosta Fritillaria, bulb Fritillaria, leaves Garlic, bulb Garlic, great-headed, bulb Garlic, Serpent, bulb Kurrat Lady's leek Leek Leek, wild Lily, bulb Onion, Beltsville bunching Onion, bulb Onion, Chinese, bulb Onion, fresh Onion, green Onion, macrostem Onion, pearl Onion, potato, bulb Onion, tree, tops Onion, Welsh, tops Shallot, bulb Shallot, fresh leaves Cultivars, varieties and/or hybrids of these | Botrytis leaf blight (<i>Botrytis</i> spp.) Purple blotch (<i>Alternaria porri</i>) | 6.8 | 6 | 41 (In-furrow plus foliar) | 7 |

Application Directions. For control of purple blotch and Botrytis leaf blight, begin applications of **Endura** prior to disease development and continue on a 7- to 14-day interval.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than six (6) applications of **Endura** per season.

DO NOT make more than two (2) applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

Instructions for In-furrow Use to Aid in the Control of Soilborne White Rot (*Sclerotium cepivorum*) in the Bulb Vegetables Group, including Garlic and Onions

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|--|--|---|---|---|
| Bulb vegetable group Chive, fresh leaves Chive, Chinese, fresh leaves Daylily, bulb Elegans hosta Fritillaria, bulb Fritillaria, leaves Garlic, bulb Garlic, great-headed, bulb Garlic, Serpent, bulb Kurrat Lady's leek Leek Leek, wild Lily, bulb Onion, Beltsville bunching Onion, bulb Onion, Chinese, bulb Onion, fresh Onion, green Onion, macrostem Onion, pearl Onion, potato, bulb Onion, tree, tops Onion, Welsh, tops Shallot, bulb Shallot, fresh leaves Cultivars, varieties and/or hybrids of these | White rot (<i>Sclerotium cepivorum</i>) | 6.8 | 6 | 41 (In-furrow plus foliar) | 7 |

Application Directions. Apply **Endura** at planting as an in-furrow spray by directing spray pattern to the soil bed. Use a 4- to 6-inch band spray pattern applied directly over/into the seed furrow before covering with soil. Depending upon the level of potential infection, make additional foliar applications at the 6.8 ozs/A rate. Use a minimum volume of application of 5 gallons of water per acre.

Apply no more than a total of 41 ozs/A per season for a combination of the in-furrow and foliar uses.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than six (6) applications of **Endura** per season.

DO NOT make more than two (2) applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|---|--|---|---|---|
| Celery Celery (Chinese) | Alternaria leaf spot <i>(Alternaria spp.)</i> Suppression Only: Powdery mildew <i>(Erysiphe spp.)</i> | 4.5 to 9 | 2 | 18 | 0 |
| | Botrytis rot <i>(Botrytis spp.)</i> Phoma <i>(Phoma spp.)</i> Sclerotinia rot and blight <i>(Sclerotinia spp.)</i> Pink rot <i>(Sclerotinia sclerotiorum)</i> Suppression Only: Crater rot <i>(Rhizoctonia solani)</i> | 8 to 9 | | | |

Application Directions. Begin applications of **Endura** prior to the onset of disease development and continue on a 7-day interval. For pink rot and crater rot, make the first application just prior to row closure with a subsequent application two weeks later.

Use the higher rate when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|---|--|---|---|---|
| <p>Cucurbit vegetables group</p> <p>Includes all types and hybrids of:</p> <ul style="list-style-type: none"> Chayote Chinese waxgourd Citron melon Cucumber Gherkin Pumpkin Watermelon <p>Edible gourd</p> <ul style="list-style-type: none"> Chinese okra Cucuzza Hytan <p>Momordica spp.</p> <ul style="list-style-type: none"> Balsam apple Balsam pear Bitter melon Chinese cucumber <p>Muskmelon</p> <ul style="list-style-type: none"> Cantaloupe Casaba Crenshaw melon Golden pershaw melon Honeydew melon Honey balls Mango melon Persian melon Pineapple melon Santa Claus melon Snake melon <p>Summer squash</p> <ul style="list-style-type: none"> Crookneck squash Scallop squash Straightneck squash Vegetable marrow Zucchini <p>Winter squash</p> <ul style="list-style-type: none"> Acorn squash Butternut squash Calabaza Hubbard squash Spaghetti squash | <p>Alternaria blight (<i>Alternaria cucumerina</i>)</p> <p>Gummy stem blight (<i>Didymella bryoniae</i>)</p> <p>Suppression Only:</p> <p>Powdery mildew (<i>Sphaerotheca fuliginea</i>, <i>Erysiphe cichoracearum</i>)</p> | <p>6.5</p> | <p>4</p> | <p>26</p> | <p>0</p> |

Application Directions. Begin applications of **Endura** prior to disease development and continue on a 7- to 14-day interval. Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Endura** per season.

DO NOT make more than one (1) application of **Endura** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|---|--|---|---|---|
| Fruiting vegetable group African eggplant Bush tomato Bell pepper Cocona Currant tomato Eggplant Garden huckleberry Goji berry Groundcherry Martynia Naranjilla Okra Pea eggplant Pepino Pepper (all varieties) Nonbell pepper Roselle Scarlet eggplant Sunberry Tomatillo Tree tomato Cultivars, varieties and/or hybrids of these | Early Blight <i>(Alternaria solani)</i> | 2.5 to 3.5 or 2.5 to 3.5 ozs per 100 gallons of spray volume (dilute)* | 6 | 21 | 0 |
| Target spot <i>(Corynespora cassiicola)</i> Suppression Only: Botrytis gray mold** <i>(Botrytis cinerea)</i> | 3.5 | | | | |
| Tomato | Black mold <i>(Alternaria alternata)</i> | 3.5 to 5 | 5 | 25 | |
| | Botrytis gray mold <i>(Botrytis cinerea)</i> | 9 to 12.5 | 2 | | |

Application Directions. Begin applications of **Endura** prior to disease development and continue on a 7- to 14-day interval for early blight, Botrytis gray mold, and black mold.

Use the higher rate and the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than six (6) applications of **Endura** at the low rate or two (2) applications of **Endura** at the high rate per season. **DO NOT** make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

* For applications based on dilute volume, plants should be sprayed to runoff. Apply a minimum of 20 gallons of spray volume per acre, and increase the spray volume as the plants grow during the season. Spray volume should be proportional to the amount of plant tissue to such that 100 gallons of spray per acre is used on mature plants.

** Not registered for use in California.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--------|---|--|---|---|---|
| Grapes | Powdery mildew (<i>Uncinula necator</i>) | 4.5 | 5 | 24 | 14 |
| | Botrytis gray mold (<i>Botrytis cinerea</i>) | 8 | 3 | | |

Application Directions. For powdery mildew control, begin applications of **Endura** at budbreak or prior to the onset of disease and continue on a 10- to 14-day interval.

For the control of Botrytis gray mold, begin applications of **Endura** prior to disease development and when conditions favor disease development during early bloom, bunch pre-closure or veraison.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Endura** at the low rate or three (3) applications of **Endura** at the high rate per season.

DO NOT make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|--|--|---|---|---|
| Leafy greens* (except Brassica, head lettuce, and leaf lettuce) Amaranth Arugula Chervil Chrysanthemum (edible-leaved and garland) | Alternaria leaf spot (<i>Alternaria</i> spp.) Ascochyta leaf spot (<i>Ascochyta</i> spp.) Phoma (<i>Phoma</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.) | 4.5 to 9 | 2 | 18 | 14 |
| Corn salad Cress (garden and upland) Dandelion Dock Endive Orach Parsley Purslane (garden and winter) Radicchio (red chicory) | Botrytis rot (<i>Botrytis</i> spp.) Sclerotinia rot and blight (<i>Sclerotinia</i> spp.) | 7 to 9 | | | |

Application Directions. Begin applications of **Endura** prior to the onset of disease development and continue on a 7-day interval.

Use the higher application rate when disease pressure is high.

No restriction on livestock grazing or feeding.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

* Not registered for use in California except on radicchio for control of Sclerotinia rot and blight when applied at a rate of 9 ozs/A.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|--|--|---|---|---|
| Leafy petioles* (except Brassica) Cardoon Celtuce Fennel (Florence) Rhubarb Swiss chard | Alternaria leaf spot (<i>Alternaria</i> spp.) Ascochyta leaf spot (<i>Ascochyta</i> spp.) Phoma (<i>Phoma</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.) | 4.5 to 9 | 2 | 18 | 0 |
| | Botrytis rot (<i>Botrytis</i> spp.) Sclerotinia rot and blight (<i>Sclerotinia</i> spp.) | 7 to 9 | | | |

Application Directions. Begin applications of **Endura** prior to the onset of disease development and continue on a 7-day interval.

Use the higher application rate when disease pressure is high.

No restriction on livestock grazing or feeding.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

* Not registered for use in California.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---------------------------------|---|--|---|---|---|
| Lettuce Head and Leaf | Lettuce drop (<i>Sclerotinia minor</i> , <i>S. sclerotiorum</i>) Botrytis rot (<i>Botrytis cinerea</i>) Phoma basal rot (<i>Phoma exigua</i>) Suppression Only: Rhizoctonia bottom rot (<i>Rhizoctonia solani</i>) Lettuce powdery mildew (<i>Erysiphe cichoracearum</i>) | 8 to 11 | 2 | 22 | 14 |

Application Directions. A protective fungicide barrier is needed to maximize disease control. Use the higher rate when disease pressure is high.

At Planting or Preemergence Applications for Sclerotinia Control

For chemigation applications, apply as outlined in the **Directions For Use Through Sprinkler Irrigation Systems** section in this label. Light incorporation by cultivation can increase the performance.

Post-emergence Applications for Control of Sclerotinia and other diseases

Apply **Endura** on direct-seeded lettuce immediately after emergence or prior to disease development. On transplanted lettuce, make the first application immediately after transplanting or prior to the onset of disease.

Make a second application if the soil surface is disturbed by cultivation or thinning and if conditions continue to favor disease development.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|---|--|---|---|---|
| Low growing berry subgroup (except cranberry and strawberry) Bearberry Bilberry Cloudberry Muntries Partridgeberry | Botrytis gray mold <i>(Botrytis cinerea)</i> | 8 | 5 | 40 | 0 |

Application Directions. Begin applications of **Endura** no later than 10% bloom or prior to the onset of disease and continue on a 7- to 14-day interval.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Endura** per season.

DO NOT make more than three (3) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|---|--|---|---|---|
| Oilseed crops (except rapeseed and sunflower) Flax seed | PasmO (<i>Septoria linicola</i>) | 5 to 6 | 2 | 12 | 21 |
| Borage Calendula Castor oil plant Chinese tallowtree Crambe Cuphea Echium Euphorbia Evening primrose Gold of pleasure Hare's ear mustard Jojoba Lesquerella Lunaria Meadowfoam Milkweed Mustard seed Niger seed Oil radish Poppy seed Rose hip Safflower Sesame Stokes aster Sweet rocket Tallowwood Tea oil plant Vernonia | Sclerotinia rot and blight (<i>Sclerotinia</i> spp.) Suppression Only: Rust (<i>Puccinia helianthi</i>) | | | | |

Application Directions. For optimal disease control, begin applications of **Endura** prior to disease development and continue on a 7- to 14-day interval if conditions are conducive for disease development.

Use the higher rate and shorter interval when disease pressure is high.

No restriction on livestock grazing or feeding.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--------|---|--|---|---|---|
| Peanut | Early leaf spot (<i>Cercospora arachidicola</i>) Late leaf spot (<i>Cercosporidium personatum</i>) | 6.5 to 10 | 3 | 30 | 14 |
| | Sclerotinia blight (<i>Sclerotinia minor</i>) Web blotch (<i>Phoma arachidicola</i>) Suppression Only: Sclerotium stem rot, Southern stem rot (<i>Sclerotium rolfsii</i>) | 8 to 10 | | | |

Application Directions. For control of early and late leaf spot and web blotch, begin applications of **Endura** prior to the onset of disease and continue on a 14-day interval.

For control of Sclerotinia blight, begin applications of **Endura** prior to the onset of disease or at 45 to 60 days after planting. Make a second application 14 to 21 days later.

For suppression of Southern stem rot (*Sclerotium rolfsii*), apply **Endura** prior to disease development or at 45 to 60 days after planting. Two additional applications may be made at 14-day intervals. For improved control of Southern stem rot, **Endura** may be mixed with other labeled, effective fungicides.

Use the higher rate and/or shorter spray interval when disease pressure is high or in fields with a history of disease.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than three (3) applications of **Endura** per season.

DO NOT make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Restrictions. DO NOT feed peanut hay to livestock. DO NOT graze or harvest for forage use.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days). |
|---|---|--|---|---|--|
| Peas, dried shelled Lentil (<i>Lens</i>) Pea (<i>Pisum</i>) Garden pea Green pea Pigeon pea | Alternaria leaf and pod spot (<i>Alternaria</i> spp.) | 6 | 2 | 22 | 21 |
| | Ascochyta blight (<i>Phoma exigua</i> , <i>Ascochyta</i> spp.) Botrytis gray mold (<i>Botrytis cinerea</i>) Mycosphaerella blight (<i>Mycosphaerella</i> spp.) White mold (<i>Sclerotinia sclerotiorum</i>) Suppression Only: Powdery mildew (<i>Erysiphe polygoni</i>) | | | | |

Application Directions. For optimal disease control, begin applications of **Endura** prior to disease development or at the beginning of flowering and repeat on a 5- to 14-day interval if conditions are conducive for disease development.

Use the higher rate and shorter interval when disease pressure is high.

Ascochyta blight in lentils develops quickly once established, so early detection and application is essential to reduce losses.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

Restrictions. DO NOT use on cowpea, field pea, and grain lupin. DO NOT feed treated pea commodities to livestock.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|--|--|---|---|---|
| Pome fruits group Apple Azarole Crabapple Loquat Mayhaw Medlar Pear Pear, Asian Pear, Oriental Quince Quince, Chinese Quince, Japanese Tejocote Cultivars, varieties and/or hybrids of these | Apple scab (<i>Venturia inaequalis</i>) Pear scab (<i>Venturia pirina</i>) Powdery mildew (<i>Podosphaera leucotricha</i>) Fly speck (<i>Zygothiala jamaicensis</i>) Alternaria blotch (<i>Alternaria mali</i>) | 6.5 | 4 | 26 | 0 |

Application Directions for scab and powdery mildew. Begin applications of **Endura** prior to disease development and continue on a 7- to 10-day interval.

Use the shorter interval when disease pressure is high.

Application Directions for fly speck and Alternaria blotch. Begin applications of **Endura** prior to disease development and continue on a 7- to 14-day interval. Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Endura** per season.

DO NOT make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--------|---|--|---|---|---|
| Potato | Brown Spot (<i>Alternaria alternata</i>) Early blight (<i>Alternaria</i> spp.) | 3.5 to 4.5 | 4 | 20 | 10 |
| | White mold (<i>Sclerotinia sclerotiorum</i>) Suppression Only: Gray mold (<i>Botrytis cinerea</i>) | 5.5 to 10 | 2 | | |

Application Directions. For control of *Sclerotinia* white mold, begin applications of **Endura** prior to infection. Generally, the first infections begin when the rows start to close and/or when the plants start flowering. Make a second application 14 days later if conditions continue to be favorable for disease development.

For control of *Alternaria* early blight or suppression of *Botrytis* gray mold, begin applications of **Endura** prior to the onset of disease and continue applications at 7- to 14-day intervals if conditions continue to be favorable for disease development.

Use the higher rates when disease has been confirmed in your area or weather conditions are conducive to disease development.

The use of additives or adjuvants may improve the performance of **Endura**. For additional details and precautions, refer to **Additives and General Tank Mixing Information**.

No restriction on livestock grazing or feeding.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Endura** at the low rate or two (2) applications of **Endura** at the high rate per season.

DO NOT make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|--|--|---|---|---|
| Rapeseed (cultivars, varieties and/or hybrids, including canola and crambe) | White mold/Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>) | 5 to 6 | 2 | 12 | 21 |

Application Directions. Begin applications of **Endura** at 20% to 50% flowering or prior to the onset of disease. Use the higher rate for extended protection. Apply a second time if conditions continue to be favorable for disease development.

No restriction on livestock grazing or feeding.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|--|--|---|---|---|
| Root and tuber vegetables Root vegetables subgroup 1A Carrot Celeriac Ginseng Horseradish Skirret | Alternaria leaf spot (<i>Alternaria dauci</i> , <i>Alternaria</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.) | 4.5 | 5 | 23.4 | 0 |
| | Cottony rot, White rot of rhizomes, Watery soft rot (<i>Sclerotinia sclerotiorum</i>) Gray mold rot (<i>Botrytis cinerea</i>) | 7.8 | 3 | | |

Application Directions. For optimal disease control, begin applications of **Endura** prior to disease development and repeat on a 7- to 14-day interval if conditions are conducive for disease development.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Endura** per season. **DO NOT** make more than two (2) applications of **Endura** before alternating to a labeled fungicide with a different mode of action for at least one application.

Restrictions. **DO NOT** use on garden beet, sugar beet, radish or turnip roots.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|---|--|---|---|---|
| Root and tuber vegetables | Early blight (<i>Alternaria solani</i>) | 2.5 to 4.5 | 4 | 20 | 10 |
| Tuberous and corm vegetables subgroup 1C Arrowroot Chinese artichoke Jerusalem artichoke Edible canna Chayote (root) Ginger Leren Sweet potato Turmeric Yam bean True yam | Sclerotinia white mold (<i>Sclerotinia sclerotiorum</i>) | 5.5 to 10 | 2 | | |

Application Directions. For optimal disease control, begin applications of **Endura** prior to disease development and repeat on a 7- to 14-day interval if conditions are conducive for disease development.

Use the higher rate and shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Endura** at the low rate or two (2) applications of **Endura** at the high rate per season. **DO NOT** make more than two (2) applications of **Endura** before alternating to a labeled fungicide with a different mode of action for at least one application.

Restrictions. DO NOT use on garden beet, sugar beet, radish or turnip roots.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|--|--|---|---|---|
| Small fruit, vine climbing subgroup (except fuzzy kiwifruit and grape) | Powdery mildew (<i>Uncinula necator</i>) | 4.5 | 5 | 24 | 14 |
| Amur river grape Gooseberry Kiwifruit, hardy Maypop Schisandra berry | Botrytis gray mold (<i>Botrytis cinerea</i>) | 8 | 3 | | |

Application Directions. For powdery mildew control, begin applications of **Endura** at budbreak or prior to the onset of disease and continue on a 10- to 14-day interval.

For the control of Botrytis gray mold, begin applications of **Endura** prior to disease development and when conditions favor disease development during early bloom, bunch pre-closure or veraison.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Endura** at the low rate or three (3) applications of **Endura** at the high rate per season.

DO NOT make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|----------|--|--|---|---|---|
| Soybean* | Alternaria leaf spot (<i>Alternaria</i> spp.) | 3.5 to 5.5 | 2 | 22 | 21 |
| | Suppression Only: Brown spot (<i>Septoria glycines</i>) Cercospora blight and leaf spot (<i>Cercospora kikuchii</i>) Frogeye leaf spot (<i>Cercospora sojina</i>) Rhizoctonia aerial blight (<i>Rhizoctonia solani</i>) | 5.5 | | | |
| | Suppression Only: White mold (<i>Sclerotinia sclerotiorum</i>) | 5.5 to 11 | | | |

Application Directions. Begin applications of **Endura** prior to disease development or when conditions are conducive for disease development and continue on a 7- to 14-day interval.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

* Not registered for use in California.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|--|--|---|---|---|
| Spinach Spinach (New Zealand and vine) | Alternaria leaf spot (<i>Alternaria</i> spp.) Ascochyta leaf spot (<i>Ascochyta</i> spp.) Suppression Only: Powdery mildew (<i>Erysiphe</i> spp., <i>Phyllactinia</i> spp., <i>Sphaerotheca</i> spp.) | 4.5 to 9 | 2 | 18 | 0 |
| | Botrytis rot (<i>Botrytis</i> spp.) Phoma (<i>Phoma</i> spp.) Sclerotinia rot and blight (<i>Sclerotinia</i> spp.) | 8 to 9 | | | |
| <p>Application Directions. Begin applications of Endura prior to the onset of disease development and continue on a 7-day interval.</p> <p>Resistance Management. To limit the potential for development of resistance, DO NOT make more than two (2) applications of Endura per season.</p> | | | | | |

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|--|--|---|---|---|
| Stone fruits group Apricot Cherry (sweet and tart) Nectarine Peach Plum (all varieties) Plumcot Prune | Alternaria leaf spot (<i>Alternaria</i> spp.) Brown rot (<i>Monilinia</i> spp.) Monilinia blossom blight (<i>Monilinia</i> spp.) Ripe fruit rot (<i>Botrytis cinerea</i> , <i>Monilinia fructicola</i> , <i>Monilinia laxa</i>) Scab (<i>Cladosporium carpophilum</i>) Suppression Only: Powdery mildew (<i>Sphaerotheca</i> spp., <i>Podosphaera</i> spp.) | 5.3 | 5 | 26.5 | 0 |

Application Directions. Begin applications of **Endura** at pink bud or prior to the onset of disease and continue on a 7- to 14-day interval.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Endura** per season.

DO NOT make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

For aerial application to stone fruits trees, use no less than 10 gallons of spray solution per acre.

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---------------------|--|--|---|---|---|
| Strawberries | Botrytis gray mold (<i>Botrytis cinerea</i>) | 8 | 5 | 40 | 0 |

Application Directions. Begin applications of **Endura** no later than 10% bloom or prior to the onset of disease and continue on a 7- to 14-day interval.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Endura** per season.

DO NOT make more than three (3) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|------------|--|--|---|---|---|
| Sunflower* | Alternaria leaf spot <i>(Alternaria spp.)</i> Botrytis gray mold <i>(Botrytis cinerea)</i> Powdery mildew <i>(Erysiphe cichoracearum)</i> | 4.5 to 9 | 2 | 18 | 21 |
| | <u>Suppression Only:</u> Sclerotinia head rot <i>(Sclerotinia sclerotiorum)</i> | 6 to 9 | | | |

Application Directions. For optimal disease control, begin applications of **Endura** prior to disease development and continue on a 7- to 14-day interval if conditions are conducive for disease development.

Use the higher rate and shorter interval when disease pressure is high.

No restriction on livestock grazing or feeding.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

* Not registered for use in California.

Table 2. Endura® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|---|--|---|---|---|
| Tree nuts group Almond Beech nut Brazil nut Butternut Cashew Chestnut Chinquapin Filbert Hickory nut Macadamia nut Pecan Pistachio Walnut (black and English) | Alternaria late blight (<i>Alternaria alternata</i>) Alternaria leaf spot (<i>Alternaria</i> spp.) Blossom blight (<i>Monilinia</i> spp.) Green fruit rot (<i>Botrytis cinerea</i>) Suppression Only: Scab (<i>Cladosporium carpophilum</i> , <i>C. caryigenum</i>) | 5.3 | 4 | 21.2 | 14 (for almond - 25) |

Application Directions. In almond, begin applications of **Endura** at pink bud and continue on a 7- to 14-day interval up to 5 weeks after petal fall. In pecan, begin application of **Endura** prior to disease development and continue on a 7- to 21-day interval for the suppression of scab. For all other crops listed above, apply **Endura** prior to the onset of disease and continue on a 7- to 28-day interval.

In all cases, use the shorter interval when disease pressure is high or shoot growth is very rapid.

No restriction on livestock grazing or feeding for almond hulls.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Endura** per season.

DO NOT make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

For aerial application to tree nuts, use no less than 10 gallons of spray solution per acre.

Table 2. **Endura**® fungicide Crop-Specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|----------------------|--|--|---|---|---|
| Turnip greens | Alternaria blight (Black spot) <i>(Alternaria spp.)</i> Gray mold <i>(Botrytis cinerea)</i> Sclerotinia stem rot <i>(Sclerotinia sclerotiorum, S. minor)</i> Suppression Only: Powdery mildew <i>(Erysiphe polygoni)</i> Rhizoctonia bottom rot <i>(Rhizoctonia solani)</i> | 6 to 9 | 2 | 18 | 14 |

Application Directions. Begin applications of **Endura** prior to disease development and continue on a 7- to 14-day interval.

Use the higher rate and the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Endura and Pristine are registered trademarks of BASF.

© 2013 BASF Corporation
All rights reserved.

007969-00197.20131105b.NVA 2013-04-201-0144

Supersedes: NVA 2013-04-201-0070

Supplemental: NVA 2012-04-201-0130

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



Group 7 Fungicide

Supplemental Label



For use in the following crops: berries; bulb vegetables; fruiting vegetables; low growing berry; oilseed crops; pome fruits; rapeseed; small fruit, vine climbing; sunflower; and turnip greens

This supplemental label expires November 30, 2016 and must not be used or distributed after this date.

Active Ingredient:

boscalid*: 3-pyridinecarboxamide,2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl)- 70.0%

Other Ingredients: 30.0%

Total: 100.0%

*0.7 oz [0.044 lb ai] boscalid in 1 oz of product

EPA Reg. No. 7969-197

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Endura® fungicide** container label, EPA Reg. No. 7969-197, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Endura** before applying.
- Use of **Endura** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Endura**.

Application Instructions

Refer to the **Endura** container label for additional application instructions and restrictions.

ACCEPTED
DEC 17 2013
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg No. **7969-197**

Table 1. Endura® fungicide Crop-specific Restrictions and Limitations Overview*

| Crop** | Minimum Time from Application to Harvest (PHI) (days) | Maximum Rate per Acre per Application (ozs product) | Maximum Number of Applications per Season**** | Maximum Rate per Acre per Season (ozs product) |
|--|--|--|--|---|
| Berry subgroups | 0 | 8 | 4 | 32 |
| Bulb vegetable group | 7 | 6.8 | 6 | 41 |
| Fruiting vegetable group | 0 | 3.5 | 6 | 21 |
| Tomato*** | | 12.5 | 2 | 25 |
| Low growing berry subgroup (except cranberry) | 0 | 8 | 5 | 40 |
| Oilseed crops (except rapeseed and sunflower) | 21 | 6 | 2 | 12 |
| Pome fruits group | 0 | 6.5 | 4 | 26 |
| Rapeseed | 21 | 6 | 2 | 12 |
| Small fruit, vine climbing subgroup (except fuzzy kiwifruit) | 14 | 8 | 5 | 24 |
| Sunflower | 21 | 9 | 2 | 18 |
| Turnip greens | 14 | 9 | 2 | 18 |

*See Table 2. Endura® fungicide Crop-specific Requirements for complete directions and exceptions.

**For a complete list of crops within a crop group or crop subgroup, see Table 2. Endura® fungicide Crop-specific Requirements.

***Maximum rate per acre per application, number of applications per season, and maximum rate per acre per season vary for tomato depending on the target disease. Refer to Table 2. Crop-specific Requirements for fruiting vegetables for maximum rates and number of applications by target disease.

****At the maximum use rate only, except for tomato.

Table 2. Endura® fungicide Crop-specific Requirements

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|---|--|---|---|---|
| <p>Berry subgroups</p> <p>Bushberry subgroup Aronia berry Black currant Blueberry (highbush and lowbush) Buffalo currant Chilean guava Currant Elderberry European barberry Gooseberry Highbush cranberry Honeysuckle, edible Huckleberry Jostaberry Juneberry Lingonberry Native currant Red currant Salal Sea buckthorn</p> <p>Caneberry subgroup Blackberry (all varieties) Loganberry Raspberry (black and red) Wild raspberry</p> | <p>Botrytis gray mold (<i>Botrytis cinerea</i>)</p> | <p>8</p> | <p>4</p> | <p>32</p> | <p>0</p> |

Application Directions. Begin applications of **Endura** prior to disease development and continue on a 7 to 14 day interval.

DO NOT apply by air.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Endura** per season.

DO NOT make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|---|--|---|---|---|
| Bulb vegetable group Chive, fresh leaves Chive, Chinese, fresh leaves Daylily, bulb Elegans hosta Fritillaria, bulb Fritillaria, leaves Garlic, bulb Garlic, great-headed, bulb Garlic; Serpent, bulb Kurrat Lady's leek Leek Leek, wild Lily, bulb Onion, Beltsville bunching Onion, bulb Onion, Chinese, bulb Onion, fresh Onion, green Onion, macrostem Onion, pearl Onion, potato, bulb Onion, tree, tops Onion, Welsh, tops Shallot, bulb Shallot, fresh leaves Cultivars, varieties and/or hybrids of these | Botrytis leaf blight (<i>Botrytis</i> spp.) Purple blotch (<i>Alternaria porri</i>) | 6.8 | 6 | 41 (In-furrow plus foliar) | 7 |

Application Directions. For control of purple blotch and Botrytis leaf blight, begin applications of **Endura** prior to disease development and continue on a 7 to 14 day interval.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than six (6) applications of **Endura** per season.

DO NOT make more than two (2) applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

(continued)

Table 2. Endura® fungicide | Crop-specific Requirements (continued)

Instructions for In-furrow Use to Aid in the Control of Soilborne White Rot (*Sclerotium cepivorum*) in the Bulb Vegetables Group, including Garlic and Onions

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|--|--|---|---|---|
| Bulb vegetable group Chive, fresh leaves Chive, Chinese, fresh leaves Daylily, bulb Elegans hosta Fritillaria, bulb Fritillaria, leaves Garlic, bulb Garlic, great-headed, bulb Garlic, Serpent, bulb Kurrat Lady's leek Leek Leek, wild Lily, bulb Onion, Beltsville bunching Onion, bulb Onion, Chinese, bulb Onion, fresh Onion, green Onion, macrostem Onion, pearl Onion, potato, bulb Onion, tree, tops Onion, Welsh, tops Shallot, bulb Shallot, fresh leaves Cultivars, varieties and/or hybrids of these | White rot (<i>Sclerotium cepivorum</i>) | 6.8 | 6 | 41 (In-furrow plus foliar) | 7 |

Application Directions. Apply **Endura** at planting as an in-furrow spray by directing spray pattern to the soil bed. Use a 4 to 6 inch band spray pattern applied directly over/into the seed furrow before covering with soil. Depending upon the level of potential infection, make additional foliar applications at the 6.8 ozs/A rate. Use a minimum volume of application of 5 gallons of water per acre.

Apply no more than a total of 41 ozs/A per season for a combination of the in-furrow and foliar uses.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than six (6) applications of **Endura** per season.

DO NOT make more than two (2) applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|--|--|---|---|---|
| Fruiting vegetable group African eggplant Bush tomato Bell pepper Cocona Currant tomato Eggplant Garden huckleberry Goji berry Groundcherry Martynia Naranjilla Okra Pea eggplant Pepino Pepper (all varieties) Nonbell pepper Roselle Scarlet eggplant Sunberry Tomatillo Tree tomato Cultivars, varieties and/or hybrids of these | Early blight (<i>Alternaria solani</i>) | 2.5 to 3.5 or 2.5 to 3.5 ozs per 100 gal of spray volume (dilute)* | 6 | 21 | 0 |
| | Target spot (<i>Corynespora cinerea</i>) Suppression Only: Botrytis gray mold** (<i>Botrytis cinerea</i>) | 3.5 | | | |
| Tomato | Black mold (<i>Alternaria alternata</i>) | 3.5 to 5 | 5 | 25 | |
| | Botrytis gray mold (<i>Botrytis cinerea</i>) | 9 to 12.5 | 2 | | |
| <p>Application Directions. Begin applications of Endura prior to disease development and continue on a 7 to 14 day interval for early blight, Botrytis gray mold, and Black mold.</p> <p>Use the higher rate and the shorter interval when disease pressure is high.</p> <p>Resistance Management. To limit the potential for development of resistance, DO NOT make more than six (6) applications of Endura at the low rate or two (2) applications of Endura at the high rate per season. DO NOT make more than two (2) sequential applications of Endura before alternating to a labeled fungicide with a different mode of action.</p> <p>*For applications based on dilute volume, plants should be sprayed to runoff. Apply a minimum of 20 gallons of spray volume per acre, and increase the spray volume as the plants grow during the season. Spray volume should be proportional to the amount of plant tissue to such that 100 gallons of spray per acre is used on mature plants.</p> <p>**Not registered for use in California.</p> | | | | | |

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|--|--|---|---|---|
| Low growing berry subgroup (except cranberry) Bearberry Bilberry Cloudberry Muntries Partridgeberry Strawberry | Botrytis gray mold (<i>Botrytis cinerea</i>) | 8 | 5 | 40 | 0 |
| <p>Application Directions. Begin applications of Endura no later than 10% bloom or prior to the onset of disease and continue on a 7 to 14 day interval.</p> <p>Use the shorter interval when disease pressure is high.</p> <p>Resistance Management. To limit the potential for development of resistance, DO NOT make more than five (5) applications of Endura per season.</p> <p>DO NOT make more than three (3) sequential applications of Endura before alternating to a labeled fungicide with a different mode of action.</p> | | | | | |

Table 2. Endura® fungicide crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|---|--|---|---|---|
| Oilseed crops (except rapeseed and sunflower) Flax seed | Pasmó <i>(Septoria linicola)</i> | 5 to 6 | 2 | 12 | 21 |
| Borage Calendula Castor oil plant Chinese tallowtree Crambe Cuphea Echium Euphorbia Evening primrose Gold of pleasure Hare's ear mustard Jojoba Lesquerella Lunaria Meadowfoam Milkweed Mustard seed Niger seed Oil radish Poppy seed Rose hip Safflower Sesame Stokes aster Sweet rocket Tallowwood Tea oil plant Vernonia | Sclerotinia rot and blight <i>(Sclerotinia spp.)</i> <u>Suppression Only:</u> Rust <i>(Puccinia helianthi)</i> | | | | |
| <p>Application Directions. For optimal disease control, begin applications of Endura prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development.</p> <p>Use the higher rate and shorter interval when disease pressure is high.</p> <p>No restriction on livestock grazing or feeding.</p> <p>Resistance Management. To limit the potential for development of resistance, DO NOT make more than two (2) applications of Endura per season.</p> | | | | | |

(continued)

Table 2. Endura® fungicide | Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|--|--|---|---|---|
| Pome fruits group Apple Azarole Crabapple Loquat Mayhaw Medlar Pear Pear, Asian Pear, Oriental Quince Quince, Chinese Quince, Japanese Tejocote Cultivars, varieties and/or hybrids of these | Apple scab (<i>Venturia inaequalis</i>) Pear scab (<i>Venturia pirina</i>) Powdery mildew (<i>Podosphaera leucotricha</i>) Fly speck (<i>Zygothiala jamaicensis</i>) Alternaria blotch (<i>Alternaria mali</i>) | 6.5 | 4 | 26 | 0 |

Application Directions for scab and powdery mildew. Begin applications of **Endura** prior to disease development and continue on a 7 to 10 day interval.

Use the shorter interval when disease pressure is high.

Application Directions for fly speck and Alternaria blotch. Begin applications of **Endura** prior to disease development and continue on a 7 to 14 day interval. Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Endura** per season.

DO NOT make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---|--|--|---|---|---|
| Rapeseed (cultivars, varieties, and/or hybrids, including canola and crambe) | White mold/Sclerotinia stem rot <i>(Sclerotinia sclerotiorum)</i> | 5 to 6 | 2 | 12 | 21 |
| <p>Application Directions. Begin applications of Endura at 20% to 50% flowering or prior to the onset of disease. Use the higher rate for extended protection. Apply a second time if conditions continue to be favorable for disease development.</p> <p>No restriction on livestock grazing or feeding.</p> <p>Resistance Management. To limit the potential for development of resistance, DO NOT make more than two (2) applications of Endura per season.</p> | | | | | |

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|--|---|--|---|---|---|
| Small fruit, vine climbing subgroup (except fuzzy kiwifruit) Amur river grape Gooseberry Grape Kiwifruit, hardy Maypop Schisandra berry | Powdery mildew <i>(Uncinula necator)</i> | 4.5 | 5 | 24 | 14 |
| | Botrytis gray mold <i>(Botrytis cinerea)</i> | 8 | 3 | | |

Application Directions. For powdery mildew control, begin applications of **Endura** at budbreak or prior to the onset of disease and continue on a 10 to 14 day interval.

For the control of Botrytis gray mold, begin applications of **Endura** prior to disease development and when conditions favor disease development during early bloom, bunch pre-closure or veraison.

Use the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Endura** at the low rate or three (3) applications of **Endura** at the high rate per season.

DO NOT make more than two (2) sequential applications of **Endura** before alternating to a labeled fungicide with a different mode of action.

Table 2. Endura® fungicide Crop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|------------|--|--|---|---|---|
| Sunflower* | Alternaria leaf spot (<i>Alternaria</i> spp.) Botrytis gray mold (<i>Botrytis cinerea</i>) Powdery mildew (<i>Erysiphe cichoracearum</i>) | 4.5 to 9 | 2 | 18 | 21 |
| | <u>Suppression Only:</u> Sclerotinia head rot (<i>Sclerotinia sclerotiorum</i>) | 6 to 9 | | | |

Application Directions. For optimal disease control, begin applications of **Endura** prior to disease development and continue on a 7 to 14 day interval if conditions are conducive for disease development.

Use the higher rate and shorter interval when disease pressure is high.

No restriction on livestock grazing or feeding.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

*Not registered for use in California.

Table 2. Endura® fungicide | Drop-specific Requirements (continued)

| Crop | Target Disease | Product Use Rate per Application (ozs/A) | Maximum Number of Applications per Season | Maximum Product Rate per Season (ozs/A) | Minimum Time from Application to Harvest (PHI) (days) |
|---------------|--|--|---|---|---|
| Turnip greens | Alternaria blight (Black spot) (<i>Alternaria</i> spp.) Gray mold (<i>Botrytis cinerea</i>) Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i> , <i>S. minor</i>) Suppression Only: Powdery mildew (<i>Erysiphe polygoni</i>) Rhizoctonia bottom rot (<i>Rhizoctonia solani</i>) | 6 to 9 | 2 | 18 | 14 |

Application Directions. Begin applications of **Endura** prior to disease development and continue on a 7 to 14-day interval.

Use the higher rate and the shorter interval when disease pressure is high.

Resistance Management. To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Endura** per season.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Endura is a registered trademark of BASF.

©2013 BASF Corporation
All rights reserved.

007969-00197.20120621e.NVA 2012-04-201-0130

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



The Chemical Company