



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

November 10, 2021

Quinn Peacock  
Product Registration Manager  
BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709

Subject: Label Amendment – Addition of California restriction to subgroup 6C and crop-specific restriction section; Incorporation of ID mitigation language for boscalid and pyraclostrobin; Other minor label revisions  
Product Name: Pristine Fungicide  
EPA Registration Number: 7969-199  
Application Date: October 12, 2020  
Decision Number: 567443

Dear Ms. Peacock:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Page 2 of 2  
EPA Reg. No. 7969-199  
Decision No. 567443

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Lindsay DeMers via email at demers.lindsay@epa.gov.

Sincerely,

A handwritten signature in black ink, reading "Shaja B. Joyner". The signature is written in a cursive style with a large initial 'S'.


Shaja B. Joyner, Product Manager 20  
Fungicide-Herbicide Branch  
Registration Division 7505P

Enclosure



We create chemistry

Boscalid	Group	7	Fungicide
Pyraclostrobin	Group	11	Fungicide



# Pristine®

## Fungicide

For use in disease control and plant health in the following crops: alfalfa; Belgium endive; berries; *Brassica*, head and stem; *Brassica*, leafy greens (except watercress); bulb vegetables; carrot; celery; celtuce; citrus fruit; cotton; cucurbit vegetables; dill seed; Florence fennel; fruiting vegetables; globe artichoke; grape; herbs; hops; kohlrabi; leaf petiole vegetables; leafy greens; low growing berry (except cranberry); oilseed crops; pea and bean, dried shelled, except soybean, subgroup 6C; peanut; persimmon; pome fruit; root vegetables (except sugar beet); small fruit, vine climbing (except fuzzy kiwifruit); soybean; spinach; stone fruit; strawberry; tree nut; tropical fruits (listed); and turnip greens

### Active Ingredients:

pyraclostrobin\*: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester) . . . . . 12.8%  
boscalid\*\*: 3-pyridinecarboxamide,2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl)- . . . . . 25.2%

**Other Ingredients:** . . . . . 62.0%

**Total:** . . . . . 100.0%

\* 0.128 oz (0.008 lb) of pyraclostrobin in 1 oz of product

\*\* 0.252 oz (0.0158 lb) of boscalid in 1 oz of product

EPA Reg. No. 7969-199

EPA Est. No.

### KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See full label for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

**In case of an emergency endangering life or property involving this product, call day or night, 1-800-832-HELP (4357).**

### Net Contents:

BASF Corporation  
26 Davis Drive, Research Triangle Park, NC 27709

<b>FIRST AID</b>	
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• <b>DO NOT</b> induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• <b>DO NOT</b> give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).	

## Precautionary Statements

### Hazards to Humans and Domestic Animals

**CAUTION.** Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Protective eyewear (goggles, face shield or safety glasses)
- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks

### User Safety Requirements

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## USER SAFETY RECOMMENDATIONS

#### Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove clothing/PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

### Groundwater Advisory

Boscalid and pyraclostrobin are known to leach through soil into groundwater under certain conditions as a result of label use. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of boscalid and

pyraclostrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

## Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours** for all crop uses except when performing cane tying, cane turning, cane girdling or leaf pulling tasks on grapes. The REI is **5 days** for treated grapes when conducting cane tying, cane turning, cane girdling or leaf pulling tasks.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Protective eyewear (goggles, faceshield or safety glasses)
- Coveralls
- Chemical-resistant gloves made of: barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, natural rubber  $\geq$  14 mils, polyethylene, polyvinyl chloride  $\geq$  14 mils, or viton  $\geq$  14 mils
- Shoes plus socks

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food, or feed by storage or disposal.

### Pesticide Storage

Store in original containers only. Keep container closed when not in use. **DO NOT** store near food or feed.

### Pesticide Disposal

Wastes resulting from using this product may be disposed of on-site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your state pesticide agency or environmental control agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

### Container Handling

**Nonrefillable Container. DO NOT reuse or refill this container.** Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

#### Triple rinse containers small enough to shake (capacity $\leq$ 50 pounds) as follows:

Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

#### Triple rinse containers too large to shake (capacity $>$ 50 pounds) as follows:

Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or mix tank. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

## In Case of Emergency

In case of large-scale spillage regarding this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

### Steps to be taken in case material is released or spilled:

- In case of spill on floor or paved surfaces, mop or sweep spill; then remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.
- Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing, and wash affected skin areas with soap and water.
- Wash clothing before reuse.
- Keep the spill out of all sewers and open bodies of water.

---

## Product Information

---

This package contains **Pristine® fungicide**, a water dispersible granule (WG). The active ingredients in **Pristine** belong to two classes of fungicides, the strobilurins and anilides. Preventive applications optimize disease control resulting in improved plant health. The increase in plant health comes from the combined effect of disease control (including fungal diseases listed in Crop-specific directions), improved growth efficiency and improved stress response. Overall increased plant health may result in an improvement in crop growth and crop quality as well as increased crop yields. **Pristine** is effective against pathogens resistant to other fungicides. **Pristine** has a protective effect because it inhibits spore germination. It also has a curative effect because it inhibits mycelial growth and sporulation of the fungus on the leaf surface. However, optimum disease control is achieved when **Pristine** is applied in a regularly scheduled protective spray program and is used in a rotation program with other fungicides. Because of its high specific activity and rain-fastness, **Pristine** has good residual activity against target fungi.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>.

**Pristine** is not for use in greenhouse or transplant production.

### Sensitive Crop Precaution

**Grapes** - **DO NOT** use on Concord or Noiret (NY73.0136.17) due to foliar injury. Possible foliar injury could occur to Worden, Fredonia, Niagara, Steuben,

Rougeon or related grape varieties. Use special care when applying **Pristine** to prevent contact with these sensitive varieties. Not all varieties have been thoroughly tested. Consult a BASF representative for more information concerning these sensitive grape varieties. Thoroughly rinse spray equipment, including the inside of the tank, hoses and nozzles after and before using the same equipment on grape varieties sensitive to **Pristine**.

**Blueberry (highbush and lowbush)** - **DO NOT** apply **Pristine** to blueberries as a tank mix with other pesticide products except fungicide products that contain captan (N-Trichloromethylthio-4-cyclohexene-1,2-dicarboxamide) as the **ONLY** active ingredient. **DO NOT** apply **Pristine** as a tank mix with adjuvants, liquid fertilizers, nutrients or other additives. Only use water as the spray carrier.

### Modes of Action

Pyraclostrobin and boscalid, the active ingredients of **Pristine**, belong to the groups of respiration inhibitors classified as target site of action **Group 11** and **Group 7** fungicides, respectively.

### Resistance Management

For resistance management, please note that **Pristine** contains both a **Group 7** (boscalid) and **Group 11** (pyraclostrobin) fungicide. Any fungal population may contain individuals naturally resistant to **Pristine** and other **Group 7** or **Group 11** fungicides. A gradual or total loss of pest control may occur over time if fungicides from these groups are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

**Pristine** provides effective resistance management of most of its target pathogens because it is a premix of two fungicides with different modes of action. **Pristine** is effective against pathogens resistant to fungicides with modes of action different from those of target site **Group 7** and **Group 11**, including dicarboximides, sterol inhibitors, benzimidazoles, or phenylamides. **Pristine** is also effective against certain pathogens with resistance to **Group 11** fungicides. However, fungal isolates resistant to **Group 7** or **Group 11** fungicides may eventually dominate the fungal population if **Group 7** or **Group 11** fungicides are used predominantly and repeatedly in the same field in successive years as the primary method of control for the targeted pathogen species. This may result in reduction of disease control by **Pristine** or other **Group 7** or **Group 11** fungicides, especially if resistance to either **Group 7** or **Group 11** fungicides is already present in the pathogen population. To maintain the performance of **Pristine** in the field, **DO NOT** exceed the specified number of applications of **Pristine** and the total number of applications of **Pristine** per year stated in **Restrictions and Limitations** and crop-specific use requirements. Adhere to the label instructions regarding the sequential use of **Pristine** or other target site of action **Group 7** and **Group 11** fungicides that have a similar site of action on the same pathogens.

To delay the development of fungicide resistance:

- **Rotate** - Rotate the use of **Pristine® fungicide** or other **Group 7** and **Group 11** fungicides within a growing season sequence with different mode of action groups that control the same pathogens.
- **Tank Mixtures** - Use tank mixtures with fungicides from different target site of action groups that are registered/ permitted for the same use and that are effective against the pathogens of concern. Use at least the minimum labeled rates of each fungicide in the tank mix.
- **Integrated Pest Management (IPM)** - Integrate **Pristine** into an overall disease and pest management program. Follow cultural practices known to reduce disease development. Consult your local extension specialist, certified crop advisor and/or BASF representative for additional IPM strategies established for your area. **Pristine** may be used in Agricultural Extension advisory (disease forecasting) programs which base application timing on environmental factors favorable for disease development.
- **Monitoring** - Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development. Monitor treated fungal populations for resistance development.
- **Reporting** - If a **Group 7** or **Group 11** target site fungicide, including **Pristine**, appears to be less effective or no longer effective against a pathogen that it previously controlled or suppressed, contact a BASF representative, local extension specialist, or certified crop advisor to assist in determining the cause of reduced performance.

## Cleaning Spray Equipment

Clean spray equipment thoroughly before and after applying this product, particularly if a product with the potential to injure crops was used prior to **Pristine**.

---

## Application Instructions

---

Apply directed rates of **Pristine** as instructed by **Table 3. Pristine® fungicide Crop-specific Requirements**. Under low-level disease conditions, use the minimum application rates. Use maximum application rates and the shorter application interval for severe or threatening disease conditions. Ground application is advised for thorough coverage. **DO NOT** apply by air to dill seed, herb, peanut, and turnip greens. For all other crops listed on this label, aerial application can be made, including conditions where applications are not possible using ground equipment. **Pristine** can be applied through sprinkler irrigation equipment, except for use on dill seed, herb, and turnip greens. Check equipment frequently for calibration.

## Ground Application

Apply **Pristine** in sufficient water to ensure thorough coverage of foliage, bloom, and fruit. Thorough coverage is required for optimum disease control.

## Directed or Banded Sprays

The application rates on the **Pristine** label reflect the amount of product uniformly applied over an acre of ground on a broadcast basis.

In some crops, apply **Pristine** as a directed or banded spray over the rows or plant beds with the alleys or row middles left unsprayed. For such uses, reduce the labeled **Pristine** rates in proportion to the area actually sprayed. This adjustment is necessary to avoid applying the product at use rates higher than permitted according to label directions.

Use the following formula to determine the broadcast equivalent rate for doing directed or banded sprays:

$$\begin{array}{r} \text{sprayed bed} \\ \text{width} \end{array} + \begin{array}{r} \text{unsprayed row} \\ \text{middles width} \end{array} = \begin{array}{r} \text{total} \\ \text{row width} \end{array}$$

$$\frac{\begin{array}{r} \text{sprayed bed} \\ \text{width in inches} \end{array}}{\begin{array}{r} \text{total row} \\ \text{width in inches} \end{array}} \times \frac{\begin{array}{r} \text{broadcast rate} \\ \text{treated acre} \end{array}}{\begin{array}{r} \text{band rate} \\ \text{field acre} \end{array}} = \frac{\begin{array}{r} \text{band rate} \\ \text{field acre} \end{array}}{\begin{array}{r} \text{band rate} \\ \text{field acre} \end{array}}$$

**Example:** A directed spray application will be made to 45-inch plant beds that are separated by 15-inch unsprayed row middles.

$$\begin{array}{r} 45 \text{ inches} \\ \text{sprayed} \\ \text{bed width} \end{array} + \begin{array}{r} 15 \text{ inches} \\ \text{unsprayed} \\ \text{row middles} \end{array} = \begin{array}{r} 60 \text{ inches} \\ \text{total} \\ \text{row width} \end{array}$$

The calculation to determine the appropriate equivalent rate of product to use for this situation based on a label broadcast rate of 12 ozs/acre follows:

$$\frac{\begin{array}{r} 45 \text{ inches} \\ \text{sprayed} \\ \text{bed width} \end{array}}{\begin{array}{r} 60 \text{ inches} \\ \text{total row width} \end{array}} \times \frac{\begin{array}{r} 12 \text{ ozs} \\ \text{Pristine} \\ \text{treated acre} \end{array}}{\begin{array}{r} 9 \text{ ozs} \\ \text{Pristine} \\ \text{field acre} \end{array}} = \frac{\begin{array}{r} 9 \text{ ozs} \\ \text{Pristine} \\ \text{field acre} \end{array}}{\begin{array}{r} 9 \text{ ozs} \\ \text{Pristine} \\ \text{field acre} \end{array}}$$

## Aerial Application

**For aerial application in New York State, DO NOT apply within 100 feet of aquatic habitats (including, but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).**

**DO NOT** apply by air to dill seed, herb, peanut, and turnip greens. For all other crops listed on this label, aerial application can be made and thorough coverage is required to obtain optimum disease control. Avoid applications under conditions when uniform coverage cannot be obtained or when spray drift may occur. Use no less than 5 gallons of spray solution per acre. For aerial application to citrus fruit, grape, hops, pome fruit, stone fruit, and tree nut, use no less than 10 gallons of spray solution per acre. Thorough coverage is required for optimum disease control.

## MANDATORY SPRAY DRIFT MANAGEMENT

### Aerial Applications

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzle and pressure that deliver a medium or coarser droplet size (ASABE S641).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site. If the windspeed is greater than 10 miles per hour, the boom length must be 65% or less of the wingspan for fixed-wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- If the windspeed is 10 miles per hour or less, applicators must use 1/2 swath displacement upwind at the downwind edge of the field. When the windspeed is between 11 to 15 miles per hour, applicators must use 3/4 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

### Airblast Applications

- Sprays must be directed into the canopy.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

### Ground Boom Applications

- User must only apply with the release height recommended by the manufacturer, but no more than 4 ft above the ground or crop canopy.
- Applicators are required to use nozzles and pressure that deliver a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

## SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NONTARGET SITES AND ENVIRONMENTAL CONDITIONS.

### Boomless Ground Applications

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

### Handheld Technology Applications

Take precautions to minimize spray drift.

## Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### Controlling Droplet Size - Aircraft

**Adjust Nozzles** - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### Release Height - Aircraft

Higher release heights increase the potential for spray drift.

### Boom Height - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

### Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### Wind

AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Drift potential generally increases with wind speed. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.



## Directions for Use Through Sprinkler Irrigation Systems

### Sprayer Preparation

Clean chemical tank and injector system thoroughly. Flush system with clean water.

### Application Instructions

Apply **Pristine® fungicide** at rates and timings as described in this label.

### Use Precautions for Sprinkler Irrigation Applications

- **DO NOT** apply by sprinkler irrigation (chemigation) to dill seed, herb, and turnip greens. For all other crops, this product can be applied through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. **DO NOT** apply this product through any other type of irrigation system.
- Add this product to the pesticide supply tank containing sufficient water to maintain a continuous flow by the injection equipment. In continuous moving systems, inject this product-water mixture continuously, applying the labeled rate per acre for that crop. **DO NOT** exceed 1/2 inch (13,577 gallons) per acre. In stationary or non-continuous moving systems, inject the product-water mixture in the last 15 to 30 minutes of each set allowing sufficient time for all of the required pesticide to be applied by all the sprinkler heads and applying the labeled rate per acre for that crop. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. Thorough coverage of foliage is required for good control. Maintain good agitation during the entire application period.
- If you have questions about calibration, contact a state extension service specialist, equipment manufacturers or other experts.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, including a positive displacement injection pump (e.g., diaphragm pump),

effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, must shut the system down and make necessary adjustments when appropriate.
- **DO NOT** connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

### Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ) backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There must be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, including a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

---

## Additives and Tank Mixing Information

---

Under some conditions, the use of additives or adjuvants may improve the performance of **Pristine® fungicide**. However, all varieties and cultivars have not been tested with possible tank mix combinations. Local conditions can also influence crop response and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing **Pristine** with other products. Therefore, before using any tank mix (fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

Consult a BASF representative or local agricultural authorities for more information concerning additives.

If tank mixtures are used, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Pristine** can be tank mixed with most fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives. See **Table 3**.

**Pristine® fungicide Crop-specific Requirements** for exceptions.

### Compatibility Test for Tank Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of label rate per acre:

1. **Water** - For 100 gallons per acre spray volume, use 16 cups (1 gallon) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
2. **Water-dispersible products** (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions). Cap the jar and invert 10 cycles.
3. **Water-soluble products** - Cap the jar and invert 10 cycles.
4. **Emulsifiable concentrates** (oil concentrate or methylated seed oil when applicable). Cap the jar and invert 10 cycles.
5. **Water-soluble additives** - Cap the jar and invert 10 cycles.
6. Let the solution stand for 15 minutes.
7. **Evaluate** the solution for uniformity and stability. The spray solution must not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. **DO NOT** use any spray solution that could clog spray nozzles.

## Mixing Order

1. **Water** - Begin by filling a thoroughly clean sprayer tank 3/4 full of clean water.
2. **Agitation** - Maintain constant agitation throughout mixing and application.
3. **Inductor** - If an inductor is used, rinse it thoroughly after each component has been added.
4. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. **Water-dispersible products** (including **Pristine**, dry flowables, wettable powders, suspension concentrates, or suspo-emulsions)
6. **Water-soluble products**
7. **Emulsifiable concentrates** (including oil concentrates when applicable)
8. **Water-soluble additives** (including ammonium sulfate [AMS] or urea ammonium nitrate [UAN] when applicable)
9. **Remaining quantity of water**

Make sure that each component is thoroughly mixed and suspended before adding tank mix partners. Maintain constant agitation during application. See **Table 3**.

**Pristine® fungicide Crop-specific Requirements** for more details.

---

## Restrictions and Limitations

---

- **DO NOT** exceed the maximum product rate (ozs/A) per year, the maximum product rate per application, or the total number of applications of **Pristine** per year as stated in **Table 1. Pristine® fungicide Restrictions and Limitations Overview** and **Table 3. Pristine® fungicide Crop-specific Requirements**. Preharvest interval (PHI) restrictions are also included in these tables.
- **DO NOT** apply more than the maximum annual use rate of ai/acre or ozs of product/acre for each specific crop from any combination of products containing pyraclostrobin or boscalid (including **Pristine**, **Endura® fungicide**, **Cabrio® EG fungicide**, **Headline® fungicide**). To determine lbs of pyraclostrobin per acre, multiply ozs of product/acre by 0.008. To determine lbs of boscalid per acre, multiply ozs of product/acre by 0.0158. See **Table 2. Pristine® fungicide Rate Conversions** for corresponding pounds active ingredient per acre.
- **Pristine** is not for use in greenhouse or transplant production.
- **Blueberry (highbush and lowbush) - DO NOT** apply **Pristine** to blueberries as a tank mix with other pesticide products except fungicide products that contain captan (N-Trichloromethylthio-4-cyclohexene-1,2-dicarboxamide) as the ONLY active ingredient. **DO NOT** apply **Pristine** as a tank mix with adjuvants, liquid fertilizers, nutrients, or other additives. Only use water as the spray carrier.

- **Grape - DO NOT** use on Concord or Noiret (NY73.0136.17) due to foliar injury. Possible foliar injury could occur to Worden, Fredonia, Niagara, Steuben, Rougeon or related grape varieties. Not all varieties have been thoroughly tested.
- **Aerial application in hops - DO NOT** make more than one (1) aerial application of **Pristine® fungicide** per year and include a myclobutanil product as a tank mix.
- **DO NOT** apply by air to dill seed, herb, peanut, and turnip greens.
- **DO NOT** apply through any type of irrigation system (chemigation) to dill seed, herb, and turnip greens.
- **For aerial application in New York State, DO NOT apply within 100 feet of aquatic habitats (including, but not limited to lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).**
- **DO NOT** use on **sugar beet**.

### **Crop Rotation Restriction**

Crops listed on the **Pristine** label may be planted immediately following the last application. For all other crops, **DO NOT** plant sooner than 14 days after the last application.

**Table 1. Pristine® fungicide Restrictions and Limitations Overview\***

<b>Crop/Crop Group**</b>	<b>Minimum Time from Application to Harvest (PHI) (days)</b>	<b>Maximum Rate per Application (ozs product/A)</b>	<b>Maximum Number of Applications per Year***</b>	<b>Maximum Rate per Year (ozs product/A)</b>
<b>Alfalfa (including alfalfa grown for seed)</b>	14	18	3	54
<b>Belgium endive</b>	19	1.6 (cold storage)	1	3.4
		1.8 (forcing)	1	
<b>Berry subgroups</b>				
<b>Bushberry [subgroup 13-07B]</b>	0	23	4	92
<b>Caneberry [subgroup 13-07A]</b>	0	23	4	92
<b>Low growing berry (except cranberry) [subgroup 13-07G]</b>	0	23	5	115
<b>Small fruit, vine climbing except fuzzy kiwifruit [subgroup 13-07F]</b>	14	23	5	69
<b>Brassica, head and stem [group 5-16]</b>	0	25	2	50
<b>Brassica, leafy greens (except watercress), [subgroup 4-16B]</b>	14	25	2	50
<b>Bulb vegetables [group 3-07]</b>	7	18.5	6	111
<b>Carrot</b>	0	10.5	6	63
<b>Celery</b>	0	25	2	50
<b>Celtuce</b>	0	25	2	50
<b>Citrus fruit [group 10-10]</b>	0	18.5	4	74
<b>Cotton</b>	30	25	2	50
<b>Cucurbit vegetables [group 9]</b>	0	18.5	4	74
<b>Dill seed****</b>	0	18.5	2	37
<b>Florence fennel</b>	0	25	2	50
<b>Fruiting vegetables [group 8-10]</b>	0	9.7	6	58.2
<b>Tomato</b>		25	2	69
<b>Globe artichoke</b>	0	23	3	69
<b>Grape</b>	14	23	3	69
<b>Herb [subgroup 19A]****</b>	0	18.5	2	37
<b>Hops****</b>	14	28	3	84
<b>Kohlrabi</b>	0	25	2	50
<b>Leaf petiole vegetables [subgroup 22B]</b>	0	25	2	50
<b>Leafy greens [subgroup 4-16A]</b>	14	25	2	50
<b>Oilseed [group 20]</b>	21	24.5	2	49
<b>Pea and bean, dried shelled, except soybean, [subgroup 6C]</b>	21	25	2	50
<b>Peanut</b>	14	28	3	84
<b>Persimmon</b>	0	23	3	69
<b>Pome fruit [group 11-10]</b>	0	18.5	4	74

(continued)

**Table 1. Pristine® fungicide Restrictions and Limitations Overview\*** (continued)

Crop/Crop Group**	Minimum Time from Application to Harvest (PHI) (days)	Maximum Rate per Application (ozs product/A)	Maximum Number of Applications per Year***	Maximum Rate per Year (ozs product/A)
Root vegetables (except sugar beet) [subgroup 1B]	0	10.5	6	63
Soybean	21	16	2	32
Spinach	14	25	2	50
Stone fruit [group 12-12]	0	14.5	5	72.5
Strawberry	0	23	5	115
Tree nut [group 14-12]	14 (for almond - 25 days)	14.5	4	58
Tropical fruits (listed in Table 3)	0	18.5	2	37
Turnip greens****	14	25	2	50

\* See **Table 3. Pristine® fungicide Crop-specific Requirements** for complete directions and exceptions, including restrictions and information regarding crop sensitivity and tank mixtures.

\*\* For a complete list of crops labeled within a group, see **Table 3. Pristine® fungicide Crop-specific Requirements**.

\*\*\* See **Table 3. Pristine® fungicide Crop-specific Requirements** for exceptions.

\*\*\*\* For additional ground, aerial, and sprinkler irrigation (chemigation) application restrictions and limitations, see **Table 3. Pristine® fungicide Crop-specific Requirements**.

**Table 2. Pristine® fungicide Rate Conversions\***

<b>Product Use Rate (oz/A)</b>	<b>lb ai boscalid</b>	<b>lb ai pyraclostrobin</b>
0.8	0.013	0.006
0.9	0.014	0.007
1.6	0.025	0.013
1.8	0.028	0.014
3.4	0.054	0.027
8	0.126	0.064
9.7	0.153	0.078
10	0.158	0.080
10.5	0.166	0.084
12	0.190	0.096
12.5	0.198	0.100
14	0.221	0.112
14.5	0.229	0.116
15	0.237	0.120
16	0.253	0.128
16.7	0.264	0.134
18	0.284	0.144
18.5	0.292	0.148
23	0.363	0.184
24.5	0.387	0.196
25	0.395	0.200
28	0.442	0.224
32	0.506	0.256
33.4	0.528	0.267
37	0.585	0.296
49	0.774	0.392
50	0.790	0.400
54	0.853	0.432
58	0.916	0.464
58.2	0.920	0.466
63	0.995	0.504
69	1.09	0.552
72.5	1.15	0.580
74	1.17	0.592
84	1.33	0.672
92	1.45	0.736
111	1.75	0.888
115	1.82	0.920

\* Corresponding pounds active ingredient per acre for Product Use Rates (oz/A) in **Table 3. Pristine® fungicide Crop-specific Requirements**. Also see **Table 3. Pristine® fungicide Crop-specific Requirements** for the maximum product use rate per year in ozs/A and lb ai.

## Crop-specific Requirements

**Table 3. Pristine® fungicide Crop-specific Requirements**

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Alfalfa (including alfalfa grown for seed)</b>	Anthracnose <i>Colletotrichum trifolii</i> Common leaf spot <i>Pseudopeziza medicaginis</i> Downy mildew <i>Peronospora trifoliorum</i> Leaf spot <i>Leptosphaerulina briosiani</i> Powdery mildew <i>Erysiphe pisi</i> Rhizoctonia blight/ Black patch <i>Rhizoctonis</i> spp. Rust <i>Uromyces</i> spp. Spring black stem and Leaf spot <i>Phoma medicaginis</i> Stagonospora leaf spot <i>Stagonospora meliloti</i> Stemphyllium leaf spot <i>Stemphyllium</i> spp. Summer black stem and Leaf spot <i>Cercospora medicaginis</i> Yellow leaf blotch <i>Leptotrichila medicaginis</i>	12 to 18	2 per cutting  or  3 total per year*	54	14
	White mold/Sclerotinia crown and Stem rot <i>Sclerotinia sclerotiorum</i> , <i>S. trifoliorum</i> <b>Suppression Only:</b> Southern blight <i>Sclerotium rolfsii</i>	14 to 18			

(continued)

**Table 3. Pristine® fungicide Crop-specific Requirements** *(continued)*

**Alfalfa (including alfalfa grown for seed)** *(continued)*

**Application Directions.** Begin **Pristine** applications when conditions favorable for disease are expected, but prior to onset of disease development. For stand establishment of fall-seeded alfalfa, begin applications in fall through early winter prior to first snowfall or extended cool, wet conditions. For seed pod protection, begin applications at 10% to 30% bloom. Use the higher rate and shorter interval when disease pressure is high.

Using higher rates may improve disease control performance as the crop canopy volume and density increases. Disease control can also be improved when application equipment and spray volume is adjusted to achieve thorough canopy penetration and coverage.

Repeat application on a 14-day to 21-day interval if conditions are favorable for disease development.

Under some conditions, additives or adjuvants may improve the performance of **Pristine**.

No livestock feeding restrictions.

**Resistance Management.** To limit development of resistance, **DO NOT** make more than two (2) sequential **Pristine** applications per cutting or three (3) **Pristine** applications per year. Alternate to a labeled **non-Group 7** or **non-Group 11** fungicide with different mode of action following two (2) sequential **Pristine** applications.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (18 ozs/A contains 0.284 lb boscalid, 0.144 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (54 ozs/A contains 0.853 lb boscalid, 0.432 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 18 ozs/A (0.284 lb boscalid, 0.144 lb pyraclostrobin).
- Maximum annual application rate is 54 ozs/A (0.853 lb boscalid, 0.432 lb pyraclostrobin) per year.
- **DO NOT** make more than two (2) **Pristine** applications per cutting or more than three (3) **Pristine** applications per year at the high application rate.
- Minimum retreatment interval is 14 days.



**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application	Maximum Number of Applications per Year	Maximum Product Rate per Crop per Year (ozs product per 1000 lbs roots)	Minimum Time from Application to Harvest (PHI) (days)
Belgium endive*	Root and crown rot <i>Sclerotinia sclerotiorum</i>	<b>Prior to cold storage:</b> 0.8 to 1.6 ozs per 1000 lbs roots	1	3.4	19
		<b>Prior to forcing:</b> 0.9 to 1.8 ozs per 70 square feet of forcing tray	1		

**Application Directions. Dosage and frequency/timing of applications.** Make one application to the roots when brought into cold storage prior to forcing. Apply again at the beginning of forcing after the roots have been packed in forcing trays.

**Prior to Cold Storage.** Make one application as a spray to the roots as they move along a conveyor belt used to bring roots from field transportation into cold storage bins. Apply 0.8 to 1.6 ozs **Pristine** in 3.0 to 3.5 gals of water per 1000 lbs roots.

**Prior to Forcing.** Make one application as a spray to the roots at the beginning of forcing, after they have been packed into forcing trays. Apply at the rate of 0.9 to 1.8 ozs of **Pristine** in approximately 100 fl ozs of water per 70 square feet of forcing tray. Approximately 1000 lbs of roots will fill 70 square feet of forcing tray.

**Restrictions and Limitations**

- Maximum single application rate is 1.6 ozs (0.025 lb boscalid, 0.013 lb pyraclostrobin) prior to cold storage and 1.8 ozs (0.028 lb boscalid, 0.014 lb pyraclostrobin) prior to forcing.
- Maximum annual application rate is 3.4 ozs **Pristine** (0.054 lb boscalid, 0.027 lb pyraclostrobin) per 1000 lbs roots per crop per year.
- **DO NOT** make more than two (2) **Pristine** applications per year, one (1) **Pristine** application prior to cold storage and one (1) **Pristine** application prior to forcing.
- **DO NOT** apply after the beginning of forcing.
- Minimum time from application to harvest (PHI) is 19 days.

\* For use in California only.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Berry subgroups</b></p> <p><b>Bushberry [subgroup 13-07B]*</b>                      Aronia berry                      Black currant                      Blueberry**                      (highbush and lowbush)                      Buffalo currant                      Chilean guava                      Currant                      Elderberry                      European barberry                      Gooseberry                      Highbush cranberry                      Honeysuckle, edible                      Huckleberry                      Jostaberry                      Juneberry                      Lingonberry                      Native currant                      Red currant                      Salal                      Sea buckthorn</p> <p><b>Caneberry [subgroup 13-07A]*</b>                      Blackberry                      (all varieties)                      Loganberry                      Raspberry                      (black and red)                      Wild raspberry</p>	<p>Alternaria leaf spot and fruit rot  <i>Alternaria</i> spp.</p> <p>Anthracnose  <i>Colletotrichum</i> spp.,  <i>Elsinoe</i> spp.</p> <p>Botrytis gray mold  <i>Botrytis cinerea</i></p> <p>Leaf spot and blotch  <i>Mycosphaerella</i> spp.,  <i>Septoria</i> spp.</p> <p>Monilinia blight and mummy berry  <i>Monilinia</i> spp.</p> <p>Phomopsis leaf spot, twig blight, and fruit rot  <i>Phomopsis</i> spp.</p> <p>Powdery mildew  <i>Microsphaera</i> spp.,  <i>Oidium</i> spp.,  <i>Sphaerotheca</i> spp.</p> <p>Spur blight  <i>Didymella</i> spp.,  <i>Phoma</i> spp.</p> <p><b>Suppression Only:</b></p> <p>Rust  <i>Arthuriomyces</i> spp.,  <i>Kuehneola</i> spp.,  <i>Phragmidium</i> spp.,  <i>Pucciniastrum</i> spp.</p>	18.5 to 23	4	92	0

(continued)

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

**Berry subgroups** (continued)

**Application Directions.** Begin applications of **Pristine** prior to onset of disease development and continue on a 7-day to 14-day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

\* **For the berries listed in this table (except blueberry),** it is impossible for BASF to test all bushberry and caneberry crops for sensitivity to **Pristine** under all environments and all potential product mixture combinations. Local conditions can also influence crop response and may not match those under which BASF has conducted testing. Proceed with caution with regard to **Pristine** use, particularly in tank mixes and/or adjuvant combinations on bushberry and caneberry crops. To reduce the risk of berry crop injury, BASF advises testing **Pristine** or **Pristine** tank mixtures on a small portion of the crop before broad scale use. To the extent consistent with applicable law, the user assumes all risks associated with adding products to the **Pristine** spray solution. Refer also to the **Conditions of Sale and Warranty** section of this label.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

- Maximum single application rate is 23 ozs/A (0.363 lb boscalid, 0.184 lb pyraclostrobin).
- Maximum annual application rate is 92 ozs/A (1.45 lbs boscalid, 0.736 lb pyraclostrobin) per year.
- **DO NOT** make more than four (4) applications of **Pristine** per year.
- Minimum retreatment interval is 7 days.

\*\* **Blueberry (highbush and lowbush)** is not registered for use in California. For all other states, **DO NOT** apply **Pristine** to blueberries as a tank mix with other pesticide products except fungicide products that contain captan (N-Trichloromethylthio-4-cyclohexene-1,2-dicarboxamide) as the **ONLY** active ingredient. **DO NOT** apply **Pristine** as a tank mix with adjuvants, liquid fertilizers, nutrients, or other additives. Only use water as the spray carrier.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Berry subgroups</b> (continued)  <b>Low growing berry (except cranberry) [subgroup 13-07G]**</b>  Bearberry Bilberry Cloudberry Muntries Partridgeberry	Anthracnose <i>Colletotrichum</i> spp.  Botrytis gray mold <i>Botrytis cinerea</i>  Leaf spot <i>Mycosphaerella fragariae</i>  Powdery mildew <i>Sphaerotheca macularis</i>	18.5 to 23	5*	115	0

**Application Directions.** Begin applications of **Pristine** no later than 10% bloom, or prior to disease development, and continue on a 7-day to 14-day interval. Use the higher rate and the shorter interval when disease pressure is high.

**For the low growing berries listed in this table,** it is impossible for BASF to test all low growing berry crops for sensitivity to **Pristine** under all environments and all potential product mixture combinations. Local conditions can also influence crop response and may not match those under which BASF has conducted testing. Proceed with caution with regard to **Pristine** use, particularly in tank mixes and/or adjuvant combinations on low growing berry crops. To reduce the risk of berry crop injury, BASF advises testing **Pristine** or **Pristine** tank mixtures on a small portion of the crop before broad scale use. To the extent consistent with applicable law, the user assumes all risks associated with adding products to the **Pristine** spray solution. Refer also to the **Conditions of Sale and Warranty** section of this label.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (23 ozs/A contains 0.363 lb boscalid, 0.184 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (115 ozs/A contains 1.82 lbs boscalid, 0.920 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 23 ozs/A (0.363 lb boscalid, 0.184 lb pyraclostrobin).
- Maximum annual application rate is 115 ozs/A (1.82 lbs boscalid, 0.920 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.

\*\* See separate crop table for strawberry use rates and application directions.

(continued)

**Table 3. Pristine® fungicide Crop-specific Requirements** *(continued)*

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Berry subgroups</b> <i>(continued)</i></p> <p><b>Small fruit, vine climbing except fuzzy kiwifruit [subgroup 13-07F]**</b></p> <p>Amur river grape Gooseberry Kiwifruit, hardy Maypop Schisandra berry</p> <p>Cultivars, varieties and/or hybrids of these</p>	<p>Angular leaf spot <i>Mycosphaerella angulata</i></p> <p>Anthracnose <i>Elsinoe ampelina</i></p> <p>Black rot <i>Guignardia bidwellii</i></p> <p>Downy mildew <i>Plasmopara viticola</i></p> <p>Leaf blight <i>Pseudocercospora vitis</i></p> <p>Phomopsis cane and leaf spot <i>Phomopsis viticola</i></p> <p>Powdery mildew <i>Uncinula necator</i></p> <p>Ripe rot <i>Colletotrichum gloeosporioides</i></p> <p><b><u>Aids in Control Only:</u></b></p> <p>Summer bunch rot (Sour rot) <i>Aspergillus</i> spp. and <i>Cladosporium</i> spp.</p> <p><b><u>Suppression Only:</u></b></p> <p>Botrytis gray mold <i>Botrytis cinerea</i></p>	8 to 12.5	5*	69	14
	<p>Botrytis gray mold <i>Botrytis cinerea</i></p>	18.5 to 23	3		

*(continued)*

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

**Berry subgroups** (continued)

**Small fruit, vine climbing except fuzzy kiwifruit [subgroup13-07F]\*\*** (continued)

**Application Directions. For powdery mildew control,** begin applications of **Pristine** as of bud break prior to onset of disease, using 8 ozs per acre on a 10-day to 14-day interval. Use 10 to 12.5 ozs per acre on a 14-day to 21-day interval.

**For black rot and downy mildew control,** begin applications of **Pristine** as of pre-bloom prior to onset of disease and continue applications on a 10-day to 14-day interval.

**For all other diseases listed except for Botrytis gray mold,** begin applications of **Pristine** prior to onset of disease and continue applications on a 10-day to 14-day interval. **Pristine** applied at rates of 8 to 12.5 ozs per acre for control of the listed diseases will also suppress Botrytis gray mold.

**For control of Botrytis gray mold,** apply 18.5 to 23 ozs per acre of **Pristine** prior to onset of disease development when conditions favor disease development during early bloom, bunch pre-closure and veraison up to 14 days before harvest. Continue applications on a 10-day to 14-day interval.

Use the higher rate and the shorter interval when disease pressure is high.

**For the small fruit, vine climbing berries listed in this table,** it is impossible for BASF to test all small fruit, vine climbing berry crops for sensitivity to **Pristine** under all environments and all potential product mixture combinations. Local conditions can also influence crop response and may not match those under which BASF has conducted testing. Proceed with caution with regard to **Pristine** use, particularly in tank mixes and/or adjuvant combinations on small fruit, vine climbing berry crops. To reduce the risk of small fruit, vine climbing berry crop injury, BASF advises testing **Pristine** or **Pristine** tank mixtures on a small portion of the crop before broad scale use. To the extent consistent with applicable law, the user assumes all risks associated with adding products to the **Pristine** spray solution. Refer also to the **Conditions of Sale and Warranty** section of this label.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (23 ozs/A contains 0.363 lb boscalid, 0.184 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (69 ozs/A contains 1.09 lbs boscalid, 0.552 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 23 ozs/A (0.363 lb boscalid, 0.184 lb pyraclostrobin).
- Maximum annual application rate is 69 ozs/A (1.09 lbs boscalid, 0.552 lb pyraclostrobin) per year.
- Minimum retreatment interval is 10 days.
- Minimum time from application to harvest (PHI) is 14 days.

\*\* See separate crop table for grape use rates and application directions.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Brassica, head and stem [group 5-16]**</b> Broccoli Brussels sprouts Cabbage Chinese cabbage (napa) Cauliflower Cultivars, varieties and/or hybrids of these	Alternaria leaf spot <i>Alternaria</i> spp.	10 to 15	2*	50	0
	Black leg <i>Phoma lingam</i>				
	Cercospora leaf spot <i>Cercospora</i> spp.				
	Powdery mildew <i>Erysiphe</i> spp.				
	Ring spot <i>Mycosphaerella</i> spp.				
	White leaf spot <i>Pseudocercospora</i> spp.				
Anthracnose <i>Colletotrichum</i> spp.	15 to 20				
White rust <i>Albugo</i> spp.					
Downy mildew <i>Peronospora</i> spp.	15 to 25				
Gray mold <i>Botrytis</i> spp.					
Sclerotinia stem rot <i>Sclerotinia</i> spp.					
Rhizoctonia bottom rot <i>Rhizoctonia solani</i>					

**Application Directions.** Begin applications of **Pristine** prior to disease development and continue on a 7-day to 14-day interval. Use the higher rate and the shorter interval when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (25 ozs/A contains 0.395 lb boscalid, 0.200 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (50 ozs/A contains 0.790 lb boscalid and 0.400 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A (0.790 lb boscalid and 0.400 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.

\*\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Brassica, leafy greens (except watercress) [subgroup 4-16B]**</b> Arugula Broccoli, Chinese Broccoli raab Cabbage, abyssinian Cabbage, Chinese, bok choy Cabbage, seakale Collards Cress (garden and upland) Hanover salad Kale Maca, leaves Mizuna Mustard greens Radish, leaves Rape greens Rocket, wild Shepherd's purse Cultivars, varieties and/or hybrids of these	Alternaria leaf spot <i>Alternaria</i> spp. Anthracnose <i>Colletotrichum</i> spp. Ascochyta leaf spot <i>Ascochyta</i> spp. Cercospora leaf spot <i>Cercospora</i> spp. Downy mildew <i>Peronospora</i> spp. Phoma <i>Phoma</i> spp. Powdery mildew <i>Erysiphe</i> spp., <i>Phyllactinia</i> spp., <i>Sphaerotheca</i> spp.	10 to 15	2*	50	14
	Botrytis rot <i>Botrytis</i> spp. Sclerotinia rot and blight <i>Sclerotinia</i> spp.	15 to 25			

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7-day interval. Use the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (25 ozs/A contains 0.395 lb boscalid, 0.200 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (50 ozs/A contains 0.790 lb boscalid, 0.400 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A (0.790 lb boscalid and 0.400 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.
- Minimum time from application to harvest (PHI) is 14 days.

\*\* Not registered for use in California.



**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Bulb vegetables [group 3-07]</b> Chive, fresh leaves Chive, Chinese, fresh leaves Daylily, bulb Elegans hosta Fritillaria, bulb Fritillaria, leaves Garlic, bulb Garlic, great-headed, bulb Garlic, serpent, bulb Kurrat Lady's leek Leek Leek, wild Lily, bulb Onion, Beltsville bunching Onion, bulb Onion, Chinese, bulb Onion, fresh Onion, green Onion, macrostem Onion, pearl Onion, potato, bulb Onion, tree, tops Onion, Welsh, tops Shallot, bulb Shallot, fresh leaves Cultivars, varieties and/or hybrids of these	Botrytis leaf blight <i>Botrytis</i> spp.	14.5 to 18.5	6*	111	7
	Botrytis neck rot** <i>Botrytis</i> spp.	10.5 to 18.5			
	Purple blotch and leaf blight <i>Alternaria porri</i>	18.5			
	<b>Suppression Only:</b> Downy mildew <i>Peronospora destructor</i>				

(continued)

**Table 3. Pristine® fungicide Crop-specific Requirements** *(continued)*

**Bulb vegetables [group 3-07]** *(continued)*

**Application Directions.** For control of neck rot, purple blotch and leaf blight, begin applications of **Pristine** prior to onset of disease development and continue on a 14-day interval. If application intervals shorter than 14 days are needed, rotate to another fungicide with a different mode of action. Use the higher rate when disease pressure is high.

Applications made to control purple blotch, leaf blight and stalk rot will also suppress downy mildew. If downy mildew occurs during a **Pristine** application for these diseases, immediately follow the **Pristine** application with a downy mildew fungicide with a different mode of action.

**For downy mildew,** rotate each application of **Pristine** with an application of a labeled fungicide with a different mode of action.

No restriction on livestock grazing or feeding.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (18.5 ozs/A contains 0.292 lb boscalid, 0.148 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (111 ozs/A contains 1.75 lbs boscalid, 0.888 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 18.5 ozs/A (0.292 lb boscalid, 0.148 lb pyraclostrobin).
- Maximum annual application rate is 111 ozs/A (1.75 lbs boscalid, 0.888 lb pyraclostrobin) per year.
- Minimum retreatment interval is 14 days.
- Minimum time from application to harvest (PHI) is 7 days.

\*\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Carrot	Alternaria leaf spot <i>Alternaria</i> spp.  Cercospora leaf spot <i>Cercospora</i> spp.  Powdery mildew <i>Erysiphe</i> spp.  <b>Suppression Only:</b>  Southern root rot <i>Sclerotium rolfsii</i>	8 to 10.5	6*	63	0

**Application Directions.** Begin applications of **Pristine** prior to onset of disease development and continue on a 7-day to 14-day interval. Use the higher rate and the shorter interval when disease pressure is high.

No restriction on livestock grazing or feeding for carrot culls.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (10.5 ozs/A contains 0.166 lb boscalid, 0.084 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (63 ozs/A contains 0.995 lb boscalid, 0.504 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 10.5 ozs/A (0.166 lb boscalid, 0.084 lb pyraclostrobin).
- Maximum annual application rate is 63 ozs/A (0.995 lb boscalid, 0.504 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Celery**</b> <b>Celery (Chinese)**</b>	Alternaria leaf spot <i>Alternaria</i> spp.	10 to 15	2*	50	0
	Anthracnose <i>Colletotrichum</i> spp.				
	Ascochyta leaf spot <i>Ascochyta</i> spp.				
	Cercospora leaf spot <i>Cercospora</i> spp.				
	Downy mildew <i>Bremia</i> spp., <i>Peronospora</i> spp.				
	Phoma <i>Phoma</i> spp.				
	Powdery mildew <i>Erysiphe</i> spp.				
	Rust <i>Puccinia</i> spp.				
	Septoria leaf spot <i>Septoria</i> spp.				
	White rust <i>Albugo</i> spp.				
Botrytis rot <i>Botrytis</i> spp.	25				
Sclerotinia rot and blight <i>Sclerotinia</i> spp.					

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7-day interval. Use the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (25 ozs/A contains 0.395 lb boscalid, 0.200 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (50 ozs/A contains 0.790 lb boscalid, 0.400 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A (0.790 lb boscalid, 0.400 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.

\*\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Celtuce**	Alternaria leaf spot <i>Alternaria</i> spp. Anthracnose <i>Colletotrichum</i> spp. Ascochyta leaf spot <i>Ascochyta</i> spp. Cercospora leaf spot <i>Cercospora</i> spp. Downy mildew <i>Bremia</i> spp., <i>Peronospora</i> spp. Phoma <i>Phoma</i> spp. Powdery mildew <i>Erysiphe</i> spp. Rust <i>Puccinia</i> spp. Septoria leaf spot <i>Septoria</i> spp. White rust <i>Albugo</i> spp.	10 to 15	2*	50	0
	Botrytis rot <i>Botrytis</i> spp. Sclerotinia rot and blight <i>Sclerotinia</i> spp.	15 to 25			

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7-day interval. Use the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (25 ozs/A contains 0.395 lb boscalid, 0.200 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (50 ozs/A contains 0.790 lb boscalid, 0.400 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A (0.790 lb boscalid, 0.400 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.

\*\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Citrus fruit [group 10-10]</b></p> <p>Australian desert lime                      Australian finger lime                      Australian round lime                      Brown River finger lime                      Calamondin                      Chironja                      Citron                      Citrus hybrids                      Grapefruit                      Japanese summer grapefruit                      Kumquat                      Lemon                      Lime                      Mediterranean mandarin                      Mount White lime                      New Guinea wild lime                      Orange, sour                      Orange, sweet                      Pummelo                      Russell River lime                      Satsuma mandarin                      Sweet lime                      Tachibana orange                      Tahiti lime                      Tangelo                      Tangerine (mandarin)                      Tangor                      Trifolate orange                      Uniq fruit                      Cultivars, varieties and/or hybrids of these</p>	<p>Alternaria brown spot  <i>Alternaria alternata</i>,  <i>Alternaria</i> spp.</p> <p>Citrus black spot*  <i>Guignardia citricarpa</i></p> <p>Greasy spot  <i>Mycosphaerella citri</i></p> <p>Melanose  <i>Diaporthe citri</i></p> <p>Scab  <i>Elsinoe fawcettii</i></p>	<p>16 to 18.5</p>	<p>4</p>	<p>74</p>	<p>0</p>

(continued)

**Table 3. Pristine® fungicide Crop-specific Requirements** *(continued)*

**Citrus fruit [group 10-10]** *(continued)*

**Application Directions.** Apply **Pristine** in a regularly scheduled protective fungicide program. Begin **Pristine** applications prior to infection and continue on a 10-day to 21-day interval. Use the higher rate and shorter interval when disease pressure is high.

Disease control from **Pristine** depends on disease pressure and various cultural practices that influence rind maturation and disease susceptibility. Improved disease performance may result when **Pristine** is used in a crop management program that minimizes rind overmaturity and rind damage.

No livestock feeding restrictions.

For aerial application to citrus fruit trees, use no less than 10 gallons of spray solution per acre.

**Resistance Management.** To limit development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with different modes of action.

**Restrictions and Limitations**

- Maximum single application rate is 18.5 ozs/A (0.292 lb boscalid, 0.148 lb pyraclostrobin).
- Maximum annual application rate is 74 ozs/A (1.17 lbs boscalid, 0.592 lb pyraclostrobin) per year.
- **DO NOT** make more than four (4) applications of **Pristine** per year.
- Minimum retreatment interval is 10 days.

\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Cotton**</b>	Alternaria leaf spot <i>Alternaria</i> spp. Anthracnose, boll rot <i>Glomerella</i> spp. Ascochyta blight, boll rot <i>Ascochyta</i> spp. Cercospora leaf spot <i>Cercospora</i> spp. Diplodia boll rot <i>Diplodia</i> spp. Hard lock, boll rot <i>Fusarium</i> spp.	12.5 to 25	2*	50	30

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7-day to 14-day interval. Use the higher rate and the shorter interval when disease pressure is high.

Feed containing commodities from cotton production and processing can be fed to livestock.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (25 ozs/A contains 0.395 lb boscalid, 0.200 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (50 ozs/A contains 0.790 lb boscalid, 0.400 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A (0.790 lb boscalid, 0.400 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.
- Minimum time from application to harvest (PHI) is 30 days.

\*\* Includes cottonseed tolerance.



**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Cucurbit vegetables [group 9]</b>                      Chayote                      Chinese waxgourd                      Citron melon                      Cucumber                      Gherkin                      Pumpkin                      Watermelon</p> <p><b>Edible Gourd</b>                      Chinese okra                      Cucuzza                      Hechima                      Hyotan</p> <p><b>Momordica spp.</b>                      Balsam apple                      Balsam pear                      Bitter melon                      Chinese cucumber</p> <p><b>Muskmelon</b>                      Cantaloupe                      Casaba                      Crenshaw melon                      Golden pershaw melon                      Honey balls                      Honeydew melon                      Mango melon                      Persian melon                      Pineapple melon                      Santa Claus melon                      Snake melon</p> <p><b>Summer Squash</b>                      Crookneck squash                      Scallop squash                      Straightneck squash                      Vegetable marrow                      Zucchini</p> <p><b>Winter Squash</b>                      Acorn squash                      Butternut squash                      Calabaza                      Hubbard squash                      Spaghetti squash</p> <p>Cultivars, varieties and/or hybrids of these</p>	<p>Alternaria blight <i>Alternaria cucumerina</i></p> <p>Cercospora leaf spot <i>Cercospora citrulina</i></p> <p>Downy mildew <i>Pseudoperonospora cubensis</i></p> <p>Gummy stem blight <i>Didymella bryoniae</i></p> <p>Powdery mildew <i>Erysiphe cichoracearum</i>, <i>Sphaerotheca fuliginea</i></p> <p>Anthracnose <i>Colletotrichum orbiculare</i></p>	<p>12.5 to 18.5</p> <p>18.5</p>	<p>4</p>	<p>74</p>	<p>0</p>

(continued)

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

**Cucurbit vegetables [group 9]** (continued)

**Application Directions.** Begin applications of **Pristine** prior to onset of disease development and continue on a 7-day to 14-day interval. Use the higher rate and the shorter interval when disease pressure is high.

Use the highest labeled rate for anthracnose.

**Tank Mixes with Adjuvants and Other Products.** BASF evaluations indicate that tank mixes of additives, adjuvants, and/or other products with **Pristine** may result in injury. This is particularly true for muskmelon crops including cantaloupe and honeydew. Users need to be aware of this, proceed with caution, and test for crop safety when tank mixing, as stated below.

Applications of additives, adjuvants, and/or other products that increase penetration may cause injury when mixed with **Pristine**. Injury potential from these kinds of tank mixes may decrease with lower rates of the tank mix partner. Users are advised to test for crop safety, as stated below.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives, adjuvants, and/or other products. Local environmental conditions also influence crop response and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing **Pristine** with other products.

To minimize the likelihood of crop injury, BASF advises testing **Pristine** in combination with additives, adjuvants, and/or other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

**Resistance Management.** To limit the potential of development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Restrictions and Limitations**

- Maximum single application rate is 18.5 ozs/A (0.292 lb boscalid, 0.148 lb pyraclostrobin).
- Maximum annual application rate is 74 ozs/A (1.17 lbs boscalid, 0.592 lb pyraclostrobin) per year.
- **DO NOT** make more than four (4) applications of **Pristine** per year.
- **DO NOT** tank mix **Pristine** with chlorpyrifos, dicofol, endosulfan, malathion, methomyl, potassium salts of fatty acids, or dicloran as crop injury may result.
- Minimum retreatment interval is 7 days.

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Dill seed	Phoma blight <i>Cercosporidium</i> spp.	18.5	2	37	0

**Application Directions.** Begin **Pristine** applications prior to the onset of disease development and repeat application 7 days later as needed or rotate to another fungicide with a different mode of action.

**Restrictions and Limitations**

- Maximum single application rate is 18.5 ozs/A (0.292 lb boscalid, 0.148 lb pyraclostrobin).
- Maximum annual application rate is 37 ozs/A (0.585 lb boscalid, 0.296 lb pyraclostrobin) per year.
- **DO NOT** make more than two (2) applications of **Pristine** per year.
- **DO NOT** apply by air.
- **DO NOT** apply through any type of irrigation system (chemigation).
- Minimum retreatment interval is 7 days.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Fennel (Florence)**</b>	Alternaria leaf spot <i>Alternaria</i> spp.  Anthracnose <i>Colletotrichum</i> spp.  Ascochyta leaf spot <i>Ascochyta</i> spp.  Cercospora leaf spot <i>Cercospora</i> spp.  Downy mildew <i>Bremia</i> spp., <i>Peronospora</i> spp.  Phoma <i>Phoma</i> spp.  Powdery mildew <i>Erysiphe</i> spp.  Rust <i>Puccinia</i> spp.  Septoria leaf spot <i>Septoria</i> spp.  White rust <i>Albugo</i> spp.	10 to 15	2*	50	0
	Botrytis rot <i>Botrytis</i> spp.  Sclerotinia rot and blight <i>Sclerotinia</i> spp.	15 to 25			

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7-day interval. Use the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (25 ozs/A contains 0.395 lb boscalid, 0.200 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (50 ozs/A contains 0.790 lb boscalid, 0.400 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A (0.790 lb boscalid, 0.400 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.

\*\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Fruiting vegetables [group 8-10]</b></p> <p>African eggplant                      Bell pepper                      Bush tomato                      Cocona                      Currant tomato                      Eggplant                      Garden huckleberry                      Goji berry                      Groundcherry                      Martynia                      Naranjilla                      Okra                      Pea eggplant                      Pepino                      Pepper (all varieties)                      Nonbell pepper                      Roselle                      Scarlet eggplant                      Sunberry                      Tomatillo                      Tree tomato</p> <p>Cultivars, varieties and/or hybrids of these</p>	<p>Black mold  <i>Alternaria alternata</i></p> <p>Early blight  <i>Alternaria solani</i></p>	<p>9.7</p> <p>or</p> <p>9.7 ozs per 100 gal of spray volume (dilute)*</p>	<p>6</p>	<p>58.2</p>	<p>0</p>
<p>Tomato</p> <p>Cultivars, varieties and/or hybrids</p>	<p>Anthracnose  <i>Colletotrichum</i> spp.</p> <p>Black mold  <i>Alternaria alternata</i></p> <p>Botrytis gray mold  <i>Botrytis cinerea</i></p> <p>Early blight  <i>Alternaria solani</i></p> <p>Late blight  <i>Phytophthora infestans</i></p> <p>Powdery mildew  <i>Leveillula taurica</i></p> <p>Septoria leaf spot  <i>Septoria lycopersici</i></p> <p>Target spot  <i>Corynespora cassiicola</i></p>	<p>12.5 to 25</p>	<p>5 at 12.5 ozs/A</p> <p>2 at 25 ozs/A</p>	<p>69</p>	

(continued)

**Table 3. Pristine® fungicide Crop-specific Requirements** *(continued)*

**Fruiting vegetables [group 8-10]** *(continued)*

**Application Directions.** Begin **Pristine** applications prior to disease development and continue on a 7-day to 14-day interval for anthracnose, black mold, botrytis gray mold, early blight, powdery mildew, Septoria leaf spot, and target spot. For control of late blight, begin applications prior to disease development; then follow each **Pristine** application with a labeled fungicide with a different mode of action 5 to 7 days later. Use the higher rate and shorter interval when disease pressure is high.

\* For applications based on dilute volume, spray plants to runoff. Apply a minimum of 20 gallons of spray volume per acre, and increase the spray volume as the plants grow. Spray proportional volume to the amount of plant tissue to be covered such that 100 gallons of spray per acre is used on mature plants.

**Use of Adjuvants.** Additive or adjuvant use may improve the performance of **Pristine** on fruiting vegetables. However, BASF evaluations also indicate that under some conditions (particularly high temperatures and/or high additive rates), **Pristine** application in combination with certain rates of silicone based or oil-containing (petroleum or crop) additives or adjuvants can cause injury.

BASF has not tested all varieties and cultivars with all possible tank mix combinations and rates of additives or adjuvants. Local environmental conditions also influence crop response and may not match those under which BASF has conducted testing. Physical incompatibility, reduced disease control, or crop injury may result from mixing **Pristine** with other products.

To the extent consistent with applicable law, the user assumes all risks associated with adding products to the **Pristine** spray solution. BASF cannot be held responsible for crop injury, reduced disease control or incompatibility because of additives, adjuvants or other products used in combination with **Pristine** (see **Conditions of Sale and Warranty**).

To minimize the likelihood of crop injury, test **Pristine** in combination with other products for crop safety on a small portion of the crop. However, environmental variability precludes direct and consistent projection of small area test results to future use.

Consult a BASF representative for more information concerning additives or adjuvants.

**Resistance Management.** To limit development of resistance, **DO NOT** make more than two (2) sequential **Pristine** applications before alternating to a labeled **non-Group 7** or **non-Group 11** fungicide with a different mode of action.

For additional resistance management information, see **Resistance Management** in the **Product Information** section.

**Restrictions and Limitations**

- **DO NOT** apply more than 58.2 ozs/A (0.920 lb boscalid, 0.466 lb pyraclostrobin) per year for fruiting vegetables, except Tomato. For Tomato, **DO NOT** apply more than 69 ozs/A (1.09 lbs boscalid, 0.552 lb pyraclostrobin) per year.
- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- **DO NOT** make more than six (6) applications of **Pristine** per year for fruiting vegetables, except Tomato. For Tomato, **DO NOT** make more than five (5) applications of **Pristine** per year at 12.5 ozs/A or two (2) applications of **Pristine** per year at 25 ozs/A.
- Minimum retreatment interval is 7 days.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Globe artichoke	Bud rot <i>Botrytis cinerea</i>  Ramularia leaf spot <i>Ramularia</i> spp.	18.5 to 23.0	3	69	0

**Application Directions. Dosage and frequency/timing of applications.** Begin applications of **Pristine** prior to onset of disease development and continue on a 7-day to 14-day interval. For artichoke bud rot, begin applications at the initiation of the bud protection phase when approximately 25% of the plants have bolted. Use the shorter interval and/or the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** exceed the specified number of applications of **Pristine** or other **Group 7** or **Group 11** fungicides per year. Adhere to the label instructions regarding the consecutive use of **Pristine** or other target site of action **Group 7** and **Group 11** fungicides that have a similar site of action on the same pathogens.

**Restrictions and Limitations**

- Maximum single application rate is 23 ozs/A (0.363 lb boscalid, 0.184 lb pyraclostrobin).
- Maximum annual application rate is 69 ozs/A (1.09 lbs boscalid, 0.552 lb pyraclostrobin) per year.
- **DO NOT** make more than three (3) applications of **Pristine** per year.
- **DO NOT** apply **Pristine** to artichokes as a tank mix with any other pesticide products (including fungicides, insecticides, herbicides), adjuvants, liquid fertilizers, nutrients, any other additives or anything other than water. Mix **Pristine** with water only for applications to artichokes.
- Minimum retreatment interval is 7 days.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Grape (except Concord or Noiret (NY73.0136.17) due to foliar injury.</b> It is possible that foliar injury could occur on related grape varieties. See comments in the <b>Application Directions</b> for more information.)</p>	<p>Angular leaf spot <i>Mycosphaerella angulata</i></p> <p>Anthracnose <i>Elsinoe ampelina</i></p> <p>Black rot <i>Guignardia bidwellii</i></p> <p>Downy mildew <i>Plasmopara viticola</i></p> <p>Leaf blight <i>Pseudocercospora vitis</i></p> <p>Phomopsis cane and leaf spot <i>Phomopsis viticola</i></p> <p>Powdery mildew <i>Uncinula necator</i></p> <p>Ripe rot <i>Colletotrichum gloeosporioides</i></p> <p><b>Aids in Control Only:</b></p> <p>Summer bunch rot (Sour rot) <i>Aspergillus</i> spp. and <i>Cladosporium</i> spp.</p> <p><b>Suppression Only:</b></p> <p>Botrytis gray mold <i>Botrytis cinerea</i></p>	8 to 12.5	5*	69	14
	<p>Botrytis gray mold <i>Botrytis cinerea</i></p>	18.5 to 23	3		

(continued)

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

**Grape** (continued)

**Application Directions. For powdery mildew control**, begin applications of **Pristine** as of bud break prior to onset of disease, using 8 ozs per acre on a 10-day to 14-day interval. Use 10 to 12.5 ozs per acre on a 14-day to 21-day interval.

**For black rot and downy mildew control**, begin applications of **Pristine** as of pre-bloom prior to onset of disease and continue applications on a 10-day to 14-day interval.

**For all other diseases listed except for Botrytis gray mold**, begin applications of **Pristine** prior to onset of disease and continue applications on a 10-day to 14-day interval. **Pristine** applied at rates of 8 to 12.5 ozs per acre for control of the listed diseases will also suppress Botrytis gray mold.

**For control of Botrytis gray mold**, apply 18.5 to 23 ozs per acre of **Pristine** prior to onset of disease development when conditions favor disease development during early bloom, bunch pre-closure and veraison up to 14 days before harvest. Continue applications on a 10-day to 14-day interval.

Use the higher rate and the shorter interval when disease pressure is high.

For aerial application to grape, use no less than 10 gallons of spray solution per acre.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than five (5) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per year. **DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (23 ozs/A contains 0.363 lb boscalid, 0.184 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (69 ozs/A contains 1.09 lbs boscalid, 0.552 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 23 ozs/A (0.363 lb boscalid, 0.184 lb pyraclostrobin).
- Maximum annual application rate is 69 ozs/A (1.09 lbs boscalid, 0.552 lb pyraclostrobin) per year.
- **DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours except when performing cane tying, cane turning, cane girdling or leaf pulling tasks. The REI is 5 days for treated grapes when conducting cane tying, cane turning, cane girdling or leaf pulling tasks. Notify workers of this prohibition.**
- **DO NOT** use on **Concord or Noiret (NY73.0136.17)** due to foliar injury. Possible foliar injury could occur to **Worden, Fredonia, Niagara, Steuben, Rougeon** or related grape varieties. Not all varieties have been thoroughly tested. Consult a BASF representative for more information concerning these sensitive grapes.
- Minimum retreatment interval is 10 days.
- Minimum time from application to harvest (PHI) is 14 days.



**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Herb [subgroup 19A]</b> Angelica Balm Basil Borage Burnet Catnip Chamomile Chervil (dried) Chive Chive, Chinese Clary Coriander (leaf) Costmary Culantro (leaf) Curry (leaf) Dillweed Horehound Hyssop Lavender Lemongrass Lovage (leaf) Marigold Marjoram Nasturtium Parsley (dried) Pennyroyal Rosemary Rue Sage Savory, summer and winter Sweet bay Tansy Tarragon Thyme Wintergreen Woodruff Wormwood	Phoma blight <i>Cercosporidium</i> spp.  Powdery mildew <i>Erysiphe</i> spp., <i>Sphaerotheca</i> spp.  Rust <i>Puccinia</i> spp.	18.5	2	37	0

**Application Directions.** Begin **Pristine** applications prior to the onset of disease development and repeat application 7 days later as needed or rotate to another fungicide with a different mode of action.

**Restrictions and Limitations**

- Maximum single application rate is 18.5 ozs/A (0.292 lb boscalid, 0.148 lb pyraclostrobin).
- Maximum annual application rate is 37 ozs/A (0.585 lb boscalid, 0.296 lb pyraclostrobin) per year.
- **DO NOT** make more than two (2) applications of **Pristine** per year.
- **DO NOT** apply by air.
- **DO NOT** apply through any type of irrigation system (chemigation).
- Minimum retreatment interval is 7 days.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application	Maximum Number of Ground Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Hops (Ground Application)</b>	Powdery mildew <i>Erysiphe cichoracearum</i> , <i>Sphaerotheca</i> spp.  Downy mildew <i>Pseudoperonospora humuli</i>	14 ozs per 100 gallons of dilute spray  <b>DO NOT</b> use more than 28 ozs per acre.	3  (2 if one aerial application is made)	84  (70 ozs/A if one aerial application is made)	14

**Application Directions.** Begin applications of **Pristine** prior to disease development and continue on a 10-day to 21-day interval. Use the shorter interval when disease pressure is high.

Application rates are based on 100 gallons of dilute spray applied to runoff. Adjust water volume to maintain thorough coverage. Use 25 to 50 gallons of dilute spray per acre prior to trellising and 100 to 200 gallons of dilute spray per acre thereafter. **DO NOT** use more than 200 gallons per acre of this mixture. If additional spray volume is needed for thorough coverage, use 28 ozs of **Pristine** per acre in the required spray volume.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

- **DO NOT** apply more than 84 ozs/A (1.33 lbs boscalid, 0.672 lb pyraclostrobin) per year. If one aerial application is made, **DO NOT** apply more than 70 ozs/A (1.11 lbs boscalid, 0.560 lb pyraclostrobin) per year.
- **DO NOT** make more than three (3) applications of **Pristine** per year (counting both ground and aerial applications).
- **DO NOT** use more than 200 gallons per acre of this mixture. If additional spray volume is needed for thorough coverage, use 28 ozs of **Pristine** per acre in the required spray volume.
- Maximum single application rate is 28 ozs/A (0.442 lb boscalid, 0.224 lb pyraclostrobin).
- Minimum retreatment interval is 10 days.
- Minimum time from application to harvest (PHI) is 14 days.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Aerial Product Use Rate per Application	Maximum Number of Aerial Applications per Year	Aerial Application Timing Growth Stage	Minimum Time from Aerial Application to Harvest (PHI) (days)
<b>Hops (Aerial Application)</b>	Powdery mildew <i>Erysiphe cichoracearum</i> , <i>Sphaerotheca humuli</i> , <i>Sphaerotheca macularis</i> , <i>Sphaerotheca</i> spp.	14 ozs/A as a tank mix with a myclobutanil fungicide product (see myclobutanil rate following)	1	Wire to 14 days preharvest	14

**Application Directions.** Aerial application may result in reduced control due to lack of canopy penetration and coverage. Use aerial application only when ground application is not possible.

Apply a preventive spray of **Pristine** at 14 ozs (0.221 lb boscalid, 0.112 lb pyraclostrobin) as a tank mix with a myclobutanil fungicide product at a rate **equivalent to 0.15 lb per acre of active ingredient** for resistance management.

Avoid applications under conditions when uniform coverage cannot be obtained or when spray drift may occur. Use a minimum of 10 gallons of water per acre when applying by air. Thorough coverage is essential.

Because complete coverage is important for effective disease control, **aerial application at low volumes may result in reduced control due to lack of canopy penetration and coverage.**

Mixing **Pristine** with surfactants or foliar fertilizers is not advised when applying by air. Similarly, adjuvants that enhance pesticide penetration may cause phytotoxicity when used with **Pristine** applied by air.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) aerial application of **Pristine** per year and include a myclobutanil product as a tank mix as described.

**Restrictions and Limitations**

- **DO NOT** apply more than 14 ozs/A (0.221 lb boscalid, 0.112 lb pyraclostrobin) per application
- **DO NOT** make more than one (1) application of **Pristine** per year.
- Minimum time from application to harvest (PHI) is 14 days.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Kohlrabi**	Alternaria leaf spot <i>Alternaria</i> spp.	10 to 15	2*	50	0
	Black leg <i>Phoma lingam</i>				
	Cercospora leaf spot <i>Cercospora</i> spp.				
	Powdery mildew <i>Erysiphe</i> spp.				
	Ring spot <i>Mycosphaerella</i> spp.				
	White leaf spot <i>Pseudocercospora</i> spp.				
	Anthracnose <i>Colletotrichum</i> spp.				
White rust <i>Albugo</i> spp.	15 to 25				
Downy mildew <i>Peronospora</i> spp.					
Gray mold <i>Botrytis</i> spp.					
Sclerotinia stem rot <i>Sclerotinia</i> spp.					
Rhizoctonia bottom rot <i>Rhizoctonia solani</i>					

**Application Directions.** Begin applications of **Pristine** prior to disease development and continue on a 7-day to 14-day interval. Use the higher rate and the shorter interval when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (25 ozs/A contains 0.395 lb boscalid, 0.200 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (50 ozs/A contains 0.790 lb boscalid and 0.400 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A(0.790 lb boscalid, 0.400 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.

\*\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Leaf petiole vegetables [subgroup 22B]**</b> Cardoon Fuki Rhubarb Udo Zuiki Cultivars, varieties and/or hybrids of these	Alternaria leaf spot <i>Alternaria</i> spp. Anthracnose <i>Colletotrichum</i> spp. Ascochyta leaf spot <i>Ascochyta</i> spp. Cercospora leaf spot <i>Cercospora</i> spp. Downy mildew <i>Bremia</i> spp., <i>Peronospora</i> spp. Phoma <i>Phoma</i> spp. Powdery mildew <i>Erysiphe</i> spp. Rust <i>Puccinia</i> spp. Septoria leaf spot <i>Septoria</i> spp. White rust <i>Albugo</i> spp.	10 to 15	2*	50	0
	Botrytis rot <i>Botrytis</i> spp. Sclerotinia rot and blight <i>Sclerotinia</i> spp.	15 to 25			

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7-day interval. Use the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (25 ozs/A contains 0.395 lb boscalid, 0.200 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (50 ozs/A contains 0.790 lb boscalid, 0.400 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A (0.790 lb boscalid, 0.400 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.

\*\* Not registered for use in California. See separate crop table for celery and Chinese celery use rates and application directions.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Leafy greens [subgroup 4-16A]**</b> Amaranth, Chinese Amaranth, leafy Aster, Indian Blackjack Cat's whiskers Cham-chwi Cham-na-mul Chervil, fresh leaves Chipilin Chrysanthemum, garland Cilantro, fresh leaves Corn salad Cosmos Dandelion, leaves Dang-gwi, leaves Dillweed Dock Dol-nam-mul Ebolo Endive Escarole Fameflower Feather cockscomb Good King Henry Huauzontle Jute, leaves Lettuce, bitter Lettuce, head Lettuce, leaf Orach Parsley, fresh leaves Plantain, buckhorn Primrose, English Purslane, garden Purslane, winter Radicchio Swiss chard Violet, Chinese, leaves Cultivars, varieties, and/or hybrids of these	Alternaria leaf spot <i>Alternaria</i> spp. Anthracnose <i>Colletotrichum</i> spp. Ascochyta leaf spot <i>Ascochyta</i> spp. Cercospora leaf spot <i>Cercospora</i> spp. Downy mildew <i>Peronospora</i> spp. Phoma <i>Phoma</i> spp. Powdery mildew <i>Erysiphe</i> spp., <i>Phyllactinia</i> spp., <i>Sphaerotheca</i> spp. Rust <i>Puccinia</i> spp. Septoria leaf spot <i>Septoria</i> spp. White rust <i>Albugo</i> spp.	10 to 15	2*	50	14
	Botrytis rot <i>Botrytis</i> spp. Sclerotinia rot and blight <i>Sclerotinia</i> spp.	15 to 25			

(continued)

**Table 3. Pristine® fungicide Crop-specific Requirements** *(continued)*

**Leafy greens [subgroup 4-16A]\*\*** *(continued)*

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7-day interval. Use the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (25 ozs/A contains 0.395 lb boscalid, 0.200 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (50 ozs/A contains 0.790 lb boscalid, 0.400 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A (0.790 lb boscalid, 0.400 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.
- Minimum time from application to harvest (PHI) is 14 days.

\*\* Not registered for use in California, except on radicchio for control of Sclerotinia rot and blight when applied at a rate of 25 ozs/A.

**Table 3. Pristine® fungicide Crop-specific Requirements** *(continued)*

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Oilseed [group 20]†* Flax	Pasmo <i>Septoria linicola</i>	16.7	2	33.4	21
Rapeseed (canola varieties only)	Blackleg <i>Leptosphaeria maculans</i>  Blackspot <i>Alternaria</i> spp.  Sclerotinia rot and blight <i>Sclerotinia</i> spp.				
Sunflower	Alternaria leaf spot <i>Alternaria</i> spp.  Cercospora leaf spot <i>Cercospora helianthi</i>  Downy mildew <i>Plasmopara halstedii</i>  Powdery mildew <i>Erysiphe cichoracearum</i>  Rust <i>Puccinia helianthi</i> , <i>Uromyces</i> spp.  Sclerotinia rot and blight <i>Sclerotinia</i> spp.  Septoria leaf spot <i>Septoria</i> spp.  White rust <i>Albugo tragopogonis</i>	24.5	2	49	21

*(continued)*



**Table 3. Pristine® fungicide Crop-specific Requirements** *(continued)*

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Oilseed [group 20]†*</b>  <i>(continued)</i>                      Borage                      Calendula                      Castor oil plant                      Chinese tallowtree                      Crambe                      Cuphea                      Echium                      Euphorbia                      Evening primrose                      Gold of pleasure                      Hare's ear mustard                      Jojoba                      Lesquerella                      Lunaria                      Meadowfoam                      Milkweed                      Mustard seed                      Niger seed                      Oil radish                      Poppy seed                      Rose hip                      Safflower                      Sesame                      Stokes aster                      Sweet rocket                      Tallowwood                      Tea oil plant                      Vernonia</p>	<p><i>Alternaria</i> spp.  <i>Septoria</i> spp.</p>	<p>16.7</p>	<p>2</p>	<p>33.4</p>	<p>21</p>

*(continued)*

**Table 3. Pristine® fungicide Crop-specific Requirements** *(continued)*

**Oilseed [group 20]†\*** *(continued)*

**Application Directions.** For optimal disease control, begin applications of **Pristine** prior to disease development and continue on a 7-day to 14-day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

**Rapeseed. For control of blackleg,** apply **Pristine** at 2- to 4-leaf stage. **For optimal control of blackspot,** apply **Pristine** at early pod development. A second application 7 to 10 days later may be made if disease persists or if weather conditions are favorable for disease development.

**Flax.** Apply **Pristine** at mid-flowering (7 to 10 days after flower initiation). Make a second application 7 to 10 days later if disease persists or if weather conditions are favorable for disease development.

† See separate crop table for cotton/cottonseed use rates and application directions.

**Pristine** may be used with adjuvants.

No livestock feeding restrictions.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

- **DO NOT** apply more than 33.4 ozs/A (0.528 lb boscalid, 0.267 lb pyraclostrobin) per year for oilseed crop, except sunflower. **DO NOT** apply more than 49 ozs/A (0.774 lb boscalid, 0.392 lb pyraclostrobin) per year for sunflower.
- Maximum single application rate is 24.5 ozs/A (0.381 lb boscalid, 0.196 lb pyraclostrobin).
- **DO NOT** make more than two (2) applications of **Pristine** per year.
- Minimum retreatment interval is 7 days.
- Minimum time from application to harvest (PHI) is 21 days.

\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Pea and bean, dried shelled, except soybean, subgroup 6C**</b></p> <p><b>Lupinus spp.</b> Grain lupin Sweet lupin White lupin White sweet lupin</p> <p><b>Phaseolus spp.</b> Field bean Kidney bean Lima bean (dry) Navy bean Pink bean Pinto bean Tepary bean</p> <p><b>Pisum spp.</b> Field pea</p>	<p>Alternaria leaf and pod spot <i>Alternaria</i> spp.</p> <p>Ascochyta blight <i>Ascochyta</i> spp., <i>Phoma exigua</i></p> <p>Cercospora leaf spot <i>Cercospora</i> spp.</p> <p>Downy mildew <i>Phytophthora nicotianae</i></p> <p>Mycosphaerella blight <i>Mycosphaerella</i> spp.</p> <p>Powdery mildew <i>Erysiphe polygoni</i></p> <p>Rust <i>Uromyces appendiculatus</i></p> <p>Septoria leaf spot <i>Septoria</i> spp.</p>	10 to 15	2*	50	21
<p><b>Vigna spp.</b> Adjuki bean Blackeyed pea Catjang Cowpea Crowder pea Moth bean Mung bean Rice bean Southern pea Urd bean</p>	<p>Anthracnose <i>Colletotrichum</i> spp.</p> <p>Botrytis gray mold <i>Botrytis cinerea</i></p> <p>White mold <i>Sclerotinia sclerotiorum</i></p>	15 to 25			

**Application Directions.** For optimal disease control, begin applications of **Pristine** prior to onset of disease development or at the beginning of flowering and repeat on a 5-day to 14-day interval if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high.

**Resistance Management. DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (25 ozs/A contains 0.395 lb boscalid, 0.200 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (50 ozs/A contains 0.790 lb boscalid, 0.400 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A (0.790 lb boscalid, 0.400 lb pyraclostrobin) per year.
- Minimum retreatment interval is 5 days.
- Minimum time from application to harvest (PHI) is 21 days.
- **DO NOT** feed treated pea commodities to livestock.

\*\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** *(continued)*

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Peanut**	Early leaf spot <i>Cercospora arachidicola</i>  Late leaf spot <i>Cercosporidium personatum</i>  Pepperspot <i>Leptosphaerulina crassiasca</i>  Rust <i>Puccinia arachidis</i>  Web blotch <i>Phoma arachidicola</i>	12.5 to 18.5	3*	84	14
	Rhizoctonia limb rot, peg rot, and pod rot <i>Rhizoctonia solani</i>  Sclerotinia blight <i>Sclerotinia minor</i>  Sclerotium rot, Southern blight, Southern stem rot, and White mold <i>Sclerotium rolfsii</i>	18.5 to 28			

*(continued)*

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

**Peanut\*\*** (continued)

**Application Directions.** For control of **pepperspot, rust, web blotch, early and late leaf spot**, begin applications of **Pristine** prior to onset of disease development and continue on a 14-day interval.

For control of **Rhizoctonia and Sclerotium rot**, begin applications of **Pristine** prior to onset of disease development and continue on a 14-day interval.

For control of **Sclerotinia blight**, begin applications of **Pristine** prior to onset of disease development or 45 to 60 days after planting. Make a second application 14 to 21 days later.

Use the higher rate and/or shorter spray interval when disease pressure is high or in fields with a history of disease.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (28 ozs/A contains 0.442 lb boscalid, 0.224 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (84 ozs/A contains 1.33 lbs boscalid, 0.672 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 28 ozs/A (0.442 lb boscalid, 0.224 lb pyraclostrobin).
- Maximum annual application rate is 84 ozs/A (1.33 lbs boscalid, 0.672 lb pyraclostrobin) per year.
- Use of **Pristine** with silicone-based adjuvants may cause crop injury.
- **DO NOT** apply by air.
- **DO NOT** feed treated peanut hay to livestock.
- **DO NOT** graze livestock or harvest for forage use.
- Minimum retreatment interval is 14 days.
- Minimum time from application to harvest (PHI) is 14 days.

\*\* Not registered for use in California.

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Persimmon	Cercospora leaf spot <i>Cercospora</i> spp.	18.5 to 23.0	3	69	0

**Application Directions.** Dosage and frequency/timing of applications. Begin applications of **Pristine** prior to the onset of disease development and continue on a 7-day to 14-day interval. Use the shorter interval and/or higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** exceed the specified number of applications of **Pristine** or other **Group 7** or **Group 11** fungicides per year. Adhere to the label instructions regarding the consecutive use of **Pristine** or other target site of action **Group 7** and **Group 11** fungicides that have a similar site of action on the same pathogens.

**Restrictions and Limitations**

- Maximum single application rate is 23 ozs/A (0.363 lb boscalid, 0.184 lb pyraclostrobin).
- Maximum annual application rate is 69 ozs/A (1.09 lbs boscalid, 0.552 lb pyraclostrobin) per year.
- **DO NOT** make more than three (3) applications of **Pristine** per year.
- Minimum retreatment interval is 7 days.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Pome fruit [group 11-10]</b></p> <p>Apple Azarole Crabapple Loquat Mayhaw Medlar Pear Pear, Asian Pear, Oriental Quince Quince, Chinese Quince, Japanese Tejocote Cultivars, varieties and/or hybrids of these</p>	<p>Alternaria blotch <i>Alternaria mali</i></p> <p>Apple scab <i>Venturia inaequalis</i></p> <p>Bitter rot <i>Colletotrichum</i> spp.</p> <p>Black rot/Frogeye leaf spot <i>Botryosphaeria obtusa</i></p> <p>Blue mold* <i>Penicillium</i> spp.</p> <p>Brooks spot <i>Mycosphaerella pomi</i></p> <p>Flyspeck <i>Zygophiala jamaicensis</i></p> <p>Gray mold* <i>Botrytis</i> spp.</p> <p>Pear scab <i>Venturia pirina</i></p> <p>Powdery mildew <i>Podosphaera leucotricha</i></p> <p>Sooty blotch (disease complex)</p> <p>White rot <i>Botryosphaeria dothidea</i></p> <p><b>Suppression Only:</b></p> <p>Cedar apple rust <i>Gymnosporangium juniperi-virginianae</i></p> <p>Quince rust <i>Gymnosporangium clavipes</i></p>	14.5 to 18.5	4**	74	0

(continued)

**Table 3. Pristine® fungicide Crop-specific Requirements** *(continued)*

**Pome fruit [group 11-10]** *(continued)*

**Application Directions for scab, powdery mildew, frog-eye leaf spot and rust.** Begin applications of **Pristine** prior to disease development and continue on a 7-day to 10-day interval.

**Application Directions for blue mold, gray mold, sooty blotch, flyspeck, white rot, black rot, bitter rot and Alternaria blotch.** Begin applications of **Pristine** prior to disease development and continue on a 7-day to 14-day interval.

Use the higher rate and shorter interval when disease pressure is high.

No restriction on livestock grazing or feeding for pome fruit feed items.

For aerial application to pome fruit trees, use no less than 10 gallons of spray solution per acre.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than four (4) applications of **Pristine** or other **Group 7** or **Group 11** fungicides per year. **DO NOT** make more than two (2) sequential applications of **Pristine** before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\*\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (18.5 ozs/A contains 0.292 lb boscalid, 0.148 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (74 ozs/A contains 1.17 lbs boscalid, 0.592 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 18.5 ozs/A (0.292 lb boscalid, 0.148 lb pyraclostrobin).
- Maximum annual application rate is 74 ozs/A (1.17 lbs boscalid, 0.592 lb pyraclostrobin) per year.
- For pears, **DO NOT** use **Pristine** with a horticultural mineral oil as crop response to foliage and/or fruit can occur under certain conditions.
- Minimum retreatment interval is 7 days.

\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Root vegetables (except sugar beet) [subgroup 1B]**</b></p> <p>Beet, garden                      Burdock, edible                      Celeriac                      Chervil, turnip-rooted                      Chicory                      Ginseng                      Horseradish                      Parsley, turnip-rooted                      Parsnip                      Radish                      Radish, oriental                      Rutabaga                      Salsify                      Salsify, black                      Salsify, Spanish                      Skirret                      Turnip</p>	<p>Alternaria leaf spot  <i>Alternaria</i> spp.</p> <p>Cercospora leaf spot  <i>Cercospora</i> spp.</p> <p>Powdery mildew  <i>Erysiphe</i> spp.</p> <p><b>Suppression Only:</b></p> <p>Southern root rot  <i>Sclerotium rolfsii</i></p>	8 to 10.5	6*	63	0

**Application Directions.** Begin applications of **Pristine** prior to onset of disease development and continue on a 7-day to 14-day interval. Use the higher rate and the shorter interval when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one application.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (10.5 ozs/A contains 0.166 lb boscalid, 0.084 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (63 ozs/A contains 0.995 lb boscalid, 0.504 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 10.5 ozs/A (0.166 lb boscalid, 0.084 lb pyraclostrobin).
- Maximum annual application rate is 63 ozs/A (0.995 lb boscalid, 0.504 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.

\*\* Not registered for use in California.



**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
Soybean**	Alternaria leaf spot <i>Alternaria</i> spp.	8 to 16	2*	32	21
	Anthracnose <i>Colletotrichum truncatum</i>				
	Brown spot <i>Septoria glycines</i>				
	Cercospora blight <i>Cercospora kikuchii</i>				
	Frogeye leaf spot <i>Cercospora sojina</i>				
	Pod and stem blight <i>Diaporthe phaseolorum</i>				
	Rhizoctonia aerial blight <i>Rhizoctonia solani</i>				
Asian soybean rust <i>Phakopsora pachyrhizi</i>	12.5 to 16				
Southern blight <i>Sclerotium rolfsii</i>	16				
White mold <i>Sclerotinia sclerotiorum</i>					

**Application Directions.** For optimal disease control, apply **Pristine** at early flowering (R1 to R3 growth stage) or prior to disease development, whichever is earlier. Make a second application 7 to 21 days later if monitoring shows disease development or if conditions are conducive for disease infection. Use the higher labeled rate and shorter interval when disease pressure is high.

**Pristine** may be applied with adjuvants.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (16 ozs/A contains 0.253 lb boscalid, 0.128 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (32 ozs/A contains 0.506 lb boscalid, 0.256 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 16 ozs/A (0.253 lb boscalid, 0.128 lb pyraclostrobin).
- Maximum annual application rate is 32 ozs/A (0.506 lb boscalid, 0.256 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.
- **DO NOT** use soybean forage as feed earlier than 14 days after last application.
- **DO NOT** use soybean hay as feed earlier than 21 days after last application.
- Minimum time from application to harvest (PHI) is 21 days.

\*\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Spinach</b> <b>Spinach (Malabar, New Zealand, and tanier)</b>	Alternaria leaf spot <i>Alternaria</i> spp.	10 to 15	2*	50	14
	Anthracnose <i>Colletotrichum</i> spp.				
	Ascochyta leaf spot <i>Ascochyta</i> spp.				
	Cercospora leaf spot <i>Cercospora</i> spp.				
	Downy mildew <i>Peronospora</i> spp.				
	Phoma <i>Phoma</i> spp.				
	Powdery mildew <i>Erysiphe</i> spp., <i>Phyllactinia</i> spp., <i>Sphaerotheca</i> spp.				
	Rust <i>Puccinia</i> spp.				
	Septoria leaf spot <i>Septoria</i> spp.				
	White rust <i>Albugo</i> spp.				
Botrytis rot <i>Botrytis</i> spp.	25				
Sclerotinia rot and blight <i>Sclerotinia</i> spp.					

**Application Directions.** Begin applications of **Pristine** prior to the onset of disease development and continue on a 7-day interval. Use the higher rate when disease pressure is high.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than one (1) application of **Pristine** before alternating to a labeled fungicide with a different mode of action for at least one (1) application.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (25 ozs/A contains 0.395 lb boscalid, 0.200 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (50 ozs/A contains 0.790 lb boscalid, 0.400 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A (0.790 lb boscalid, 0.400 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.
- Minimum time from application to harvest (PHI) is 14 days.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Stone fruit [group 12-12]</b> Apricot Apricot, Japanese Capulin Cherry, black Cherry, Nanking Cherry, sweet Cherry, tart Jujube, Chinese Nectarine Peach Plum Plum, American Plum, beach Plum, Canada Plum, cherry Plum, Chickasaw Plum, Damson Plum, Japanese Plum, Klamath Plum, prune Plumcot Sloe Cultivars, varieties, and/or hybrids of these	Alternaria leaf spot <i>Alternaria</i> spp. Anthracnose <i>Colletotrichum</i> spp. Blossom blight <i>Monilinia</i> spp. Brown rot <i>Monilinia</i> spp. Leaf spot <i>Blumeriella jaapii</i> Powdery mildew <i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp. Ripe fruit rot <i>Botrytis cinerea</i> , <i>Monilinia fructicola</i> , <i>Monilinia laxa</i> , <i>Rhizopus</i> spp. Rust <i>Tranzschelia discolor</i> Scab <i>Cladosporium carpophilum</i> Shothole <i>Wilsonomyces carpophilus</i>	10.5 to 14.5	5*	72.5	0
Nectarine Peach	<b>Suppression Only:</b> Leaf curl** <i>Taphrina deformans</i>				

**Application Directions.** Begin application of **Pristine** at pink bud or prior to onset of disease development and continue on a 7-day to 14-day interval. Use the shorter interval and/or the higher rate when disease pressure is high.

For aerial application to stone fruit trees, use no less than 10 gallons of spray solution per acre.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (14.5 ozs/A contains 0.229 lb boscalid, 0.116 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (72.5 ozs/A contains 1.15 lbs boscalid, 0.580 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 14.5 ozs/A (0.229 lb boscalid, 0.116 lb pyraclostrobin.)
- Maximum annual application rate is 72.5 ozs/A (1.15 lbs boscalid, 0.580 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.

\*\* Not registered for use in California.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Strawberry</b>	Anthracnose <i>Colletotrichum</i> spp.  Botrytis gray mold <i>Botrytis cinerea</i>  Leaf spot <i>Mycosphaerella fragariae</i>  Powdery mildew <i>Sphaerotheca macularis</i>	18.5 to 23	5*	115	0

**Application Directions.** Begin applications of **Pristine** no later than 10% bloom, or prior to disease development, and continue on a 7-day to 14-day interval. Use the higher rate and the shorter interval when disease pressure is high.

It is impossible for BASF to test strawberry sensitivity to **Pristine** under all environments and all potential product mixture combinations. Local conditions can also influence crop response and may not match those under which BASF has conducted testing. Proceed with caution with regard to **Pristine** use, particularly in tank mixes and/or adjuvant combinations on strawberry crops. To reduce the risk of strawberry crop injury, BASF advises testing **Pristine** or **Pristine** tank mixtures on a small portion of the crop before broad scale use. To the extent consistent with applicable law, the user assumes all risks associated with adding products to the **Pristine** spray solution. Refer also to the **Conditions of Sale and Warranty** section of this label.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (23 ozs/A contains 0.363 lb boscalid, 0.184 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (115 ozs/A contains 1.82 lbs boscalid, 0.920 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 23 ozs/A (0.363 lb boscalid, 0.184 lb pyraclostrobin).
- Maximum annual application rate is 115 ozs/A (1.82 lbs boscalid, 0.920 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Tree nut [group 14-12]</b></p> <p>African nut-tree Almond Beechnut Brazil nut Brazilian pine Bunya Bur oak Butternut Cajou nut Candlenut Cashew Chestnut Chinquapin Coconut Coquito nut Dika nut Ginkgo Guiana chestnut Hazelnut (Filbert) Heartnut Hickory nut Japanese horse-chestnut Macadamia nut Mongongo nut Monkey-pot Monkey puzzle nut Okari nut Pachira nut Peach palm nut Pecan Pequi Pili nut Pine nut Pistachio Sapucaia nut Tropical almond Walnut, black Walnut, English Yellowhorn</p> <p>Cultivars, varieties, and/or hybrids of these</p>	<p>Alternaria leaf spot <i>Alternaria</i> spp.</p> <p>Anthracnose <i>Colletotrichum</i> spp., <i>Marossonina juglandis</i></p> <p>Blossom blight <i>Monilinia</i> spp.</p> <p>Botrytis blossom and shoot blight/Green fruit rot <i>Botrytis cinerea</i></p> <p>Eastern filbert blight <i>Anisogramma anomala</i></p> <p>Leaf rust <i>Tranzschelia discolor</i></p> <p>Panicle and shoot blight <i>Botryosphaeria</i> spp.</p> <p>Scab <i>Cladosporium</i> spp.</p> <p>Shothole <i>Wilsonomyces carpophilus</i></p>	<p>10.5 to 14.5</p>	<p>4*</p>	<p>58</p>	<p>14</p> <p><b>(for almond - 25 days)</b></p>

(continued)

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

**Tree nut [group 14-12]** (continued)

**Application Directions.** In **almond**, begin applications of **Pristine** at pink bud and continue on a 7-day to 14-day interval up to 25 days before harvest. In **filbert**, begin applications at budswell to budbreak, prior to infection and onset of disease development. Continue on a 7-day to 14-day interval to cover and protect new growth. In **pecan**, begin applications of **Pristine** prior to onset of disease development and continue on a 7-day to 21-day interval for the control of scab. In **pistachio**, begin applications prior to the onset of disease development and continue on a 10-day to 30-day interval. **For all other crops listed above**, apply **Pristine** prior to disease development and continue on a 7-day to 28-day interval. In all cases, use the shorter interval when shoot growth is very rapid.

Use the shorter interval and/or the higher rate when disease pressure is high.

No restriction on livestock feeding for almond hulls.

For aerial application to tree nuts, use no less than 10 gallons of spray solution per acre.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

\* **DO NOT** make more than the Maximum Number of Applications per Year for applications made at the maximum Product Use Rate per Application (14.5 ozs/A contains 0.229 lb boscalid, 0.116 lb pyraclostrobin). Additional applications per year are permitted when a lower Product Use Rate per Application is used, as long as the Maximum Product Rate per Year (58 ozs/A contains 0.916 lb boscalid, 0.464 lb pyraclostrobin) is not exceeded.

- Maximum single application rate is 14.5 ozs/A (0.229 lb boscalid, 0.116 lb pyraclostrobin).
- Maximum annual application rate is 58 ozs/A (0.916 lb boscalid, 0.464 lb pyraclostrobin) per year.
- Minimum retreatment interval is 7 days.
- Minimum time from application to harvest (PHI) is 14 days. **(for almond - 25 days)**

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<p><b>Tropical fruits (listed)</b></p> <p>Avocado                      Black sapote                      Canistel                      Mamey sapote                      Mango                      Papaya                      Sapodilla                      Star apple</p>	<p>Anthracnose  <i>Colletotrichum gloeosporioides</i></p> <p>Black spot  <i>Alternaria alternata</i>,  <i>Asperisporium caricae</i>,  <i>Cercospora papayae</i></p> <p>Dry rot  <i>Mycosphaerella</i> spp.</p> <p>Powdery mildew  <i>Erysiphe</i> spp., <i>Oidium</i> spp.</p> <p>Pseudocercospora spot/blotch  <i>Cercospora</i> spp.,  <i>Pseudocercospora purpurea</i></p> <p>Scab  <i>Elsinoe mangiferae</i>,  <i>Sphaceloma perseae</i></p>	18.5	2	37	0

**Application Directions.** Begin application of **Pristine** prior to the onset of disease development and repeat application 7 days later, as needed, or alternate with another labeled fungicide having a different mode of action.

**Resistance Management.** To limit the potential for development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

- Maximum single application rate is 18.5 ozs/A (0.292 lb boscalid, 0.148 lb pyraclostrobin).
- Maximum annual application rate is 37 ozs/A (0.585 lb boscalid, 0.296 lb pyraclostrobin) per year.
- **DO NOT** make more than two (2) applications of **Pristine** per year.
- Minimum retreatment interval is 7 days.
- **DO NOT** apply **Pristine** as a tank mix with any pesticide product (includes any fungicide, herbicide, or insecticide), fertilizer, nutrient or additive other than nonionic surfactant (NIS). NIS rate not to exceed 11 ozs/100 gallons (0.08% v/v).

**Table 3. Pristine® fungicide Crop-specific Requirements** (continued)

Crop	Target Disease	Product Use Rate per Application (ozs/A)	Maximum Number of Applications per Year	Maximum Product Rate per Year (ozs/A)	Minimum Time from Application to Harvest (PHI) (days)
<b>Turnip greens</b>	Alternaria leaf spot and blight <i>Alternaria</i> spp. Anthracnose <i>Colletotrichum</i> spp. Cercospora leaf spot <i>Cercospora brassicicola</i> Downy mildew <i>Peronospora parasitica</i> Gray mold <i>Botrytis cinerea</i> Powdery mildew <i>Erysiphe polygoni</i> Rhizoctonia stem rot and Bottom rot <i>Rhizoctonia solani</i> Ring spot <i>Mycosphaerella brassicicola</i> Sclerotinia stem rot <i>Sclerotinia minor</i> , <i>S. sclerotiorum</i> Southern blight <i>Sclerotium rolfsii</i> White rust <i>Albugo candida</i>	18.8 to 25	2	50	14

**Application Directions.** Begin **Pristine** applications prior to disease development and continue on a 7-day to 10-day interval. Use the higher rate and shorter interval when disease pressure is high.

**Resistance Management.** To limit development of resistance, **DO NOT** make more than two (2) sequential applications of **Pristine** or other **Group 7** or **Group 11** fungicides before alternating to a labeled fungicide with a different mode of action.

**Restrictions and Limitations**

- Maximum single application rate is 25 ozs/A (0.395 lb boscalid, 0.200 lb pyraclostrobin).
- Maximum annual application rate is 50 ozs/A (0.790 lb boscalid, 0.400 lb pyraclostrobin) per year.
- **DO NOT** make more than two (2) applications of **Pristine** per year.
- Minimum retreatment interval is 7 days.
- Minimum time from application to harvest (PHI) is 14 days.
- **DO NOT** apply by air.
- **DO NOT** apply through any type of irrigation system (chemigation).



## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Cabrio, Endura, Headline and Pristine are registered trademarks of BASF.*

© 2021 BASF Corporation  
All rights reserved.

007969-00199.20200914c.**NVA 2020-04-156-0163**  
Supersedes: NVA 2017-04-156-0178

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709

  
We create chemistry