

7969-207

08/22/2007

1/13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

AUG 22 2007

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Dave G. Bolin
BASF Corporation
Agricultural Products
P.O. Box 13528
Research Triangle Park, NC 27709-3528

Dear Mr. Bolin:
Subject: Regent® 4SC Insecticide
EPA Reg. No. 7969-207
Submission D57139 dated 5/17/05

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you submit the confirmatory data specified in Attachment A and make the label changes listed below. One copy of the finished labeling must be submitted prior to releasing the product for shipment.

Under the section entitled General Precautions and Instructions, insert a subheading entitled "Use Restrictions". Under this subheading place the following statements which apply to both corn and potato use:

- Do not apply this product through any kind of irrigation system.
- Do not harvest within 90 days of application.
- Do not plant root, leafy or legume vegetables within four months following application.
- Do not plant wheat or triticale within two months following application.
- Do not plant any other rotational crops (except field corn, potato, leafy vegetables, root vegetables, legumes, wheat or triticale) within 12 months following application.
- Observe the following precautions when applying in the vicinity of aquatic areas:
 - Do not apply within 20 yards (in New York State, do not apply within 100 feet or 33 yards) of lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, commercial aquaculture facilities or other bodies of water that convey water to these areas. Protection of aquatic areas may be enhanced by maintaining all or a portion of this buffer in vegetative cover.
- Do not use this product if the corn or potato is planted by hand.

Add the following restriction to the section on potato use:
Do not apply to sweet potatoes.

Move the statement "Do not contaminate water food or feed by storage or disposal" so that it appears immediate below the Storage and Disposal heading.

Change the subheading "storage" to read "pesticide storage."

On page 3, revise the following statement "No label dosage rates should be exceeded" to read "Do not exceed any label dosage rates."

Delete the heading "Pesticide disposal" the second time it appears.

Move the compatibility statements to the section on spray preparation.

Review the application instructions on page 4, which presently are formatted to apply only to corn, for applicability to potato, and if they do apply to potato, revise the label to make clear they also apply to potato.

Revise the last sentence of the instructions appearing in the application method column to read as follows:

Do not use T-banding over the top of a closed furrow.

Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have questions, please call Ann Sibold at 703 305-6502.

Sincerely,



Richard Gebken
Product Manager 10
Insecticide Branch
Registration Division (7505P)

Data Required as Condition of Registration

Group C—Subchronic Toxicity Test Guidelines

Guideline	Title	Due Date
870.3465	Inhalation 28 day	June 8, 2009

Comment: Inhalation 28 day toxicity study in the rat. This study is required to further characterize the inhalation risk for use in the risk assessment of fipronil. The protocol for the existing 90-day inhalation toxicity study (OPPTS Guideline 870.3465) should be followed with the exposure (treatment) ending after 28 days, instead of 90 days.

RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO AQUATIC INVERTEBRATES
For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



REGENT® 4 SC

INSECTICIDE

For In-Furrow Use on Corn and Potatoes

Shake Well Before Use

ACTIVE INGREDIENT:

Fipronil: 5-amino-1-(2,6-dichloro-4-(trifluoromethyl) phenyl)-4-((1*R,S*)-(trifluoromethyl)sulfinyl)-1-*H*-pyrazole-3-carbonitrile 39.4%

OTHER INGREDIENTS: 60.6%

TOTAL: 100.0%

One gallon Regent® 4 SC insecticide contains 4 lbs fipronil

EPA Reg No. 7969-207

EPA Est. No. _____

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

In case of an emergency endangering life or property involving this product
call, day or night, 1-800-832-HELP (4357)

See inside booklet for complete **First Aid, Precautionary Statements, Directions For Use, and Conditions of Sale and Warranty**

ACCEPTED
with COMMENTS
In EPA Letter Dated:
AUG 22 2007

NET CONTENTS: _____

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7969-207

5/13

FIRST AID

- If swallowed**
 - Immediately call a poison control center or doctor for treatment advice.
 - **DO NOT** induce vomiting unless told to do so by a poison control center or doctor.
 - Have person sip a glass of water if able to swallow.
 - **DO NOT** give anything by mouth to an unconscious person.
- If on skin or clothing**
 - Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- If inhaled**
 - Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
 - Call a poison control center or doctor for further treatment advice.
- If in eyes**
 - Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
 - Call a poison control center or doctor for treatment advice.

For MEDICAL Emergencies Call 24 hours a day 1-800-832-HELP (4357). Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: There is no specific antidote. All treatment should be based on observed signs and symptoms of distress in the patient. Overexposure to materials other than this product may have occurred.

GENERAL: Contact a physician immediately in all cases of suspected poisoning. Transport patient to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

SYMPTOMS OF POISONING: In severe cases of overexposure by oral ingestion, lethargy, muscle tremors, and in extreme cases, possibly convulsions, may occur.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

MAY BE FATAL IF SWALLOWED, ABSORBED THROUGH SKIN, OR INHALED. CAUSES EYE IRRITATION. DO NOT get on skin or breathe spray mist. **DO NOT** get in eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, and using tobacco. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)
Applicators, mixer/loaders and other handlers must wear:

- Coveralls over short-sleeved shirts and short pants
- Chemical-resistant (such as nitrile or butyl) gloves
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Protective eyewear
- Respirator with an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N,R,P or HE prefilter
- Chemical-resistant apron when cleaning equipment or mixing and loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's

concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS


Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to birds, fish, and aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

DO NOT contaminate water when disposing of equipment washwaters or rinsate.



6/13

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **0 hours**.

STORAGE AND DISPOSAL

STORAGE

DO NOT store in or around the home. Store unused product in original container only, in a cool, dry area out of reach of children and animals. Store at temperatures above 32° F. If exposed to freezing temperatures, store at temperatures above freezing for 24 hours or until completely thawed. Shake well before using.

PESTICIDE DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IN CASE OF SPILL

If container is damaged, before cleaning up, put on personal protective equipment.

GENERAL PRECAUTIONS AND INSTRUCTIONS


SPRAY PREPARATION

Remove oil, rust, scale, pesticide residues and foreign matter from sprayer and strainer. Flush with clean water. Use 50 mesh screens or equivalent in spray system. Applicator tanks must be equipped with a sparge tube agitator, unless the alternative agitator designs can adequately cleanse the bottom of the tank and provide sufficient agitation required for product mixing. Pump approximately 10% of the tank capacity per minute back into the tank through the agitator.

To prepare for spraying, fill tank to approximately 1/2 the needed volume of water or liquid fertilizer and start agitation. Open product container(s) and pour contents into tank. Agitate to ensure thorough mixing for a minimum of 2 minutes or until product is thoroughly dispersed. Agitate product mixtures continuously during use. Avoid allowing tank contents to remain without agitation for periods greater than 1 hour. When tank contents have remained without agitation for 1 hour or longer, agitate contents for at least 10 minutes before starting application. Avoid allowing large quantities of product mixtures (greater than 5 gallons) to be stored in a non-agitated tank for extended periods of time, such as 12 hours or greater.

Regent® 4 SC insecticide can be tank mixed in accordance with the most restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product whose label contains a prohibition against such mixing.

If tank mixes with other products are to be used, evaluate tank mix compatibility with **Regent 4 SC** in a small container before mixing larger quantities in the application tank.



USE DIRECTIONS FOR CC

For use on conventional and conservation tillage field corn.

PESTS CONTROLLED

Crop	Pests Controlled	Pests Suppressed
Field Corn (30" row spacing or higher)	Northern, Western, Mexican, and Southern Corn Rootworm larvae European Corn Borer (First Brood) Wireworms Common Stalk Borer Seedcorn Maggots Seedcorn Beetles Billbugs Chinch Bugs Grubs Grape Colaspis Thrips	Grasshoppers
Field Corn (15" to less than 30" row spacing)	Wireworms Seedcorn Maggots	Grubs Seedcorn Beetles

- **DO NOT** apply to sweet corn or popcorn.
- **DO NOT** use this product with corn seeds that have been treated with a product containing fipronil, such as **Regent® TS insecticide**.
- **DO NOT** harvest within 90 days of application.
- **DO NOT** plant root, leafy, or legume vegetables within four months following application.
- **DO NOT** plant wheat or triticale within two months following application.
- **DO NOT** plant any other rotational crops (except field corn, potato, leafy vegetables, root vegetables, legumes, wheat, or triticale) within 12 months following application.

APPLICATION

Make one in-furrow application at planting time only.

DO NOT apply this product through any kind of irrigation system.

Apply at planting time as a solid stream using a microtube or other suitable metering orifice that is directed into the open seed furrow. Dilute **Regent® 4 SC insecticide** in a minimum of 1.0 gallon of water or liquid fertilizer per acre. If flat-fan nozzles are used for in-furrow application, the nozzles should be aligned with the row to direct the spray into the open furrow.

Regent 4 SC can be directly injected into the suction or pressure line(s) of a liquid spray system provided that the product is uniformly dispersed into the water or liquid fertilizer. Predilution of **Regent 4 SC** in a minimum of 0.5 gallons of water or liquid fertilizer per acre, prior to injection, is necessary to improve metering accuracy. The mix tank of **Regent 4 SC** plus water or liquid fertilizer must be agitated continuously. After completion of the direct injection in the mixing process, the final volume delivered to the furrow should have a minimum volume of 1.0 gallon per acre of water, liquid fertilizer or the combination of the two.

Observe the following precautions when applying in the vicinity of aquatic areas: **DO NOT** apply within 20 yards (in New York State, **DO NOT** apply within 100 feet or 33 yards) of lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, commercial aquaculture facilities or other bodies of water that convey water to these areas. Protection of aquatic areas may be enhanced by maintaining all or a portion of this buffer in vegetative cover.

8/13

Use the following chart to select the correct rate of **Regent® 4 SC insecticide** per 1,000 row feet and rate per acre:

Row Spacing (inch)	Rate/Acre (fluid ounces/acre)	Row Feet/Acre (feet)	Rate/1,000 row feet (fluid ounces/1,000 row feet)
40	3.12	13,068	0.24
38	3.29	13,756	0.24
36	3.47	14,520	0.24
30	4.16	17,424	0.24
22	4.16	23,760	0.18
20	4.16	26,133	0.16
15	4.16	34,848	0.12

- **DO NOT** apply 0.24 fluid ounces/1,000 row ft on row spacing less than 30 inches.
- Adjust rate per 1,000 row feet so that you **DO NOT** apply more than 0.13 lb. ai/acre or 4.16 fluid ounces of **REGENT 4 SC** per acre.

COMPATIBILITY

The full range of compatibilities with **Regent 4 SC** under local conditions is not known. Therefore, it is essential that before using **Regent 4 SC**, the compatibility of the mixture be established. For example: Fill a quart jar 1/2 full of water or liquid fertilizer, then add 2 teaspoons of **Regent 4 SC** and disperse. Add any other pesticides, fertilizers, or tank mixtures and shake thoroughly. **DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GREASE. FOR BEST RESULTS, SPRAY MIXTURES SHOULD BE USED IMMEDIATELY AFTER MIXING. USE ADEQUATE AGITATION.**

**USE DIRECTIONS FOR POTATO
PESTS CONTROLLED AND APPLICATION METHOD**

Crop	Pests Controlled	Application Method
Potato	Wireworms	Make one in-furrow application at time of planting only. Banding must be done in-furrow as a 5- to 7-inch wide band. The Regent must be thoroughly incorporated and covered with soil immediately after application. Use of T-banding over top of closed furrow is NOT permitted.

RATE: Regent 4 SC Maximum Allowable Rate = 3.2 fl oz of Regent 4 SC per acre

Use the following chart to select the correct rate of **Regent 4 SC** per 1,000 row ft and rate of product per acre for potato.

*Row Spacing (inches)	Row ft per Acre	Rate (fl oz) per 1,000 Row ft	*Rate (fl oz) per Acre
30	17,424	0.184	3.2
34	15,374	0.208	3.2
36	14,520	0.220	3.2
38	13,756	0.220	3.0
40	13,068	0.220	2.9

DO NOT harvest within 90 days of application.

- * **DO NOT** apply in row spacing less than 30 inches. On any row spacing greater than 36 inches, apply no more than 0.220 fl oz **Regent 4 SC** per 1,000 row feet.
- * Adjust rate per 1,000 row feet so that you **DO NOT** apply more than 3.2 fl oz of **Regent 4 SC** per acre (equal to 0.10 lb ai/A).

9/13

COMPATIBILITY

The full range of compatibilities with **Regent® 4 SC insecticide** under local conditions is not known. Therefore, it is essential that before using **Regent 4 SC**, the compatibility of the mixture be established. For example: Fill a quart jar 1/2 full of water or liquid fertilizer, then add 2 teaspoons of **Regent 4 SC** and disperse. Add any other pesticides, fertilizers, or tank mixtures and shake thoroughly. **DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GREASE. FOR BEST RESULTS, SPRAY MIXTURES SHOULD BE USED IMMEDIATELY AFTER MIXING. USE ADEQUATE AGITATION.**

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Regent is a registered trademark of BASF.

© 2007 BASF Corporation
All rights reserved.

007969-00207.20070531b.NVA 2007-04-218-0105
Supersedes: NVA 2006-04-218-0022

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709



10/13



The Chemical Company

RESTRICTED USE PESTICIDE
DUE TO TOXICITY TO AQUATIC INVERTEBRATES

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

REGENT[®] 4 SC
INSECTICIDE

**Supplemental
Labeling**

For In-Furrow Use on Potatoes

Shake Well Before Use

EPA Reg. No. 7969-207

ACTIVE INGREDIENT:

Fipronil: 5-amino-1-(2,6-dichloro-4-(trifluoromethyl) phenyl)-
4((1,R,S)-(trifluoromethyl)sulfinyl)-1-*H*-pyrazole-3-carbonitrile 39.4%

OTHER INGREDIENTS:..... 60.6%

TOTAL: 100.0%

One gallon Regent[®] 4 SC insecticide contains 4 lbs fipronil.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

GENERAL PRECAUTIONS AND INSTRUCTIONS

SPRAY PREPARATION

Remove oil, rust, scale, pesticide residues and foreign matter from sprayer and strainer. Flush with clean water. Use 50 mesh screens or equivalent in spray system. Applicator tanks must be equipped with a sparge tube agitator, unless the alternative agitator designs can adequately cleanse the bottom of the tank and provide sufficient agitation required for product mixing. Pump approximately 10% of the tank capacity per minute back into the tank through the agitator.

To prepare for spraying, fill tank to approximately ½ the needed volume of water or liquid fertilizer and start agitation. Open product container(s) and pour contents into tank. Agitate to ensure thorough mixing for a minimum of 2 minutes or until product is thoroughly dispersed. Agitate product mixtures continuously during use. Avoid allowing tank contents to remain without agitation for periods greater than 1 hour. When tank contents have remained without agitation for 1 hour or longer, agitate contents for at least 10 minutes before starting application. Avoid allowing large quantities of product mixtures (greater than 5 gallons) to be stored in a non-agitated tank for extended periods of time, such as 12 hours or greater.

Regent® 4 SC insecticide can be tank mixed in accordance with the most restrictive of label limitations and precautions. **DO NOT** exceed any label dosage rates. This product cannot be mixed with any product whose label contains a prohibition against such mixing.

If tank mixes with other products are to be used, evaluate tank mix compatibility with **Regent 4 SC** in a small container before mixing larger quantities in the application tank.

COMPATIBILITY

The full range of compatibilities with **Regent 4 SC** under local conditions is not known. Therefore, it is essential that before using **Regent 4 SC**, the compatibility of the mixture be established. For example: Fill a quart jar ½ full of water or liquid fertilizer, then add 2 teaspoons of **Regent 4 SC** and disperse. Add any other pesticides, fertilizers, or tank mixtures and shake thoroughly. **DO NOT USE MIXTURES THAT CURDLE, PRECIPITATE, OR GREASE. FOR BEST RESULTS, SPRAY MIXTURES SHOULD BE USED IMMEDIATELY AFTER MIXING. USE ADEQUATE AGITATION.**

USE RESTRICTIONS

- **DO NOT** apply this product through any kind of irrigation system.
- **DO NOT** harvest within 90 days of application.
- **DO NOT** plant root, leafy or legume vegetables within four months following application.
- **DO NOT** plant wheat or triticale within two months following application.
- **DO NOT** plant any other rotational crops (except field corn, potato, leafy vegetables, root vegetables, legume vegetables, wheat, or triticale) within 12 months following application.

Observe the following precautions when applying in the vicinity of aquatic areas: **DO NOT** apply within 20 yards (**in New York State, DO NOT apply within 100 feet or 33 yards**) of lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, commercial aquaculture facilities or other bodies of water that convey water to these areas. Protection of aquatic areas may be enhanced by maintaining all or a portion of this buffer in vegetative cover.

DO NOT use this product if planted by hand.

APPLICATION

Make one in-furrow application at planting time only. **DO NOT** apply this product through any kind of irrigation system.

Dilute **Regent 4 SC** in a minimum of 5.0 gallons of water or liquid fertilizer per acre. If flat-fan nozzles are used for in-furrow application, the nozzles should be aligned with the row to direct the spray into the open furrow.

Regent 4 SC can be directly injected into the suction or pressure line(s) of a liquid spray system provided that the product is uniformly dispersed into the water or liquid fertilizer. Predilution of **Regent 4 SC** in a minimum of 0.5 gallons of water or liquid fertilizer per acre, prior to injection, is necessary to improve metering accuracy. The mix tank of **Regent 4 SC** plus water or liquid fertilizer must be agitated continuously. After completion of the direct injection in the mixing process, the final volume delivered to the furrow should have a minimum volume of 5.0 gallons per acre of water, liquid fertilizer or the combination of the two.

12/13

USE DIRECTIONS FOR POTATO

PESTS CONTROLLED AND APPLICATION METHOD

Crop	Pests Controlled	Application Method
Potato	Wireworms	Make one in-furrow application at time of planting only. Banding must be done in-furrow as a 5- to 7-inch wide band. Regent® 4 SC insecticide must be thoroughly incorporated and covered with soil immediately after application. DO NOT use T-banding over the top of a closed furrow.

RATE: Regent 4 SC Maximum Allowable Rate = 3.2 fl oz of Regent 4 SC per acre

Use the following chart to select the correct rate of **Regent 4 SC** per 1,000 row feet and rate of product per acre for potato.

*Row Spacing (inches)	**Rate/Acre (fluid ounces/acre)	Row Feet/Acre (feet)	Rate/1,000 row feet (fluid ounces/1,000 row feet)
30	3.2	17,424	0.184
34	3.2	15,374	0.208
36	3.2	14,520	0.220
38	3.0	13,756	0.220
40	2.9	13,068	0.220

* **DO NOT** apply in row spacing less than 30 inches. On any row spacing greater than 36 inches, apply no more than 0.220 fl oz **Regent 4 SC** per 1,000 row feet.

** Adjust rate per 1,000 row feet so that you **DO NOT** apply more than 3.2 fluid ounces of **Regent 4 SC** per acre (equal to 0.10 lb ai/A).

• **DO NOT** apply to sweet potatoes.

13/13

CONDITIONS OF SALE AND WARRANTY

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

807

Regent is a registered trademark of BASF.

© 2007 BASF Corporation
All rights reserved.

007969-00207.20070920.NVA 2007-04-218-0152
Based on: NVA 2007-04-218-0105

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

