

7969-242

8-24-2007

1/17



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

AUG 24 2007

Mr. Dave G. Bolin
Product Registration Manager-
US Regulatory Affairs
BASF Corporation
Agricultural Products
PO Box 13528
26 Davis Drive
Research Triangle Park, NC 27709-3528

RE: Changes made to Final Printed Label under notification criteria were to the Warranty Statement and Correction of Typographical/Printing Errors
EPA Registration Number: 7969-242
Date of Submission: July 24, 2007

Dear Mr. Bolin:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated July 24, 2007, for the product Status Herbicide. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to be "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
 Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 7969-242	2. EPA Product Manager Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Status Herbicide	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) BASF Corp, Agricultural Products PO Box 13528 Research Triangle Park, NC 27709-3528 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	---

NOTIFICATION

AUG 24 2007

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

As per PR Notice 98-10, BASF is submitting one (1) copy of the final printed labeling via notification. Changes made to final print label under notification criteria were to the Warranty Statement and Correction of Typographical / Printing Errors.

No PRIA fee is associated with this action.

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Unit Packaging wgt. 7.813 lbs No. per container 4	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes" Package wgt _____ No. per container _____	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
---	---	--	---

* Certification must be submitted

3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container 1 jug = 7.813 lbs; 4 jugs = case	5. Location of Label Directions <input checked="" type="checkbox"/> on label
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph Paper glued Stenciled <input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Dave G. Bolin	Title Product Registration Manager	Telephone No. (include Area Code) 919-547-2108
-----------------------	---------------------------------------	---

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Product Registration Manager	6. Date Application Received (Stamped) _____
4. Typed Name Dave G. Bolin	5. Date July 24, 2007	

4/17

Storage and Disposal

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store product in original container only. Store product in a cool, dry place. **DO NOT** store this product under wet conditions. Avoid cross-contamination with other pesticides.

Pesticide Disposal: Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal:

- **Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. **DO NOT** reuse container.

I. GENERAL INFORMATION

Status® herbicide is a selective postemergence herbicide for the control of annual broadleaf weeds, control or suppression of many perennial broadleaf weeds, and suppression of annual grasses found in field corn grown for grain, seed, or silage. **Status** is recommended to be used sequentially or tank mixed with a grass herbicide for a complete weed control program (refer to section **V. General Tank Mixing Information**).

Table 1. General Weed List, Including ALS¹, Glyphosate-resistant², protox (PPO)-resistant³, and Triazine-resistant⁴ Biotypes

Annual Broadleaf Weeds	
Amaranth, Palmer	Lettuce, Prickly
Beggarweed, Florida	Mallow,
Buckwheat, Wild	Common
Buffalobur	Venice
Burcucumber	Marestail (Horseweed)
Carpetweed	Morningglory,
Chickweed, Common	Entireleaf
Cocklebur, Common	Ivyleaf
Croton, Tropic	Pitted
Devil's Claw	Smallflower
Henbit	Tall
Jimsonweed	Nightshade,
Knotweed, Prostrate	Black
Kochia	Eastern Black
Ladysthumb	Hairy
Lambsquarters, Common	Pennycress, Field

(continued)

5/17

Table 1. General Weed List, Including ALS¹, Glyphosate-resistant², protox (PPO)-resistant³, and Triazine-resistant⁴ Biotypes (continued)

Annual Broadleaf Weeds	
Pigweed, Prostrate Redroot Smooth Spiny Tumble Purslane, Common Radish, Wild Ragweed, Common Giant Sesbania, Hemp Shepherdspurse Sicklepod	Sida, Pricky (Teaweed) Smartweed, Pennsylvania Smellmelon Sowthistle, Annual Spurge, Prostrate Sunflower, Common (Wild) Volunteer Thistle, Russian Velvetleaf Waterhemp, Common Tall
Perennial Broadleaf Weeds ⁵	Annual Grasses ⁶
Alfalfa Bindweed, Field Hedge Clover, White Dandelion, Common Dock, Broadleaf Curly Dogbane, Hemp Horsenettle, Carolina Knapweed, Spotted Milkweed, Common Honeyvine Nightshade, Silverleaf Plantain, Broadleaf Pokeweed Potato, Volunteer Smartweed, Swamp Sowthistle, Perennial Thistle, Canada	Barryardgrass Foxtail, Giant Green Yellow Johnsongrass (seedling) Panicum, Fall Shattercane Signalgrass, Broadleaf

6/17

¹ ALS (acetolactate synthase)-resistant weeds include those weeds resistant to the sulfonylurea (e.g. **Spirit™** and **Exceed® herbicides**), imidazolinone (e.g. **Lightning® herbicide**), or sulfonamide (e.g. **Python™ herbicide**) family of herbicides.

² Glyphosate-resistant weeds include those weeds resistant to glyphosate (e.g. **Roundup®**, etc).

³ PPO (protoporphyrinogen)-resistant weeds include those weeds resistant to diphenyl ether herbicides (e.g. **Cobra®** and **Blazer® herbicides**).

⁴ Triazine-resistant weeds include those weeds resistant to the triazine herbicides (e.g. atrazine).

⁵ Partially controlled or suppressed.

⁶ **Status® herbicide** provides suppression of annual grasses at appropriate rates (5 oz per acre or greater). Emerged grasses up to 3" tall will cease growing but may remain green for weeks after application. Regrowth of grasses is limited once corn canopies over the row middles.

Mode of Action

Status is absorbed by leaves, roots, and shoots and translocated to the growing points of sensitive weeds to provide postemergence control of emerged weeds as well as moderate residual control of germinating weeds. **Status** controls weeds by auxin transport inhibition and auxin agonist modes of action.

Crop Tolerance

Corn is very tolerant to an application of **Status**. Corn growing under stress conditions such as drought, poor fertility, or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if **Status** is applied. Injury can be avoided by agronomic practices that promote good crop growth and minimize stress conditions, especially combinations of stress factors.

Coverage

Weeds must be thoroughly covered with spray. Dense leaf canopies shelter smaller weeds and can prevent adequate spray coverage.

Cultivation

Avoid disturbing (e.g. tillage or cultivating) treated areas for at least 7 days following application to allow best herbicide uptake, translocation, and weed control.

Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions and triple rinsing the equipment before and after applying this product.

II. Application Instructions

Best product performance is obtained when **Status** is applied to actively growing weeds. Treated weeds will stop growing soon after application. Broadleaf weeds will display epinastic twisting and crinkling symptoms before becoming necrotic. Suppressed grass weeds may display some epinasty and remain stunted and green.

Ground Application Methods and Equipment

As a wettable granule formulation, **Status** can be applied using water as the spray carrier.

Water Volume: Use 3-50 gallons of diluted spray solution per treated acre. Use the higher water volume when treating dense or tall vegetation.

Application Equipment: Use application equipment that will provide good spray coverage of weed foliage. Exercise preventative measures to avoid drift onto nontarget areas.

7/17

Spray Drift Management

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, spray nozzle/pressure combinations, spray droplet size, temperature/humidity, etc. Contact your state extension agent for spray drift prevention guidelines in your area. All ground application equipment must be properly maintained and calibrated using appropriate carriers. Avoiding spray drift at the application site is the responsibility of the applicator. Agriculturally approved drift reducing additives are recommended.

III. Additives

Adjuvants must be used with **Status® herbicide** to achieve consistent weed control. The best results under most conditions will be achieved by combining an adjuvant (NIS, MSO, or COC) with a nitrogen source (urea ammonium nitrate or ammonium sulfate). Refer to **Table 2. Additive Rates** for additive rates.

Nitrogen Source

Use a minimum of 5 quarts of urea ammonium nitrate (UAN; 28-34% nitrogen) per 100 gallons. Spray grade ammonium sulfate (AMS; 21% nitrogen) may be substituted for UAN at a rate of 5 to 17 pounds per 100 gallons of spray mix. Use high-quality AMS (spray grade) to avoid plugging of nozzles. When using AMS, dissolve **Status** in the tank before adding AMS. Because most nitrogen solutions are mildly corrosive to galvanized, mild steel, and brass spray equipment, rinse the entire spray system with water soon after use.

Table 2. Additive Rates

Adjuvant		Nitrogen source
Nonionic Surfactant (0.25% v/v or 1 qt/100 gallon)	+	5 quarts of UAN (1.25% v/v) or 5 to 17 pounds of AMS
COC (1.0 to 2.0 pt/a)	+	5 quarts of UAN (1.25% v/v) or 5 to 17 pounds of AMS
MSO (1.0 to 2.0 pt/a)	+	5 quarts of UAN (1.25% v/v) or 5 to 17 pounds of AMS

Compatibility Test for Mix Components

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

- 1) **Water:** For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
- 2) **Products in PVA bags:** Cap the jar and invert 10 cycles.
- 3) **Water-dispersible products (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions):** Cap the jar and invert 10 cycles.
- 4) **Water-soluble products (such as Status):** Cap the jar and invert 10 cycles. **Status** must be fully dissolved before adding other components.
- 5) **Emulsifiable concentrates (including COC and MSO):** Cap the jar and invert 10 cycles.
- 6) **Water-soluble additives (NIS, and AMS or UAN when applicable):** Cap the jar and invert 10 cycles.

8/17

7) Let the solution stand for 15 minutes.

8) Evaluate the solution for uniformity and stability.

The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, **DO NOT** mix the ingredients in the same tank.

IV. Mixing Order

1) **Water:** Begin by agitating a thoroughly clean sprayer tank half full of clean water.*

2) **Products in PVA bags:** Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.

3) **Water-dispersible products (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions).**

4) **Water-soluble products (such as Status® herbicide):** Status must be fully dissolved before adding other components.

5) **Emulsifiable concentrates (including COC and MSO).**

6) **Water-soluble additives (nonionic surfactant, AMS, or UAN when applicable).**

7) **Remaining quantity of water:**

Maintain constant agitation during application. For more information, refer to section V. **General Tank Mixing Information.** Refer to the drift reducing additive label for proper addition rate and mixing order.

* If the user prefers to fill the spray tank from a nurse tank containing an AMS product dissolved in water, this is acceptable. If this method is used, the AMS product must be totally dissolved before adding Status. Status must be thoroughly dissolved before adding additional products or additives. The user should verify that the AMS pre-mix water alternative is compatible with other tank mix components.

V. General Tank Mixing Information

Status is recommended to be used sequentially or tank mixed with other herbicides as part of a complete weed control program. Tank mix recommendations are for use only in states where the sequential or tank mix product and application site are registered. Refer to section VII. **Crop-Specific Information** for more details and for specific tank mix restrictions. Local agricultural authorities may be a source of information when using other than BASF-recommended tank mixes. Read and follow the applicable **Restrictions and Limitations and Directions For Use** on all products involved in tank mixing. The most restrictive labeling applies to tank mixes. Physical incompatibility, reduced weed control, or crop injury may result from mixing Status with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers.

VI. General Restrictions and Limitations

- **Maximum Seasonal Use Rate:** **DO NOT** apply more than a total of 12.5 oz of Status (0.438 pounds ae) per acre per season.
- **DO NOT** apply if corn is more than 36" tall, or V10 stage, or within 15 days before tassel emergence, whichever comes first.

9/17

- **Restricted-entry Interval (REI): 24 hours.**
- **Preharvest Interval (PHI): DO NOT** apply within 32 days of corn forage harvest. **DO NOT** apply within 72 days of corn grain and stover harvest (Table 3).
- Allow a minimum of 15 days between sequential applications of **Status® herbicide**.
- **Crop Rotation Restrictions: DO NOT** plant any crops within 120 days after the last application of **Status**, with the following exceptions. If at least 1 inch of rainfall or overhead irrigation is received following the last application of **Status** (less than or equal to 5 oz per acre only), alfalfa, cereal grain crops, cotton, grain sorghum and soybeans may be planted 30 days after the rainfall/irrigation event in all states except California. In the event of crop failure, corn can be replanted 7 or more days after application.
- **Rain-fast Period: Status** is rain fast **4 hours** after application when used with recommended adjuvants according to section III. **Additives**.
- **DO NOT** apply to corn showing injury (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications because this injury may be enhanced or prolonged.
- **DO NOT** use sprayable fluid fertilizer as the carrier for application of **Status** made after corn emergence.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** treat irrigation ditches or water used for crop irrigation or domestic uses.

Table 3. Crop-Specific Restrictions and Limitations

Crop	Minimum Time From Application to Harvest (PHI)	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season
Corn forage	32 days	10 oz	12.5 oz
Corn grain or stover	72 days		

VII. Crop-Specific Information

CORN

FIELD CORN GROWN FOR GRAIN, SEED, OR SILAGE

Before applying **Status** to seed corn, verify the selectivity of **Status** on the inbred line or hybrid with your local seed corn company (supplier). This precaution will help avoid potential injury to sensitive lines. **Status is not registered for use on sweet corn or popcorn.**

Application Rates and Timings:

Status herbicide can be applied postemergence at the rates and growth stages listed in **Table 4** for all tillage systems (e.g. conventional, no-tillage, reduced tillage).

• **Postemergence Application in Corn:**

- **Status** can be applied from 4-inch tall (V2) to 36-inch tall (V10) corn at rates from 5 to 10 oz per acre.
- Early postemergence application is recommended for best weed control and crop yield potential.
- **Status** may be tank mixed with **Lightning®, Roundup®,** glyphosate or **Liberty® herbicides.**
- Apply 2.5 to 10 oz of **Status** with the tank mix herbicides (e.g. **Lightning, Roundup,** glyphosate or **Liberty**).

10/17

- Use the minimum rate of 5 oz/acre of **Status® herbicide** for weeds that are resistant to the tank mix herbicide (ALS, glyphosate, or PPO resistance), for perennial weeds listed on the weed list, weeds taller than 6 inches, or weeds not controlled by the tank mix partner.
- **Lightning®, Roundup®, glyphosate and Liberty® herbicides** can only be used on specifically designated corn varieties (e.g. **Roundup** or glyphosate on **Roundup Ready®** corn, **Liberty** on **LibertyLink®** corn, and **Lightning** on **CLEARFIELD®** corn). **DO NOT** use on corn varieties that are not labeled for use.

• **Split Applications:**

Split applications of **Status** may be made during a growing season. **DO NOT** exceed a total of 12.5 oz of **Status** per treated acre per crop year. Allow a minimum of 15 days between sequential applications of **Status**.

Field Corn Tank Mixes and Sequential Uses

In addition to control of many broadleaf weed species, **Status** offers herbistatic activity (suppression) of several annual grasses that may enhance overall control of your grass herbicide program. For commercial control of grasses, **Status** should be used either as a sequential postemergence treatment following a preemergence grass herbicide (e.g. **Outlook®, Guardsman Max®, or G-Max Lite™ herbicides**) or in tank mix combination with a postemergence herbicide that controls grass (e.g. **Liberty, Lightning, Roundup, glyphosate, or Option® herbicides**).

Tank Mix Restrictions and Limitations

Status may be applied sequentially or in tank mixes with other herbicides registered for use in corn with the following limitations:

- Postemergence applications of **Status** are not recommended for use in tank mixes with plant growth regulating herbicides such as those products that contain dicamba, 2,4-D, or clopyralid herbicides. Additionally, sequential treatments with these products should be separated by at least 15 days.
- Tank mixes with emulsifiable concentrate (EC) formulations of chloroacetamide grass herbicides (e.g. **Dual II Magnum®, Harness®, Outlook, or Surpass® herbicides**) are not recommended.
- **Status** may be used sequentially with all soil-applied insecticides or used sequentially or in tank mixes with foliar-applied insecticides with the following limitations: **Status** is not recommended for use in foliar-applied tank mixes with **Lorsban™ insecticide**. However, sequential treatments with these products may be used if applications are separated by at least 7 days.

Table 4. Application Rates and Timing in Field Corn for Status

Application Timing	Corn Stage (inches)	Status Solo Application ¹	Status Tank Mix Applications ²
		Rate of Status per acre ³	
Postemergence	4" to 36" (V2 - V10)	5 - 10 oz,	2.5 - 10 oz

¹ Solo rate also includes tank mixes with grass herbicides with little broadleaf activity.

² Tank mixes with **Lightning, Roundup, glyphosate, or Liberty**. Refer to their respective labels for use rates.

³ Increase the rate of **Status** if target weeds are resistant to tank mix partner, are perennial weeds, are taller than 6", or weeds not controlled by tank mix partner.

11/17

Pests Listed in This Label	
Common Name	Scientific Name
Alfalfa	<i>Medicago sativa</i>
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Beggarweed, Florida	<i>Desmodium tortuosum</i>
Bindweed, Field	<i>Convolvulus arvensis</i>
, Hedge	<i>Calystegia sepium</i>
Buckwheat, Wild	<i>Polygonum convolvulus</i>
Buffalobur	<i>Solanum rostratum</i>
Burcucumber	<i>Sicyos angulatus</i>
Carpetweed	<i>Mollugo verticillata</i>
Chickweed, Common	<i>Stellaria media</i>
Clover, White	<i>Trifolium repens</i>
Cocklebur, Common	<i>Xanthium strumarium</i>
Croton, Tropic	<i>Croton glandulosus</i>
Dandelion, Common	<i>Taraxacum officinale</i>
Devil's Claw	<i>Proboscidea louisianica</i>
Dock, Broadleaf	<i>Rumex obtusifolius</i>
, Curly	<i>Rumex crispus</i>
Dogbane, Hemp	<i>Apocynum cannabinum</i>
Foxtail, Giant	<i>Setaria faberi</i>
, Green	<i>Setaria viridis</i>
, Yellow	<i>Setaria glauca</i>
Henbit	<i>Lamium amplexicaule</i>
Horsenettle, Carolina	<i>Solanum carolinense</i>
Jimsonweed	<i>Datura stramonium</i>
Johnsongrass, Seedling	<i>Sorghum halepense</i>
Knapweed, Spotted	<i>Centaurea maculosa</i>
Knotweed, Prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, Common	<i>Chenopodium album</i>
Lettuce, Pricky	<i>Lactuca serriola</i>

(continued)

12/17

Pests Listed in This Label (continued)	
Common Name	Scientific Name
Mallow, Common	<i>Malva neglecta</i>
, Venice	<i>Hibiscus trionum</i>
Marestail (Horseweed)	<i>Coryza canadensis</i>
Milkweed, Common	<i>Asclepias syriaca</i>
, Honeyvine	<i>Ampelamus albidus</i>
Morningglory, Entireleaf	<i>Ipomoea hederacea var. integruscula</i>
, Ivyleaf	<i>Ipomoea hederacea var. hederacea</i>
, Pitted (Small White)	<i>Ipomoea lacunosa</i>
, Smallflower	<i>Jacquemontia tamnifolia</i>
, Tall (Common)	<i>Ipomoea purpurea</i>
Nightshade, Black	<i>Solanum nigrum</i>
, Eastern Black	<i>Solanum ptycanthum</i>
, Hairy	<i>Solanum sarachoides</i>
, Silvertleaf	<i>Solanum elaeagnifolium</i>
Panicum, Fall	<i>Panicum dichotomiflorum</i>
Pennycress, Field	<i>Thlaspi arvense</i>
Pigweed, Prostrate	<i>Amaranthus blitoides</i>
, Redroot	<i>Amaranthus retroflexus</i>
, Smooth	<i>Amaranthus hybridus</i>
, Spiny	<i>Amaranthus spinosus</i>
, Tumble	<i>Amaranthus albus</i>
Plantain, Broadleaf	<i>Plantago major</i>
Pokeweed	<i>Phytolacca americana</i>
Potato, Volunteer	<i>Solanum tuberosum</i>
Purslane, Common	<i>Portulaca oleracea</i>
Radish, Wild	<i>Raphanus raphanistrum</i>
Ragweed, Common	<i>Ambrosia artemisiifolia</i>
, Giant	<i>Ambrosia trifida</i>
Sesbania, Hemp	<i>Sesbania exaltata</i>
Shattercane	<i>Sorghum bicolor</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>

13/17

Pests Listed in This Label (continued)	
Common Name	Scientific Name
Sicklepod	<i>Cassia obtusifolia</i>
Sida, Prickly (Teaweed)	<i>Sida spinosa</i>
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Swamp	<i>Polygonum coccineum</i>
Smellmelon	<i>Cucumis melo</i>
Sowthistle, Annual	<i>Sonchus oleraceus</i>
Perennial	<i>Sonchus arvensis</i>
Spurge, Prostrate	<i>Euphorbia humistrata</i>
Sunflower, Volunteer	<i>Helianthus annuus</i>
Wild (Common)	<i>Helianthus annuus</i>
Thistle, Canada	<i>Cirsium arvense</i>
Russian	<i>Salsola iberica</i>
Velvetleaf	<i>Abitilon theophrasti</i>
Waterhemp, Common	<i>Amaranthus rudis</i>
Tall	<i>Amaranthus tuberculatus</i>

Crops
This product can be used on the following crops:
Field Corn grown for grain, seed, or silage.
See inside for complete Restrictions and Limitations and Application Instructions .

14/17

Conditions of Sale and Warranty

The **Directions For Use** of this product reflects the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Status® herbicide is specially formulated for use on crops as specified in this label. Bayer CropScience is the owner of United States patent rights to methods and compositions covering certain uses, particularly the following United States Patents: 6,486,096; 6,569,805; 5,922,646; and 5,516,750.

Status, CLEARFIELD, Guardsman Max, Lightning, and Outlook are registered trademarks of BASF. **G-Max Lite** is a trademark of BASF.

Blazer is a registered trademark of United Phosphorus, Inc.

Cobra is a registered trademark of Valent U.S.A. Corporation.

Dual II Magnum and Exceed are registered trademarks of a Syngenta Group Company.

Harness, Roundup and Roundup Ready are trademarks of Monsanto Technology LLC.

Liberty, LibertyLink and Option are registered trademarks of Bayer.

Lorsban, Surpass and Python are trademarks of Dow AgroSciences LLC.

Spirit is a trademark of a Syngenta Group Company.

© 2007 BASF Corporation

All rights reserved.

007969-00242.20070601.NVA 2007-04-279-0102

Based on: NVA 2006-04-279-0188

Supersedes: NVA 2006-04-279-0266

BASF Corporation

Agricultural Products


26 Davis Drive

Research Triangle Park, NC 27709



The Chemical Company

15/17



16/17

BASF

The Chemical Company



FOR USE IN FIELD CORN GROWN FOR GRAIN, SEED, OR SILAGE.
A broadspectrum corn herbicide, in a safened formulation, specifically designed for use alone or with glyphosate combinations.

Active Ingredients:

Sodium salt of diflufenzopyr: 2-[1-[(3,5-difluorophenylamino) carbonyl]-hydrazono]ethyl]-3-pyridinecarboxylic acid, sodium salt	17.1%
Sodium salt of dicamba: 3,6-dichloro-2-methoxybenzoic acid, sodium salt**	44.0%
Other Ingredients:	38.9%
Total:	100.0%

*This product contains 16% 2-[1-[(3,5-difluorophenylamino) carbonyl]-hydrazono]ethyl]-3-pyridinecarboxylic acid (diflufenzopyr).
**This product contains 40% 3,6-dichloro-2-methoxybenzoic acid.

EPA Reg. No. 7969-242

EPA Est. No. 11773-IA-001

**KEEP OUT OF REACH OF CHILDREN.
CAUTION/PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID: If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. **DO NOT** induce vomiting unless told to by a poison control center or doctor. **DO NOT** give anything by mouth to an unconscious person. **If on skin:** Take off contaminated clothing. Rinse immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **HOT LINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357). **PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION:** Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. **ENVIRONMENTAL HAZARDS: DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters. This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination. **See attached booklet for complete Environmental Hazards & Ground and Surface Water Protection.** **STORAGE AND DISPOSAL: DO NOT** contaminate water, food, or feed by storage or disposal. **Pesticide Storage:** Store product in original container only. Store product in a cool, dry place. **DO NOT** store this product under wet conditions. Avoid cross-contamination with other pesticides. **Pesticide Disposal:** Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **Container Disposal: Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. **DO NOT** reuse container. See attached booklet for complete First Aid, Precautionary Statements, Directions For Use, and Conditions of Sale and Warranty. In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357). Product of U.S.A.

Net Contents: 125 ounces

Produced for: BASF Corporation, Agricultural Products
26 Davis Drive, Research Triangle Park, NC 27709

2088315
NVA 2007-05-279-0208



The Chemical Company

Agricultural Products

17/17

Ms. Joanne Miller, Room Number 4900
U.S. Environmental Protection Agency
Registration Division (7505P), PM Team 23
One Potomac Yard (South Building)
2777 S. Crystal Drive
Arlington, VA 22202

July 24, 2007

**Re: Status® Herbicide (EPA Registration No. 7969-242)
Notification of Final Print Labeling.**

Dear Ms. Miller,

BASF hereby submits one (1) copy of the final print label for Status Herbicide. This communication is submitted in accordance with PR Notice 98-10. Also included are: completed EPA form 8570-1, copy of the stamped approved label, and your comments letter dated 12/15/06.

As defined in PR Notice 98-10 BASF has made the following types of changes allowed by notification to the Agency:

- Changes in the Warranty Statement
- Correction of typographical / printing errors
- Changed "Inert Ingredients" to "Other Ingredients"

As no data review is required, BASF believes that there is no PRIA fee associated with this action.

Please feel free to contact me (919-547-2108 / david.bolin@basf.com) if you have any questions.

Respectfully Submitted,

Dave G. Bolin
Product Registration Manager
U.S. Regulatory Affairs

® Registered Trademark of BASF

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709
Tel: (800) 669-1770
www.agproducts.basf.com