

7969-255

07/03/2007

1/12



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:
7969-255

Date of Issuance:
JUL 3 2007

NOTICE OF PESTICIDE:
 Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance: Conditional

Name of Pesticide Product:
Honor Fungicide

Name and Address of Registrant (include ZIP Code):

Attention: Khalid Akkari
BASF Corporation
P.O. Box 13528
26 Davis Drive
Research Triangle Park, NC 27709

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA Section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for re-registration of your product under FIFRA Section 4.
2. Before releasing the product for shipment, revise your label's EPA Registration Number to read "EPA Reg. No. 7969-255."
3. On or by December 28, 2007, registrant must submit a valid Primary Eye Irritation study to the Agency.

Signature of Approving Official:

Tony Kish
Product Manager, Team 22
Fungicide Branch
Registration Division (7505P)

Date:

JUL 3 2007

2/12

EPA Registration No. 7969-255
Honor Fungicide
Page 2 of 2

4. Submit a GLP one year storage stability study (830.6317) and corrosion characteristics study (830.6320) by July 2, 2008.
5. Throughout the label 'Dollar Spot', must be accompanied by the treatment limitation: 'Suppression Only'. This may be indicated by appending a superscript, or an asterisk to 'dollar spot'.
6. Under the section "Conditions of Sale and Warranty," its first paragraph's final sentence must be appended by: "*To the extent permitted by applicable law....* " Likewise, the clause "*To the extent permitted by applicable law...* " is to be appended to the first sentence in the final paragraph. As currently written the statement negates and/or detracts from labeling requirements as set forth by the Agency in 40 CFR Part 156, and is, therefore; considered misbranding.
7. For your own records and for future EPA submissions, delete 'cite-all' from the "Guidelines Reference" column, on each page of the generic data matrix -- this pertains to pages 15 through 27. Presently, this selection conflicts with the selective generic data matrix.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release of the shipment of your product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted with Comments" is enclosed for your records.

Enclosures

3/12



The Chemical Company

GROUP 7 11 FUNGICIDE

Honor[®]

fungicide

For disease control on golf course turfgrass.

Active Ingredients:

Pyraclostrobin , (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester)	16.8%
Boscalid , 3-pyridinecarboxamide, 2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl)	11.2%
Other Ingredients:	72.0%
Total:	100.0%

0.168 oz (0.0105 lb) of pyraclostrobin in 1 oz of Honor[®] fungicide
0.112 oz (0.007 lb) of boscalid in 1 oz of Honor

EPA Reg. No. 7969-xxx

EPA Est. No. _____

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See the attached booklet for complete
First Aid, Precautionary Statements, Directions For Use, and
Conditions of Sale and Warranty.

Net Contents: _____

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

**ACCEPTED
with COMMENTS
In EPA Letter Dated
JUL 3 2007**
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

7969-255

4/12

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).</p>	

Precautionary Statements

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. For more options, refer to **Category A** on an EPA chemical resistance category selection chart.

Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material (such as nitrile, butyl, neoprene, and/or barrier laminate).
- Shoes plus socks.

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4 - 6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

DO NOT apply directly to water, areas where surface water is present, or intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash waters or rinsate.

5/12

Endangered Species Concerns

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is violation of federal law.

This pesticide is toxic to fish and aquatic invertebrates and should be used strictly in accordance with drift precautions on this label in order to minimize off-site exposures. **DO NOT** apply when weather conditions favor drift from treated areas to nontarget aquatic habitats. Notify state and/or federal authorities and BASF immediately if you observe any adverse environmental effects due to use of this product.

To determine whether your county has endangered aquatic species, consult the County Bulletins at <http://www.epa.gov/espp/usa-map.htm>.

Endangered Species Bulletins may also be obtained from extension offices or state pesticide agencies. If a bulletin is not available for your specific area, check with the appropriate local state agency to determine if known populations of endangered aquatic species occur in the area to be treated.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation. This label must be in the user's possession during application.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

This product is intended for golf course use only. Not for use on residential turfgrass, turfgrass being grown for sale or other commercial use such as sod production, seed production or for research purposes.

Nonagricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouses.

DO NOT enter or allow others to enter treated areas until sprays have dried.

Storage and Disposal

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original containers only. Keep container closed when not in use. **DO NOT** store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper dis-

posal can be made if product cannot be used according to label.

Pesticide Disposal: Wastes resulting from using this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

In Case of Spill

In case of large-scale spillage regarding this product, call:
CHEMTREC 1-800-424-9300
BASF Corporation 1-800-832-HELP (4357)

Steps to be taken in case material is released or spilled:

Dike and contain the spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.

Remove contaminated clothing and wash affected skin areas with soap and water.

Wash clothing before reuse.

Keep the spill out of all sewers and open bodies of water.

I. General Information

Honor® fungicide is a broad spectrum fungicide recommended for the control of many important diseases of golf course turfgrass. For maximum efficacy, **Honor** should be applied preventively. **Honor** may be applied as a solo foliar spray or in tank mixes with other registered fungicides.

DO NOT exceed the recommended application rate or fail to comply with use restrictions listed in the **Resistance Management** and **Restrictions and Limitations** sections. All applications should be made according to the use directions that follow. Failure to follow directions and precautions on this label may result in injury and/or inferior disease control.

This package contains **Honor**, a water dispersible granule (WG). The active ingredients in **Honor** are pyraclostrobin, a member of the strobilurin class of chemistry, and boscalid, a member of the carboxamide (anilide) class of chemistry. Optimum disease control is achieved when **Honor** is applied in a regularly scheduled protective spray program and used in a rotation program with other fungicides. Because of its high specific activity, **Honor** has good residual activity against target fungi.

Mode of Action: Pyraclostrobin and boscalid, the active ingredients of **Honor**, belong to the group of respiration inhibitors classified by the USEPA and Canada PMRA as Target Site of Action **Group 11 and Group 7** fungicides, respectively.

6/12

II. Application Information

Use Sites:

Use on turfgrass - Honor® fungicide may be used for disease control in the following turf use sites:

- golf courses

Application Instructions:

For maximum efficacy, **Honor** should be applied prior to or in the early stages of disease development. For maximum efficacy, apply **Honor** at the rates indicated in **Table 1. - Application Rates and Intervals for Honor fungicide on Golf Course Turfgrass** in 1 to 4 gallons of water per 1000 square feet (43 to 174 gallons per acre). Use the shorter specified application interval and/or the higher specified rate when prolonged favorable disease conditions exist. Applications should be repeated at the specified interval as necessary.

- **Honor** is most effective when applied preventively.
- Actual length of disease control will vary depending on environmental conditions, disease pressure, and management practices.
- Calibrate spray prior to use.
- After application, allow foliage to dry prior to mowing or irrigation.
- Apply **Honor** using sufficient water volume and pressure for adequate coverage of the foliage.
- Apply the recommended rate of **Honor** as instructed in the **Use Directions** section with ground spray equipment.

Aerial application of **Honor** is not permitted.

Spray Drift Management

DO NOT spray when conditions favor drift beyond area intended for application. Conditions that contribute to drift include thermal inversion, wind speed and direction, spray nozzle/pressure combinations, spray droplet size, temperature/humidity, etc. Contact your state extension agent for spray drift prevention guidelines in your area. All application equipment must be properly maintained and calibrated using appropriate carriers. Avoiding spray drift at the application site is the responsibility of the applicator.

Wind

Drift potential is lowest when wind speed does not exceed 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

Low humidity and high temperatures increase the evaporation of spray droplets and therefore the likelihood of increased spray drift.

Avoid spraying during conditions of low humidity and/or high temperatures. When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation.

Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. bodies of water or non-target crops) is minimal and when wind is blowing away from the sensitive areas.

Resistance Management

Honor fungicide contains two active ingredients: pyraclostrobin, a target site **Group 11** fungicide, and boscalid a target site **Group 7** fungicide. **Honor** is effective against pathogens resistant to fungicides with modes of action different from those of **Groups 7** and **11**, such as the dicarboximides, sterol inhibitors, benzimidazoles, or phenylamides. Fungal isolates resistant to fungicide in **Group 11** (such as pyraclostrobin, azoxystrobin, and trifloxystrobin) or **Group 7** (such as bocalid and flutolanil) may eventually dominate the population if such fungicides are used predominantly and repeatedly in the same area for successive years as the primary method of control for the targeted pathogen species. This may result in reduction of disease control by **Honor** or other **Group 7** or **11** fungicides. In order to maintain its performance, **DO NOT** exceed the total number of sequential applications of **Honor**. Adhere to the label instructions regarding consecutive use of **Honor** and other **Group 7** or **11** fungicides with a similar site of action on the same pathogens.

The following recommendations may be considered to delay the development of fungicide resistance:

- 1. Tank mixtures:** Use tank mixtures with fungicides from different target site of action groups that are registered/permitted for the same use and that are effective against the pathogens of concern. BASF recommends using at least the minimum labeled rates of each fungicide in the tank mix.
- 2. IPM: Honor** should be integrated into an overall disease and pest management program. Cultural practices known to reduce disease development should be followed. **Honor** may be used in advisory (disease forecasting) programs which recommend application timing

7/12

based on environmental factors favorable for disease development.

3. Monitoring: Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development. If a **Group 7** or **11** target site fungicide such as **Honor® fungicide** appears to be less effective against a pathogen that it previously controlled or suppressed, contact a BASF representative or local expert for further investigation.

In turfgrass, DO NOT make more than two (2) sequential applications of **Honor** for Pythium, gray leaf spot, dollar spot, or anthracnose. Then alternate to an effective fungicide from a different target site group before reapplying **Honor**. **DO NOT** make more than three (3) consecutive applications of **Honor** for all other turfgrass diseases. Then alternate to an effective fungicide from a different target site group before reapplying **Honor**.

III. Addition of Additives

DO NOT use with organosilicate-based adjuvants or injury may occur. Due to the large number of additives or adjuvants that may be used, neither the manufacturer nor the seller has determined whether **Honor** can be used safely with all additives.

IV. General Tank Mixing Information

Tank Mix Partners/Components

Honor is compatible with most fungicide, insecticide and fertilizer products. If tank mixtures are used, adhere to restrictions due to rates, label recommendations and precautions on all labels. Physical incompatibility, reduced disease control, or plant injury may result from mixing **Honor** with fungicides, herbicides, insecticides, additives, or fertilizers. To improve control of certain diseases, **Honor** may be tank mixed with other effective fungicides from a different target site group.

Compatibility Test for Tank Mix Components:

Add components in the following sequence using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre.

- 1) **Water:** For 87 gallons per acre spray volume, use 14.4 cups (3.5 liters) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
- 2) **Water-dispersible products:** (dry flowables, wettable powders, suspension concentrates, or suspo-emulsions) Cap the jar and invert 10 cycles.
- 3) **Water-soluble products:** Cap the jar and invert 10 cycles.
- 4) **Emulsifiable concentrates:** (oil concentrate or methylated seed oil when applicable) Cap the jar and invert 10 cycles.
- 5) **Water-soluble additives:** Cap the jar and invert 10 cycles.
- 6) Let the solution stand for 15 minutes.
- 7) Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface,

nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. **DO NOT** use any spray solution that could clog spray nozzles.

Mixing Order:

Limit amount of spray mixture prepared to that needed for immediate use.

- 1) **Water:** Begin by agitating a thoroughly clean sprayer tank half full of clean water.
- 2) **Products in PVA bags:** Place the water-soluble PVA bag into the mixing tank. The water-soluble PVA bag will dissolve in water to allow the contents to disperse. Wait until all water-soluble PVA bags have fully dissolved, and the product is evenly mixed in the spray tank before continuing.
- 3) **Water-dispersible products** (dry flowables such as **Honor**, wettable powders, suspension concentrates, or suspo-emulsions).
- 4) **Water-soluble products.**
- 5) **Emulsifiable concentrates** (oil concentrate or methylated seed oil when applicable).
- 6) **Water-soluble additives** (AMS or UAN when applicable).
- 7) Remaining quantity of **water**. Maintain maximum constant agitation during application. **DO NOT allow mixture to stand for extended periods prior to application.**

Cleaning Spray Equipment:

Spraying equipment must be cleaned thoroughly before and after applying this product, particularly if a product with the potential to injure turfgrass was used prior to **Honor**.

TURFGRASS USE DIRECTIONS

Honor is recommended for the control of anthracnose, bentgrass dead spot, Bermudagrass decline, brown patch, dollar spot, fairy ring, Fusarium patch, gray leaf spot, gray snow mold, large patch, leaf spot, melting out, necrotic ring spot, pink patch, pink snow mold, powdery mildew, Pythium blight, Pythium root dysfunction, rapid blight, red thread, Rhizoctonia leaf or sheath spot, rust, summer patch, take-all patch and yellow tuft (downy mildew). For optimum control of gray snow mold and pink snow mold, tank mix **Honor** with another effective fungicide from a different target site group.

Turfgrass Uses and Tolerance:

Due to variability within turfgrass species, application techniques and possible tank mixes, neither the manufacturer nor the seller has determined whether or not **Honor** can safely be used on all turfgrasses under all conditions. Therefore, it is recommended that the user determine if **Honor** can be used safely before broad use. Apply the recommended use rate of **Honor** on a small test area under conditions expected to be encountered. Monitor for any adverse effects during a 14-day period after application. **Rate:** Use the application rates specified for each disease as listed in **Table 1. - Application Rates and Intervals for Honor fungicide on Golf Course Turfgrass**. Apply **Honor** in 1 to 4 gallons of water per 1000 square feet (43-174 gallons per acre).

8/12

RESTRICTIONS AND LIMITATIONS

- **Maximum seasonal use rate: DO NOT** apply more than a total of 6.6 ounces of **Honor® fungicide** per 1000 sq ft per year (18.0 pounds of **Honor** per acre per year).
- Refer to **Table 1. - Application Rates and Intervals for Honor fungicide on Golf Course Turfgrass** for sequential application intervals for **Honor**.
- **DO NOT** apply this product to crops other than **golf course turfgrass**.
- **DO NOT** apply to turfgrass through any type of **irrigation** equipment.
- This product cannot be used to **formulate** or reformulate any other pesticide product.

9/12

Table 1. - Application Rates and Intervals for Honor® fungicide on Golf Course Turfgrass

Disease (Pathogen)	Use Rate (oz Product per 1000 sq ft)	Use Rate (oz Product per Acre)	Application Interval (days)	Comments
Anthracnose ¹ (<i>Colletotrichum graminicola</i>)	0.55 to 1.1	24 to 48	14 to 28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.
Bentgrass Dead Spot (<i>Ophiosphaerella agrostis</i>)	0.55 to 1.1	24 to 48	14 to 28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.
Bermudagrass Decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	1.1	48	Not Applicable (see comments)	Aids in control of Bermudagrass decline when integrated with appropriate cultural practices such as raised mowing height, proper fertilization and core aeration. Make one application in the spring following green-up and a second application in the fall when air temperatures remain above 80° F and humidity is 75% or higher. Apply in 4 gallons of water per 1000 sq ft.
Brown Patch (<i>Rhizoctonia solani</i>)	0.55 to 1.1	24 to 48	14 to 28	Apply when conditions are favorable for disease development.
Dollar Spot ¹ (<i>Sclerotinia homoeocarpa</i>)	0.83 to 1.1	36 to 48	14 to 21	Begin applications prior to or in the early stages of disease development. Use the shorter specified application interval and/or the higher specified rate when prolonged favorable disease conditions exist.
Fairy Ring (various <i>Basidiomycete</i> fungi)	1.1	48	28	Apply as soon as possible after fairy ring symptom development. Fairy ring symptoms may take 2 to 3 weeks to disappear following application. Reapplication after 28 days maybe required.
Fusarium Patch (<i>Microdochium nivale</i>)	0.55 to 1.1	24 to 48	14 to 28	In the absence of snow cover, use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.
Gray Leaf Spot ¹ (<i>Fyricularia grisea</i>)	0.55 to 1.1	24 to 48	14 to 28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.
Gray Snow Mold (<i>Typhula incarnata</i>)	1.1	48	14 to 28	Make 2 applications, 14 to 28 days apart in late fall just prior to snow cover. For optimum control before extended periods of snow cover, make 1 or 2 applications of Honor at 0.83 to 1.1 oz per 1000 sq ft tank mixed with another effective fungicide from a different target site group, such as Iprodione Pro or Curalan® EG fungicide .
Large Patch (brown patch of warm season turfgrasses) (<i>Rhizoctonia solani</i>)	1.1	24 to 48	14 to 28	Apply prior to or directly at initial signs of infection in fall and make at least 2 sequential applications until turfgrass goes into dormancy. Reapplication in spring at time of green-up can be made if necessary. For control of brown patch of St Augustinegrass, centipedegrass, kikuyugrass, seashore paspalum and zoysiagrass (aka Zoysia patch).
Leaf Spot (<i>Bipolaris</i> , <i>Drechslera</i> , and <i>Exserohilum</i> spp.)	0.55 to 1.1	24 to 48	14 to 28	Apply when conditions are favorable for disease development.
Melting Out (<i>Drechslera poae</i>)	0.55 to 1.1	24 to 48	14 to 28	Apply when conditions are favorable for disease development.

10/12

Table 1. - Application Rates and Intervals for Honor® fungicide on Golf Course Turfgrass (continued)

Disease (Pathogen)	Use Rate (oz Product per 1000 sq ft)	Use Rate (oz Product per Acre)	Application Interval (days)	Comments
Necrotic Ringspot (<i>Leptosphaeria korrae</i>)	1.1	48	14 to 28	Aids in control of necrotic ringspot when combined with a non-strobilurin fungicide such as thiophanate methyl or propiconazole or chlorothalonil. Make applications in spring, fall or winter when conditions are present for outbreaks.
Pink Patch (<i>Limonomyces roseipellis</i>)	0.55 to 1.1	24 to 48	14 to 28	Apply when conditions are favorable for disease development.
Pink Snow Mold (<i>Microdochium nivale</i>)	1.1	48	14 to 28	Make 2 applications, 14 to 28 days apart in late fall just prior to snow cover. For optimum control before extended periods of snow cover, make 1 or 2 applications of Honor at 0.83 to 1.1 oz per 1000 sq ft tank mixed with another effective fungicide from a different target site group, such as Iprodione Pro or Curalan® EG fungicide .
Powdery Mildew (<i>Blumeria graminis</i>)	0.55 to 1.1	24 to 48	14 to 28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.
Pythium Blight ¹ (<i>Pythium aphanidermatum</i> , <i>Pythium</i> spp.)	1.1	48	10 to 14	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development. Tank mix Honor with another non-strobilurin fungicide labeled for Pythium blight control during severe disease pressure or when symptoms are already present.
Pythium Root Dysfunction ¹ (<i>Pythium volutum</i> , <i>Pythium</i> spp.)	1.1	48	14 to 28	Apply preventatively or early curative for control. Following two sequential applications rotate to other effective fungicides for this disease prior to additional Honor applications. Irrigate immediately following application.
Rapid Blight (<i>Labyrinthula terrestris</i>)	0.55 to 1.1	24 to 48	14 to 28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development. Follow the shorter spray interval when using the lower application rate.
Red Thread (<i>Laetisaria fuciformis</i>)	0.55 to 1.1	24 to 48	14 to 28	Apply when conditions are favorable for disease development.
Rhizoctonia Leaf or Sheath Spot (<i>R. oryzae</i> , <i>R. zea</i>)	1.1	24 to 48	14 to 28	Rhizoctonia infection can occur under warm humid conditions on both cool and warm season turfgrasses. This disease has been associated with localized dry spots and necrotic (brown) ring symptoms can form. Apply when conditions are favorable for disease development. Use of a soil wetting agent may be appropriate.
Rust (<i>Puccinia</i> and <i>Uromyces</i> spp.)	0.55 to 1.1	24 to 48	14 to 28	Apply when conditions are favorable for disease development.

11/12

Table 1. - Application Rates and Intervals for Honor® fungicide on Golf Course Turfgrass (continued)

Disease (Pathogen)	Use Rate (oz Product per 1000 sq ft)	Use Rate (oz Product per Acre)	Application Interval (days)	Comments
Summer Patch (<i>Magnaporthe poae</i>)	1.1	24 to 48	14 to 28	Initiate applications in the spring when soil temperatures reach 60 to 65° F at a 2-inch soil depth, or as dictated by local recommendations.
Take-all Patch (<i>Gaeumannomyces graminis</i> var. <i>avenae</i>)	1.1	48	28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development. Make 2 applications, 28-days apart, in the fall, and 2 applications, 28-days apart, in the spring.
Yellow Tuft (Downy Mildew) (<i>Sclerophthora macrospora</i>)	0.55 to 1.1	24 to 48	14 to 28	Use preventively. Begin applications when conditions are favorable for fungal infection, prior to disease symptom development.

¹ **DO NOT** apply more than two (2) sequential applications of **Honor** for anthracnose, dollar spot, gray leaf spot or Pythium. For all other diseases, when anthracnose, dollar spot, gray leaf spot or Pythium are not present, **DO NOT** apply more than three (3) sequential applications of **Honor**. Then alternate to an effective non-strobilurin fungicide before reapplying **Honor**.

Table 2. - Dilution Table for Spray Solutions of Honor fungicide

Ounces of Honor per 100 gallons of spray solution:				
Honor Use Rate (oz per 1000 sq ft)	Spray Volume-1 gallon per 1000 sq ft	Spray Volume-2 gallons per 1000 sq ft	Spray Volume-3 gallons per 1000 sq ft	Spray Volume-4 gallons per 1000 sq ft
0.55	55	27.5	18.3	13.8
0.83	83	41.5	27.7	20.8
1.1	110	55.0	36.7	27.5

12/12

Conditions of Sale and Warranty

The **Directions For Use** of this product reflects the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. All such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Curalan and Honor are registered trademarks of BASF.

© 2006 BASF Corporation
All rights reserved.

007969-00xxx.20061219.NVA 2006-04-091-0292

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

