

7969-257

04/26/2007

Lisa Jones

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. T. R. Nelsen
Product Registration Manager
BASF Corporation, Agricultural Products
P.O. Box 13528
Research Triangle Park, NC 17709-3528

APR 26 2007

RE: Notification of Alternate Trade Name and Change in Label
EPA Registration Number: 7969-257
Date of Submission: March 28, 2007

Dear Mr. Nelsen:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated March 28, 2007, for the product BASF Corporation/ Triconazole SC Fungicide. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

STG

2/11

Print Form

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0080. Approval expires 2-28



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 7969-257	2. EPA Product Manager Ms. Mary Waller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) BASF Corporation/Triticonazole SC Fungicide	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) BASF Corporation, Agricultural Products PO Box 13528, Research Triangle Park, NC 27709-3528 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

APR 26 2007

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Alternate trade name and change in label text to reflect new ownership of the registration

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> Text <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 1, 2.5 and 5 gal.		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name T.R. Nelsen	Title Product Registration Manager	Telephone No. (Include Area Code) (919) 549-2974
---------------------	---------------------------------------	---

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

8. Date Application Received (Stamped)

2. Signature 	3. Title Product Registration Manager	8. Date Application Received (Stamped)
4. Typed Name T.R. Nelsen	5. Date 3/28/2007	

4/11

Trinity™ FUNGICIDE

NOTIFICATION

APR 26 2007

A Fungicide For The Prevention And Control Of Certain Diseases Of Residential And Commercial Turfgrass, Golf Courses And Sod Farms. Intended For Use By Commercial Applicators Only.

Active Ingredient:		
triticonazole*	19.2%
Other Ingredients:	80.8%
Total:	100.0%

* Contains 1.69 lbs triticonazole per US gallon.

EPA Reg. No. 7969-257

EPA Est. No. _____

**KEEP OUT OF REACH OF CHILDREN.
CAUTION/PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside booklet for complete **First Aid, Precautionary Statements, Directions For Use, and Conditions of Sale and Warranty.**

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net Contents: _____

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709



5/11

FIRST AID

If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • DO NOT give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Socks and shoes.
- Chemical-resistant gloves made of waterproof material.

Wash hands and exposed skin thoroughly after handling the concentrate and after application.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. **DO NOT** contaminate water when cleaning equipment or disposing of equipment washwaters.

IN CASE OF EMERGENCY:

In case of large-scale spillage regarding this product, call:
CHEMTREC 1-800-424-9300
BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment.
- Your local poison control center (hospital).
- BASF Corporation 1-800-832-HELP (4357).

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

6/11

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made out of any waterproof material
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or green houses.

Keep unprotected persons out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container. Keep container closed. Store in an area that is out of reach of children and animals, away from food, feedstuffs, fertilizers and seed. Protect from frost and freezing.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

TURF

GENERAL INFORMATION

TURF (Golf Courses, Commercial Turfgrass, Sod Farms and Residential Turf). Intended for use only by licensed commercial applicators.

Trinity™ fungicide is a foliar-applied fungicide recommended for turfgrass disease control on golf courses, commercial turfgrass, sod farms and residential turf where fine turf is grown. Intended for use only by licensed commercial applicators.

When used in conjunction with good turf management practices, **Trinity** is effective in controlling the following diseases:

- | | |
|-------------------|----------------|
| Anthracnose | Red Leafspot |
| Brown Patch | Red Thread |
| Dollar Spot | Rust |
| Gray Snow Mold | Summer Patch |
| Large Brown Patch | Take-All Patch |
| Pink Snow Mold | |

Trinity is also effective in controlling the following diseases when combined with other products:

- Pythium Blight
- Summer Stress Complex/Decline

Begin applications when local environmental conditions favor disease development unless otherwise noted. Apply with sufficient water to obtain adequate coverage (1 to 5 gallons of spray solution per 1000 sq ft).

Recommended Turf Grass Species

- Bentgrass (Creeping, Colonial)
- Fine Fescue
- Kentucky Bluegrass
- Perennial Ryegrass
- Poa Annua*
- Tall Fescue
- Zoysia

Trinity on Turf Treated with Plant Growth Regulators

Trinity has been tested and has shown no adverse effects when applied to recommended turf types with the following Plant Growth Regulators (PGRs):

PGR	Acceptable Turf Types
Chipco® Proxy® (Ethephon)	Bentgrass Kentucky Bluegrass Perennial Rye Tall Fescue
Primo® (Trinexapac-ethyl)	Bentgrass Kentucky Bluegrass Perennial Rye Tall Fescue
Embark® (Mefluidide)	Bentgrass Tall Fescue

7/11

APPLICATION DIRECTIONS

DISEASE	USE DIRECTIONS	RATE fl oz product per 1000 sq ft		NOTES
		Preventative	Curative	
Dollar Spot <i>(Lanzia spp. and Moellerodiscus spp.)</i>	Repeat applications at 14 to 28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure, use the lower curative rate and shorter interval. Under severe disease conditions, use the higher rate and shorter interval.	0.5 to 0.75 fl oz	0.75 to 1.0 fl oz	Trinity™ fungicide is compatible with ProStar®, Compass®, and Heritage®. For control of dollar spot, combine Trinity at labeled rates with ProStar, Compass or Heritage at labeled rates.
Brown Patch <i>(Rhizoctonia solani)</i>	Repeat applications at 14 to 28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under active disease conditions, use the curative rate and shorter interval.	0.5 to 1.0 fl oz	1.0 to 2.0* fl oz *2.0 fl oz rate may be applied if needed in transition areas of the South under heavy disease pressure.	In conditions favoring moderate to heavy disease pressure, Trinity can be combined with other fungicides labeled for brown patch including ProStar or 26GT® at labeled rates.
Summer Patch <i>(Magnaporthe poae)</i>	Begin applications in spring (April to May) and repeat at 14 to 28 day intervals per the following situation: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.	1.0 to 2.0 fl oz		
Anthracnose <i>(Colletotrichum sp.)</i>	Repeat applications at 14 to 28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure, use the lower curative rate and shorter interval. Under severe disease conditions, use the higher rate and shorter interval.	0.5 to 0.75 fl oz	0.75 to 1.0 fl oz	

8/11

APPLICATION DIRECTIONS *(continued)*

DISEASE	USE DIRECTIONS	RATE fl oz product per 1000 sq ft		NOTES
		Preventative	Curative	
Take-All Patch <i>(Gaeumannomyces graminis)</i>	Make 1 to 2 fall applications (September to October) and 1 to 2 spring applications (April to May) depending on local conditions. Repeat applications at 14 to 28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under active disease conditions, use the curative rate and shorter interval.	0.5 to 1.0 fl oz	1.0 fl oz	
Large Brown Patch <i>(Rhizoctonia solani)</i>	Repeat applications at 14 to 28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under active disease conditions, use the curative rate and shorter interval.	0.5 to 1.0 fl oz	1.0 fl oz	
Red Leafspot <i>(Drechslera erythrospila)</i>	Repeat applications at 14 to 28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure, use the lower curative rate and shorter interval. Under severe disease conditions, use the higher rate and shorter interval.	0.5 to 0.75 fl oz	0.75 to 1.0 fl oz	
Algae	For algae suppression , repeat applications at 14 to 28 day intervals per the following situation: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.	0.5 to 1.0 fl oz		

9/11

APPLICATION DIRECTIONS (continued)

DISEASE	USE DIRECTIONS	RATE fl oz product per 1000 sq ft		NOTES
		Preventative	Curative	
Rust (<i>Puccinia</i> spp.)	Repeat applications at 14 to 28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure, use the lower curative rate and shorter interval. Under severe disease conditions, use the higher rate and shorter interval.	0.5 to 0.75 fl oz	0.75 to 1.0 fl oz	
Gray Snow Mold (<i>Typhula</i> spp.)	Make 1 to 2 applications late in the fall before snow cover occurs. Repeat applications at 14 to 28 day intervals per the following situation: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.	0.5 to 1.0 fl oz		Trinity™ fungicide can be combined with the following fungicides for greater efficacy: ProStar®, Chlorothalonil, PCNB, Chipco® Signature™ or 26GT®. Use only at labeled rates.
Pink Snow Mold (<i>Microdochium</i>)	Make 1 to 2 applications late in the fall before snow cover occurs. Repeat applications at 14 to 28 day intervals per the following situation: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.	0.5 to 1.0 fl oz	1.0 fl oz	Trinity can be combined with the following fungicides for greater efficacy: Chlorothalonil, PCNB, Chipco Signature or 26GT. Use only at labeled rates.
Red Thread	Repeat applications at 14 to 28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure, use the lower curative rate and shorter interval. Under severe disease conditions, use the higher rate and shorter interval.	0.5 to 0.75 fl oz	0.75 to 1.0 fl oz	
DO NOT apply more than 6 fluid ounces per 1000 sq ft of Trinity annually.				

10/11

Additional Tank Mixtures	
DISEASE	USE DIRECTIONS
Pythium Blight	Combine Trinity™ fungicide with either Chipco® Signature™ or Banol® brand fungicides for control of Pythium blight. Apply at labeled rates.
Summer Stress Complex/Summer Decline	Combine Trinity with Chipco Signature for diseases associated with summer decline. Apply at labeled rates.
DO NOT apply more than 6 fluid ounces per 1000 sq ft of Trinity annually.	

MIXING RATES

Trinity at 0.5 fluid ounce per 1000 sq ft						
SPRAY VOLUME (gallons per 1000 sq ft)						
Spray Tank Capacity		1 gallons	2 gallons	3 gallons	4 gallons	5 gallons
	25 gallons	12.5 fl oz	6.3 fl oz	4.2 fl oz	3.1 fl oz	2.5 fl oz
	50 gallons	25 fl oz	12.5 fl oz	8.3 fl oz	6.3 fl oz	5 fl oz
	100 gallons	50 fl oz	25 fl oz	16.7 fl oz	12.5 fl oz	10 fl oz
	200 gallons	100 fl oz	50 fl oz	33.3 fl oz	25 fl oz	20 fl oz

Trinity at 0.75 fluid ounce per 1000 sq ft						
SPRAY VOLUME (gallons per 1000 sq ft)						
Spray Tank Capacity		1 gallons	2 gallons	3 gallons	4 gallons	5 gallons
	25 gallons	18.8 fl oz	9.4 fl oz	6.3 fl oz	4.7 fl oz	3.75 fl oz
	50 gallons	37.5 fl oz	18.8 fl oz	12.5 fl oz	9.4 fl oz	7.5 fl oz
	100 gallons	75 fl oz	37.5 fl oz	25 fl oz	18.8 fl oz	15 fl oz
	200 gallons	150 fl oz	75 fl oz	50 fl oz	37.5 fl oz	30 fl oz

Trinity at 1.0 fluid ounce per 1000 sq ft						
SPRAY VOLUME (gallons per 1000 sq ft)						
Spray Tank Capacity		1 gallons	2 gallons	3 gallons	4 gallons	5 gallons
	25 gallons	25 fl oz	12.5 fl oz	8.3 fl oz	6.3 fl oz	5 fl oz
	50 gallons	50 fl oz	25 fl oz	16.7 fl oz	12.5 fl oz	10 fl oz
	100 gallons	100 fl oz	50 fl oz	33.3 fl oz	25 fl oz	20 fl oz
	200 gallons	200 fl oz	100 fl oz	66.7 fl oz	50 fl oz	40 fl oz

NOTES

- Apply with sufficient water to obtain adequate coverage (1 to 5 gallons of spray solution per 1000 sq ft).
- Apply with a properly calibrated sprayer and maintain agitation during spray operations.

RESTRICTIONS AND LIMITATIONS

- Only for use on labeled grasses.
- For golf courses only, **DO NOT** apply to turf cut higher than 1" on golf holes where water bodies are present.
- This product is compatible with most commonly used fungicides; however, check compatibility before tank mixing. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against mixing.
- Mixture with stickers, extenders, or wetting agents is not necessary.
- **DO NOT** graze animals on treated turf.
- **DO NOT** feed clippings from treated turf to livestock or poultry.
- **DO NOT** apply **Trinity** through any type of irrigation system.

11111

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Trinity is a trademark of BASF.

26GT, Banol, Chipco, Compass, ProStar and Proxy are registered trademarks and **Signature** is a trademark of Bayer.

Embark is a registered trademark of PBI/GORDON Corporation.

Heritage and **Primo** are registered trademarks of a Syngenta Group Company.

© 2007 BASF Corporation
All rights reserved.

007969-00257.20070306.NVA 2007-04-302-0029

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

